

Supplementary Table S2. Primers used for qRT-PCR analyses.

Gene	Forward (5'-3')	Reverse (5'-3')
<i>HGF</i>	ACGCTACGAAGTCTGTGA	AAGAATTTGTGCCGGTGT
<i>TNC</i>	TCTTGAAGGCAGGCGCAAAC	CCAAATGCCCAGGTGTGGACCGAT
<i>VIM</i>	ATTCCACTTTGCGTTCAAGG	CTTCAGAGAGAGGAAGCCGA
<i>KRT18</i>	AAATCCGGGAGCACTTGGAG	CAATCTGCAGAACGATGCGG
<i>EPCAM</i>	GTGCTGGTGTGTGAACACTG	GAAGTGCAGTCCGCAAACCTT
<i>SOX2</i>	AGCTACAGCATGATGCAGGA	GGTCATGGAGTTGTACTGCA
<i>ZO1</i>	CAGAACCAAAGCCTGTGTATG	TTAGGTAGGACACCATCAGATGGA
<i>TWIST1</i>	GTCATGGCCAACGTGCGGGA	GCCGCCAGCTTGAGGGTCTG
<i>MMP1</i>	CAGAGATGAAGTCCGGTTTTTC	GGGGTATCCGTGTAGCACAT
<i>MMP3</i>	AAATCCTGATCTTTAAAGACAGGCACT	GAGATCAAATGCAATTCAGGTTCA
<i>COL6A3</i>	GAGCAGCTTGACAACATTGCC	GCCCAGAGCACTTGCAGG
<i>COL4A1</i>	CCGGATCACATTGACATGAAACC	TGGAAACCAGTCCATGCTCG

<i>c-MET</i>	TGAGAAGGCTAAAGGAAA	TCAAAGGCATGGACATAC
<i>VAV3</i>	ACCAATGGACTGCGAAGAAC	GCTTGACTGCATCACTTGGA
<i>GAPDH</i>	TGACGCTGGGGCTGGCATT	GGCTGGTGGTCCAGGGGT