Supporting information for


Synthesis and investigation of pinane-based chiral tridentate ligands in the asymmetric addition of diethylzinc to aldehydes


Mounir Raji¹, Tam Minh Le^{1,2}, Ferenc Fülöp^{1,2} and Zsolt Szakonyi^{1,3,*}

- ¹ Institute of Pharmaceutical Chemistry, University of Szeged, Interdisciplinary excellent center, H-6720 Szeged, Eötvös utca 6, Hungary; raji.mounir@pharm.u-szeged.hu (M.R); leminhtam@pharm.u-szeged.hu (T.M.L.)
- ² MTA-SZTE Stereochemistry Research Group, Hungarian Academy of Sciences, H-6720 Szeged, Eötvös utca 6, Hungary
- ³ Interdisciplinary Centre of Natural Products, University of Szeged, H-6720 Szeged, Eötvös utca 6, Hungary
- * Correspondence: szakonyi@pharm.u-szeged.hu; Tel.: +36-62-546809; Fax: +36-62-545705


Contents


¹ H, ¹³ C, HSQC, HMBC and NOESY	NMR spectra of new compounds	S3 – S83








¹³C-NMR for compound **4a**











































¹³C-NMR for compound **8**















¹H-NMR for compound **9**














COSY for compound 9


































S36




















S44

