

Supplementary Material

The Effect of Hexanoyl Glycol Chitosan on the Proliferation of Human Mesenchymal Stem Cells

Young-Hoon Jeong ^{1,†}, Hye Min Oh ^{2,†}, Man Ryul Lee ^{3,†}, C-Yoon Kim ¹, Chanyang Joo ², Soon-Jung Park ¹, Yun-Ho Song ¹, Changhee Kang ¹, Hyung-Min Chung ¹, Sun-Woong Kang ^{4,5,*}, Kang Moo Huh ^{2,*}, and Sung-Hwan Moon ^{6,*}

¹ Department of Stem Cell Biology, School of Medicine, Konkuk University, Seoul 05029, Korea; vivavets@gmail.com (C.-Y.K.); parksoonjung@gmail.com (S.-J.P.); Yoontech67@gmail.com (Y.-H.S.); changhee_k@naver.com (C.K.); stemchung@gmail.com (H.-M.C.)

² Department of Polymer Science and Engineering, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon 34134, Korea; Jooyang011@naver.com (C.J.); khuh@cnu.ac.kr (K.M.H.)

³ Soonchunhyang Institute of Medi-bio Science (SIMS), College of Medicine, Soon Chun Hyang University, Cheonan 31151, Korea

⁴ Predictive Model Research Center, Korea Institute of Toxicology, 141 Gajeong-ro, Yuseong-gu, Daejeon 34114, Korea; swkang@kitox.re.kr

⁵ Department of Human and Environmental Toxicology, University of Science and Technology, Daejeon 34114, Korea

⁶ Department of Medicine, School of Medicine, Konkuk University, Seoul 05029, Korea; sunghwanmoon@kku.ac.kr

* Correspondence: swkang@kitox.re.kr (S.-W.K.); khuh@cnu.ac.kr (K.M.H.); sunghwanmoon@kku.ac.kr (S.-H.M.); Tel.: +82-42-610-8209 (S.-W.K.); Tel.: +82-42-821-8910 (K.M.H.); Tel.: +82-2-2049-6028 (S.-H.M.)

† These authors contributed equally to this work as first author.

‡ These authors contributed equally to this work as corresponding authors.




Figure S1. Additional cell cycle analysis of 1 ug/mL hexanoyl glycol chitosan (HGC) only groups at passage 10.