

Supplementary Information

Upper critical solution temperature (UCST) behavior of coacervate cationic protamine and anionic macroions

Hyunbin Kim¹, Byoung-jin Jeon², Sangsik Kim^{1,3}, YongSeok Jho^{4,*} and Dong Soo Hwang^{1,3,*}

¹ Division of Integrative Biosciences and Biotechnology, Pohang University of Science and Technology (POSTECH), Pohang 37673, Republic of Korea ; hbkim22@postech.ac.kr

² Materials Department, University of California, Santa Barbara, Santa Barbara, CA 93106 USA ; saleh@ucsb.edu

³ Division of Environmental Science and Engineering, Pohang University of Science and Technology (POSTECH), Pohang 37673, Republic of Korea ; ukssloveu@postech.ac.kr

⁴ Department of Physics and Research Institute of Natural Science, Gyeongsang National University, Jinju 52828, Republic of Korea ; ysjho@gnu.ac.kr

* Correspondence: yongseokjho@gmail.com or dshwang@postech.ac.kr; Tel.: +82-54-279-9505

Received: date; Accepted: date; Published: date




Figure S1. Turbidity of protamine/sodium chloride solution.




Figure S2. Measurement of τ with variable radius of droplets for interfacial tension of the complex coacervate.

FOV	R (um)	t(msec)
1	3.1570	29.74
2	miss	miss
3	2.6923	21.86
4	2.7886	18.43
5	2.9603	35.15
6	4.1624	33.63
7	2.6896	22.51
8	3.2550	28.24
9	3.8314	30.44
10	4.1602	36.63
11	5.5445	46.23
12	4.1438	34.15

Table S1. Measurement of τ with variable radius of droplets for interfacial tension of the complex coacervate.