

Catalyst Speciation During ansa-Zirconocene-Catalyzed Polymerization of 1-Hexene Studied by UV-vis Spectroscopy – Formation and Partial Re- Activation of Zr-Allyl Intermediates.

Valentina N. Panchenko ^{1,2}, Dmitrii E. Babushkin ¹, John E. Bercaw ^{3 *}and Hans H. Brintzinger ⁴

¹ Boreskov Institute of Catalysis, Russian Academy of Sciences, Siberian Branch, RU-630090 Novosibirsk, Russian Federation; panchenko@catalysis.ru, dimi@catalysis.ru

² Novosibirsk State University, Pirogova Str. 2, 630090, Novosibirsk, Russian Federation

³ Arnold and Mabel Beckman Laboratories of Chemical Synthesis, California Institute of Technology, Pasadena, California 91125, USA; bercaw@caltech.edu

⁴ Fachbereich Chemie, Universität Konstanz D-78464 Konstanz, Germany; hans.brintzinger@uni-konstanz.de

* Correspondence: hans.brintzinger@uni-konstanz.de

Supporting Information

Contents:

¹ H NMR signals ascribed to species C-2	S-1
¹ H-NMR signals of CH ₃ D	S-2
UV-vis spectrum of SBIZr-CH ₂ SiMe ₃ ⁺	S-3
UV-vis spectra of SBIZr(μ-Me) ₂ AlMe ₂ ⁺ at +40°C and -20°C	S-4
Plot of ln[C-0] _t vs. t	S-5
Plot of ln[C-1] _t vs. t	S-6
Effects of initial [Al]/[Zr] ratio	S-7





Figure S-1. ¹H NMR signals arising during polymerization of 1-hexene by the catalyst system described in Figure 1. First 4 traces (2.0 , 3.6, 5.2 and 6.7 min mean time). Signals at 6.36, 5.67, 4.87 and 4.48 ppm (marked by *), possibly due to polymer-carrying cations of type SBIZr- π -(1-R-2-pol-C₃H₃)⁺ (**10**).



Additional traces after ca. 20 min.




Figure S-2. ¹H NMR spectra of CH_4 and CH_3D in solutions obtained by addition of an equivalent amount of trityl perfluorotetraphenyl borate to a solution of SBIZrMe_2 in toluene-d₈, as described in Figure 3, after reaction times of ca. 4 h (bottom) and ca. 10 h (top).




Figure S-3. UV-vis spectra of a 0.8 mM solution of SBIzr(Me)-CH₂SiMe₃ after addition of ca. 0.8 equiv. of trityl perfluorotetraphenyl borate.




Figure S-4. UV-vis spectra of a ca. 0.5 mM toluene solution of $(\text{SBI})\text{Zr}(\mu\text{-Me})_2\text{AlMe}_2^+$ $\text{B}(\text{C}_6\text{F}_5)_4^-$ at $+40^\circ\text{C}$ (broken line) and at -20°C (solid line).




Figure S-5. Plot of $\ln[C-0]_t$ vs. t for reaction stage 1 ($t < 120$ s):




Figure S-6. Plot of $\ln[C-1]_t$ vs. t for $300 \text{ s} < t < 600 \text{ s}$




Figure S-7. Effects of initial $[Al]/[Zr]$ ratio on the re-conversion of **C-2** ($\lambda_{max} = 560$ nm) to **C-0** ($\lambda_{max} = 495$ nm), conditions as described in Figure 1.