

## Supporting Information

# Synthetic Polypeptides with Cationic Arginine Moieties Showing High Antimicrobial Activity in Similar Mineral Environments to Blood Plasma

Kuen Hee Eom <sup>1</sup>, Shuwei Li <sup>1</sup>, Eun Gyeong Lee <sup>1</sup>, Jae Ho Kim <sup>2</sup>, Jung Rae Kim <sup>1</sup> and Il Kim <sup>1,\*</sup>

<sup>1</sup> School of Chemical Engineering, Pusan National University, Busandaehag-ro 63-2, Geumjeong-gu, Busan 46241, Korea; reimreim@pusan.ac.kr (K.H.E.); lishuwei0325@pusan.ac.kr (S.L.); rud6063@pusan.ac.kr (E.G.L.); j.kim@pusan.ac.kr (J.R.K.)

<sup>2</sup> Department of Physiology, School of Medicine, Pusan National University, Busandaehak-ro, Mugeum-eup, Yangsan-si 50612, Korea; jhkimst@pusan.ac.kr

\* Correspondence: ilkim@pusan.ac.kr

## Table of Contents

**Figure S1** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>43</sub>]<sub>3</sub> (Entry 1 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S2** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>31</sub>-*co*-(Phe)<sub>8</sub>]<sub>3</sub> (Entry 2 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S3** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>25</sub>-*co*-(Phe)<sub>15</sub>]<sub>3</sub> (Entry 3 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S4** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>31</sub>]<sub>3</sub> (Entry 4 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S5** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>14</sub>-*co*-(Phe)<sub>7</sub>]<sub>3</sub> (Entry 5 in Table 1) in TFA.

**Figure S6** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>19</sub>-*co*-(Phe)<sub>11</sub>]<sub>3</sub> (Entry 6 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S7** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>16</sub>-*co*-(Phe)<sub>4</sub>]<sub>3</sub> (Entry 8 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S8** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>13</sub>-*co*-(Phe)<sub>9</sub>]<sub>3</sub> (Entry 9 in Table 1) in DMSO-*d*<sub>6</sub>.

**Figure S9** <sup>1</sup>H NMR spectra of *mb*-p(Cbz-Lys)<sub>24</sub> (Entry 11 in Table 1) in TFA.


**Figure S10** FTIR spectra of *s*-p(Lys)<sub>21</sub>, *s*-p(Arg)<sub>21</sub>, *l*-p[(Lys)<sub>23</sub>-*co*-(Phe)<sub>7</sub>], and *l*-p[(Arg)<sub>23</sub>-*co*-(Phe)<sub>7</sub>].

**Figure S11** Determination of minimal inhibitory concentration (MIC) on *E. coli* cells with *l*-, *s*- and *mb*-polypeptides.


**Figure S12** Determination of minimal inhibitory concentration (MIC) on *B. substillis* cells with *l*-, *s*- and *mb*-polypeptides.

**Figure S13** Hemolytic activity of *mb*-p[(Arg)<sub>23</sub>-*co*-(Phe)<sub>7</sub>]<sub>8</sub> in fresh red blood cells of mice.


**Supplementary figures**




**Figure S1** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>43</sub>]<sub>3</sub> (Entry 1 in Table 1) in DMSO-*d*<sub>6</sub>.




**Figure S2** <sup>1</sup>H NMR spectra of *s*-p[(Cbz-Lys)<sub>31</sub>-co-(Phe)<sub>8</sub>]<sub>3</sub> (Entry 2 in Table 1) in DMSO-*d*<sub>6</sub>.




**Figure S3**  $^1\text{H}$  NMR spectra of  $s\text{-}p[(\text{Cbz-Lys})_{25}\text{-}co\text{-}(\text{Phe})_{15}]_3$  (Entry 3 in Table 1) in  $\text{DMSO-}d_6$ .




**Figure S4**  $^1\text{H}$  NMR spectra of  $s\text{-p}[(\text{Cbz-Lys})_{31}]_3$  (Entry 4 in Table 1) in  $\text{DMSO-}d_6$ .




**Figure S5**  $^1\text{H}$  NMR spectra of *s-p*[(Cbz-Lys)<sub>14</sub>-*co*-(Phe)<sub>7</sub>]<sub>3</sub> (Entry 5 in Table 1) in TFA.




**Figure S6**  $^1\text{H}$  NMR spectra of *s-p*[(Cbz-Lys)<sub>19</sub>-*co*-(Phe)<sub>11</sub>]<sub>3</sub> (Entry 6 in Table 1) in DMSO-*d*<sub>6</sub>.




**Figure S7**  $^1\text{H}$  NMR spectra of  $s\text{-}p[(\text{Cbz-Lys})_{16}\text{-}co\text{-}(\text{Phe})_4]_3$  (Entry 8 in Table 1) in  $\text{DMSO}-d_6$ .




**Figure S8**  $^1\text{H}$  NMR spectra of  $s\text{-}p[(\text{Cbz-Lys})_{13}\text{-}co\text{-}(\text{Phe})_9]_3$  (Entry 9 in Table 1) in  $\text{DMSO}-d_6$ .




**Figure S9**  $^1\text{H}$  NMR spectra of *mb*-p(Cbz-Lys)<sub>24</sub> (Entry 11 in Table 1) in TFA.




**Figure S10.** FTIR spectra of *s*-p(Lys)<sub>21</sub>, *s*-p(Arg)<sub>21</sub>, *l*-p[(Lys)<sub>23</sub>-*co*-(Phe)<sub>7</sub>], and *l*-p[(Arg)<sub>23</sub>-*co*-(Phe)<sub>7</sub>].



**Figure S11.** Determination of minimal inhibitory concentration (MIC) on *E. coli* cells with *l*-, *s*- and *mb*-polypeptides.



**Figure S12.** Determination of minimal inhibitory concentration (MIC) on *B. subtilis* cells with *l*-, *s*- and *mb*-polypeptides.



**Figure S13.** Hemolytic activity of *mb*-p[ $(\text{Arg})_{23}\text{-}co\text{-}(\text{Phe})_7$ ]<sub>8</sub> in fresh red blood cells of mice.