

Supplementary Materials: Synthesis of Highly Branched Polyolefins Using Phenyl Substituted α -Diimine Ni(II) Catalysts

Fuzhou Wang, Ryo Tanaka, Zhengguo Cai, Yuushou Nakayama and Takeshi Shiono

1. X-ray Structure Determinations

The molecular structures of complexes **C1** and **C5'** were confirmed by single-crystal X-ray diffraction and the corresponding ORTEP diagrams are shown in Figure S1. Crystal data, data collection, and refinement parameters are listed in Table S1. Selected bond distances and angles are summarized in Table S2.





Figure S1. Molecular structures of the complexes **C1** and **C5'**, where H atoms are omitted for clarity.

Table S1. Crystal data and structure refinements of complex **C1**.

Identification Code	C1
Empirical formula	C ₄₀ H ₃₂ Br ₂ N ₂ Ni
Formula mass	759.21
Temperature (K)	292
Wavelength (Å)	0.7107
Crystal size (mm ³)	0.33 × 0.31 × 0.28
Crystal system	Orthorhombic
Space group	<i>Cmcm</i>
<i>a</i> (Å)	11.7266 (8)
<i>b</i> (Å)	11.5635 (5)
<i>c</i> (Å)	27.5587 (14)
<i>V</i> (Å ³)	3,737.0 (4)
<i>Z</i>	8
Density (calcd.) (mg/cm ³)	2.699
Absorption coefficient (mm ⁻¹)	5.37
<i>F</i> (000)	3,072
Theta range for data collec. (°)	3.0–28.5
Limiting indices	-15 ≤ <i>h</i> ≤ 15, -14 ≤ <i>k</i> ≤ 15, -35 ≤ <i>l</i> ≤ 34
Reflections collected	9,590
Independent reflections	1,900
<i>R</i> _{int}	0.056
Final <i>R</i> indices [<i>I</i> > 2σ(<i>I</i>)]	<i>R</i> ₁ = 0.0640, <i>wR</i> ₂ = 0.1827
<i>R</i> indices (all data)	<i>R</i> ₁ = 0.0945, <i>wR</i> ₂ = 0.1986
Refinement method	Full-matrix least-squares on <i>F</i> ²
Goodness-of-fit on <i>F</i> ²	1.137
Max. and min. transmission	1.0000 and 0.56405
Largest diff. peak and hole (e·Å ⁻³)	0.482 and -0.301

Table S2. Selected bond lengths (\AA) and angles ($^\circ$) for complex C1.


Bond Lengths	(\AA)	Bond Angles	($^\circ$)
Br1–Ni1	2.3343(12)	Br1–Ni1–Br1i	117.34(8)
Ni1–Br1i	2.3343(12)	N1ii–Ni1–Br1i	112.96(6)
Ni1–N1ii	2.041(5)	N1–Ni1–Br1i	112.96(6)
Ni1–N1	2.041(5)	N1–Ni1–Br1	112.96(6)
N1–C8	1.427(8)	N1ii–Ni1–Br1	112.96(6)
N1–C9	1.290(9)	N1ii–Ni1–N1	82.8(3)
C1–C2i	1.308(14)	C8–N1–Ni1	130.7(4)

**Figure S2.** Cell structure of C1 parallel to the bis(imino)acenaphthene unit.


2. Experimental Sections

2.1. Synthesis and Characterizations of Complexes C1–C7

The synthesis of ligands **L**, **L1–L7** and their complexes **Pd-Cat.**, **C1–C7** are outlined in Scheme S1. After the protection of the amino group by acetic acid, the aniline derivatives were brominated. The Suzuki-coupling reaction of the bromoaniline derivatives and phenylboronic acid obtained with a Pd(II) catalyst (**Pd-Cat.**) in PEG-400/H₂O led to the corresponding phenyl-substituted aniline derivatives **1–6**. The ligands **L1–L7** were prepared by the condensation of two equivalents of the appropriate aniline with one equivalent of acenaphthoquinone, in the presence of a formic acid catalyst. The reaction of equimolar amounts of NiBr₂(DME) and the α -diimine ligands **L1–L7** in CH₂Cl₂ led to the displacement of 1,2-dimethoxyethane and afforded the Ni(II) complexes **C1–C7** as a moderately air-stable deep red microcrystalline solid in good yields, respectively. The corresponding palladium dichloride complex (**Pd-Cat.**) is accessible by the reaction of the *ortho*-*sec*-phenethyl substituted chiral ligand **L** with (CH₃CN)₂PdCl₂ in CH₂Cl₂ at ambient temperature. Compounds **L1–L7** were well characterized by IR, ¹H NMR, ¹³C NMR spectroscopy and elemental analysis.

**Scheme S1.** Synthesis of α -diimine ligands **L**, **L1-L7** and their complexes **Pd-Cat.**, **C1-C7**.

The new organic compounds were well characterized by ^1H NMR and ^{13}C NMR (see below).

**Figure S3.** ^1H and ^{13}C NMR spectra of **1**.





Figure S4. ¹H and ¹³C NMR spectra of **2**.





Figure S5. ¹H and ¹³C NMR spectra of **3**.




Figure S6. ¹H and ¹³C NMR spectra of ligand L1.




Figure S7. ¹H and ¹³C NMR spectra of ligand L2.




Figure S8. ¹H and ¹³C NMR spectra of ligand L3.




Figure S9. ¹H and ¹³C NMR spectra of ligand L.

2.2. Living Polymerization of 1-Decene with C1-MMAO

Polymerization of 1-decene was investigated with C1-MMAO at 20 °C with the [Al]/[Ni] molar ratio of 300, and the results are summarized in Table S3.

Table S3. Polymerization of 1-decene with C1-MMAO. ^a

Entry	Time (min)	Conversion (%)	M_n ^b (kg·mol ⁻¹)	M_w/M_n ^b	N ^c (mol)
1	5	7.3	18.8	1.09	5.4
2	15	19.5	51.4	1.08	5.3
3	30	45.4	110.0	1.07	5.8
4	40	58.1	151.8	1.10	5.4
5	50	75.1	186.9	1.15	5.6
6	60	91.3	233.5	1.14	5.5

^a Polymerization conditions: Ni = 10 µmol; cocatalyst MMAO, [Al]/[Ni] = 300; 1-decene = 10 mmol; 20 °C; solvent = toluene, total volume = 20 mL; ^b M_n in kg/mol, determined by GPC against polystyrene standard. ^c Number of polymer chain calculated by yield and M_n .




Figure S10. Plots of M_n (■) and M_w/M_n (▲) as a function of conversion for the polymerization of 1-decene (20 °C, Table S3).

GPC traces for the poly(1-decene)s (Table S3):




Figure 11. Cont.




Figure S11. GPC traces for the poly(1-decene)s obtained with C1-MMAO at different reaction time (entries 1–5, Table S3).

3. Microstructure Analysis of the Polymers

3.1. Equations (S1)–(S3) for the Microstructure Analysis of the Polymers

3.1.1. Calculation of the Degree of Branching

The degree of branching (B) was estimated by ^1H NMR spectroscopy and was corrected for end groups as follows (S1):

$$B = \frac{2(I_{\text{CH}_3})}{3(I_{\text{CH}} + I_{\text{CH}_2} + I_{\text{CH}_3})} \times 1000 \quad (\text{S1})$$

Branching degree, the number of methyl carbon in every 1000 carbons, CH_3 , CH_2 , CH refer to the intensities of the methyl, methylene and methine resonances in ^1H NMR spectra.

3.1.2. Calculation of $\omega,1$ -Insertions

The fraction of $\omega,1$ -insertions was calculated using the following equation reported by Brookhart *et al.* [1].

$$\omega,1\% = \frac{1000 - (\omega - 2)B}{1000 + 2B} \times 100 \quad (\text{S2})$$

where B is the total branching calculated by Equation (S1), ω is the number of carbon atoms in the monomer.

Assignments of the ^{13}C NMR spectra and equations for the quantitative analysis of the polyolefins under investigation, according to Equation (S3) reported by Azoulay *et al.* [2]. Chemical shift and assignment of the peak listed in the Table S4 are marked in the spectrum (Figure S12).

$$\begin{aligned} \text{CH}_3 &= (I_2 + I_5)/2 + I_1 + I_3 + I_6 \\ \text{Methyl branch, Me} &= I_6 - I_4 + I_7; \text{ Ethyl branch, Et} = (I_1 + I_{10})/2 \\ \text{Propyl branch, Pr} &= (I_4 + I_{23})/2; \text{ Butyl branch, Bu} = I_3 - I_8 \\ \text{sec-Butyl branch, } {}^s\text{Bu} &= (I_2 + I_5)/2; \text{ Longer branch, Lg} = (I_8 + I_{17})/2 \end{aligned} \quad (\text{S3})$$




Figure S12. ^{13}C NMR spectrum of the polyethylene obtained with C1-MMAO at 20 °C (entry 11, Table 1).

Table S4. Chemical shift and assignment of peaks. ^a

Peak No.	Chemical shift (ppm)	Assignment	Peak No.	Chemical shift (ppm)	Assignment
1	11.09	1B ₂	14	29.61	4B _n
2	11.53	^s Bu	15	29.95	CH ₂
3	14.25	1B ₄ , 1B _n	16	30.36	$\gamma\text{B}_1, \gamma\text{B}_{2+}$
4	14.66	1B ₃	17	32.09	3B _n
5	19.33	^s Bu	18	32.57	3B ₅
6	19.81	1B ₁	19	32.94	brB ₁
7	19.98	2B ₃	20	33.34	αB_2
8	22.85	2B _n	21	33.91	αB_{3+}
9	23.24	2B ₄	22	34.56	nB _n
10	26.03	2B ₂	23	36.84	3B ₃
11	26.95	(n-1)B _n	24	37.33	αB_1
12	27.30	βB_{2+}	25	37.77	brB _n
13	27.66	βB_1	26	39.03	brB ₂

^a ^{13}C NMR (CDCl_3 δ: 77.16 ppm) measurements of the polyethylene are marked in Figure S12; Note on labels: for $x\text{B}_n$ B_n is a branch of length n carbons, x is the carbon being discussed, and the methyl at the end of the branch is numbered 1. Thus, the second carbon from the end of a butyl branch is 2B₄. $x\text{B}_{n+}$ refers to branches of length n and longer.

3.2. ^1H and ^{13}C NMR Spectra of the Polyethylenes (Tables 1 and 2)

NMR spectra of the polyethylenes prepared by C1–C7 (see Figures S13–S21):




Figure S13. ^1H NMR spectra of the polyethylenes obtained with C1-Et₂AlCl (entries 3 and 7–9, Table 1).




Figure S14. ^1H NMR spectra of the polyethylenes obtained with C1-MMAO (entries 10–14, Table 1).




Figure S15. ^1H NMR spectra of the polyethylenes obtained with C2–C7/MMAO (entries 15–20, Table 1).




Figure S16. ^{13}C NMR spectra of the polyethylene obtained with C1-Et₂AlCl at 40 °C (entry 7, Table 1).




Figure S17. ^{13}C NMR spectra of the polyethylene obtained with C1-Et₂AlCl at 20 °C (entry 3, Table 1).




Figure S18. ^{13}C NMR spectra of the polyethylene obtained with C1-MMAO at 0 °C (entry 10, Table 1).




Figure S19. ^{13}C NMR spectra of the polyethylene obtained with C1-MMAO at 40 °C (entry 12, Table 1).




Figure S20. ¹³C NMR spectra of the polyethylene obtained with C1-MMAO at 60 °C (entry 13, Table 1).




Figure S21. ¹³C NMR spectra of the polyethylene obtained with C1-MMAO at 80 °C (entry 14, Table 1).

3.3. ¹H and ¹³C NMR Spectra of the Poly(1-Alkene)s (Table 3)




Figure S22. ¹H and ¹³C NMR spectra of poly(1-hexene) obtained with C1-MMAO at 20 °C (entry 1, Table 4).




Figure S23. ¹H and ¹³C NMR spectra of poly(1-hexene) obtained with C1-MMAO at 20 °C (entry 2, Table 3).




Figure S24. ¹H and ¹³C NMR spectra of poly(1-octene) obtained with C1-MMAO at 20 °C (entry 3, Table 3).




Figure S25. ¹H and ¹³C NMR spectra of poly(1-octene) obtained with C1-MMAO at 20 °C (entry 4, Table 3).




Figure S26. ¹H and ¹³C NMR spectra of poly(1-decene) obtained with C1-MMAO at 20 °C (entry 5, Table 3).




Figure S27. ^1H and ^{13}C NMR spectra of poly(1-decene) obtained with C1-MMAO at $-20\text{ }^\circ\text{C}$ (entry 6, Table 3).




Figure S28. ^1H and ^{13}C NMR spectra of poly(1-decene) obtained with C1-MMAO at $0\text{ }^\circ\text{C}$ (entry 7, Table 3).




Figure S29. ¹H and ¹³C NMR spectra of poly(1-decene) obtained with C1-MMAO at 20 °C (entry 8, Table 3).




Figure S30. ¹H NMR spectra of poly(1-hexadecene)s obtained with C1-MMAO at 20 °C (entries 12–14, Table 3).




Figure S31. ^{13}C NMR spectra of poly(1-hexadecene) obtained with C1-MMAO at 0 °C (entry 13, Table 3).




Figure S32. ^{13}C NMR spectra of poly(1-hexadecene) obtained with C1-MMAO at 20 °C (entry 14, Table 3).

4. GPC Traces of the Polymers




Figure S33. GPC trace for the polyethylene obtained with C1-MMAO at 20 °C (entry 11, Table 1).




Figure S34. GPC trace for the polyethylene obtained with C1-MMAO at 40 °C (entry 12, Table 1).




Figure S35. GPC trace for the polyethylene obtained with C1-MMAO at 60 °C (entry 13, Table 1).




Figure S36. GPC trace for the polyethylene obtained with C1-MMAO at 80 °C (entry 14, Table 1).




Figure S37. GPC trace for the poly(1-hexene) obtained with C1-MMAO at 20 °C for 30 min (entry 1, Table 3).




Figure S38. GPC trace for the poly(1-hexene) obtained with C1-MMAO at 0 °C for 30 min (entry 2, Table 3).




Figure S39. GPC trace for the poly(1-octene) obtained with C1-MMAO at 20 °C for 30 min (entry 3, Table 3).




Figure S40. GPC trace for the poly(1-octene) obtained with C1-MMAO at 0 °C for 30 min (entry 4, Table 3).




Figure S41. GPC trace for the poly(1-decene) obtained with C1-MMAO at 20 °C for 30 min (entry 5, Table 3).




Figure S42. GPC trace for the poly(1-decene) obtained with C1-MMAO at 0 °C for 30 min (entry 7, Table 3).




Figure S43. GPC trace for the poly(1-decene) obtained with C1-MMAO at 20 °C for 30 min (entry 8, Table 3).




Figure S44. GPC trace for the poly(1-hexadecene) obtained with C1-MMAO at 20 °C for 30 min (entry 12, Table 3).




Figure S45. GPC trace for the poly(1-hexadecene) obtained with C1-MMAO at 0 °C for 30 min (entry 13, Table 3).




Figure S46. GPC trace for the poly(1-hexadecene) obtained with C1-MMAO at 20 °C for 30 min (entry 14, Table 3).

5. DSC Curves of the Polymers




Figure S47. DSC curves of polyethylenes obtained by C1-MMAO at different temperatures (entries 3, 7 and 13–14, Table 1).




Figure S48. DSC curve for the poly(1-hexene) obtained with C1-MMAO at 20 °C (entry 1, Table 3).




Figure S49. DSC curve for the poly(1-octene) obtained with C1-MMAO at 20 °C (entry 3, Table 3).




Figure S50. DSC curve for the poly(1-decene) obtained with C1-MMAO at 20 °C (entry 5, Table 3).




Figure S51. DSC curve for the poly(1-hexadecene) obtained with C1-MMAO at 20 °C (entry 12, Table 3).

References

- McCord, E.F.; McLain, S.J.; Nelson, L.T.J.; Ittel, S.D.; Tempel, D.; Killian, C.M.; Johnson, L.K.; Brookhart, M. ^{13}C NMR Analysis of α -olefin enchainment in poly(α -olefins) produced with nickel and palladium α -diimine catalysts. *Macromolecules* **2007**, *40*, 410–420.
- Azoulay, J.D.; Bazan, G.C.; Galland, G.B. Microstructural characterization of poly(1-hexene) obtained using a nickel α -Keto- β -diimine initiator. *Macromolecules* **2010**, *43*, 2794–2800.



© 2016 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).