

Table S1. Nitrogen content in rape seeds and straw, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Nitrogen - seeds (g kg ⁻¹ d.w.)			Nitrogen – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	33.77 ^{e-j} ±0.06	31.38 ^a ±0.34	33.31 ^{d-i} ±0.42	9.52 ^{ijk} ±0.10	10.43 ^{lm} ±0.11	9.72 ^{jk} ±0.12
S1Pd	34.33 ^k ±0.25	32.40 ^{bcd} ±0.35	33.10 ^{b-g} ±0.41	8.50 ^{def} ±0.09	6.99 ^b ±0.08	10.53 ^m ±0.13
S2Pd	34.25 ^{jk} ±0.18	32.29 ^{abc} ±0.35	34.42 ^{jk} ±0.43	8.91 ^{gh} ±0.10	14.48 ^q ±0.16	8.20 ^d ±0.10
S3Pd	34.90 ^{kl} ±0.22	32.19 ^{ab} ±0.35	37.15 ^{jk} ±0.43	9.31 ⁱ ±0.10	8.40 ^{de} ±0.09	12.05 ^o ±0.15
S1D	34.05 ^{g-k} ±0.27	32.80 ^{b-e} ±0.35	33.21 ^{c-h} ±0.41	9.41 ^{ij} ±0.10	9.82 ^{kl} ±0.11	10.53 ^m ±0.13
S2D	34.28 ^{jk} ±0.15	33.81 ^{f-j} ±0.37	32.70 ^{bcd} ±0.41	8.81 ^{fg} ±0.10	10.93 ⁿ ±0.12	7.30 ^c ±0.09
S3D	34.48 ^{jk} ±0.23	34.52 ^{jk} ±0.37	35.74 ^l ±0.45	9.31 ⁱ ±0.10	13.57 ^p ±0.15	17.94 ^s ±0.25
S1PdD	34.19 ^{h-k} ±0.06	32.40 ^{bcd} ±0.35	32.90 ^{b-f} ±0.41	10.12 ^{hl} ±0.11	7.19 ^{bc} ±0.08	15.06 ^r ±0.19
S2PdD	34.04 ^{g-k} ±0.05	34.12 ^{h-k} ±0.37	33.81 ^{f-j} ±0.42	9.21 ^{hi} ±0.10	6.28 ^a ±0.07	14.48 ^q ±0.18
S3PdD	34.71 ^{jk} ±0.18	33.91 ^{g-k} ±0.37	37.15 ^m ±0.46	9.21 ^{hi} ±0.11	8.71 ^{efg} ±0.09	11.24 ⁿ ±0.14
Mean	34.30^c±0.36	32.98^A±1.04	34.10^B±1.40	9.23^A±0.43	9.68^B±2.61	11.72^C±3.53

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization x years, capital letters for years. ± - standard deviation

Table S2. Sulphur content in rape seeds and straw, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Sulphur - seeds (g kg ⁻¹ d.w.)			Sulphur – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	2.31 ^a ±0.02	2.73 ^{fgh} ±0.03	2.65 ^{c-f} ±0.04	1.71 ^a ±0.02	2.02 ^c ±0.02	2.04 ^c ±0.09
S1Pd	2.71 ^{e-h} ±0.03	3.04 ⁱ ±0.03	2.73 ^{fgh} ±0.03	2.41 ^e ±0.03	3.14 ^k ±0.03	2.19 ^d ±0.03
S2Pd	2.56 ^{bc} ±0.03	2.84 ⁱ ±0.03	2.71 ^{e-h} ±0.03	2.64 ^{fg} ±0.03	3.14 ^k ±0.03	2.37 ^e ±0.03
S3Pd	2.61 ^{bcd} ±0.03	2.94 ^{jk} ±0.03	2.80 ^{hi} ±0.04	4.06 ^{ho} ±0.04	4.15 ^o ±0.04	2.03 ^c ±0.03
S1D	2.59 ^{bc} ±0.03	2.73 ^{fgh} ±0.03	2.79 ^{ghi} ±0.04	2.81 ^h ±0.03	2.18 ^d ±0.02	1.86 ^b ±0.02
S2D	2.70 ^{d-g} ±0.03	2.94 ^{jk} ±0.03	2.97 ^{kl} ±0.03	2.99 ⁱ ±0.03	3.34 ^{hl} ±0.04	3.47 ^m ±0.04
S3D	2.62 ^{b-e} ±0.03	3.14 ⁱ ±0.03	2.95 ^{kl} ±0.04	3.44 ^{lm} ±0.04	3.24 ^{kl} ±0.03	2.25 ^d ±0.03
S1PdD	2.40 ^a ±0.03	2.94 ^{jk} ±0.03	2.79 ^{ghi} ±0.03	2.55 ^f ±0.03	2.63 ^{fg} ±0.03	3.15 ^{jk} ±0.04
S2PdD	2.64 ^{c-f} ±0.03	3.04 ⁱ ±0.03	2.84 ^{ij} ±0.03	2.96 ⁱ ±0.03	2.73 ^{gh} ±0.03	2.83 ^h ±0.04
S3PdD	2.54 ^b ±0.03	2.94 ^{jk} ±0.03	2.94 ^{jk} ±0.04	3.06 ^{ij} ±0.04	4.35 ^p ±0.05	4.01 ⁿ ±0.05
Mean	2.57^A±0.12	2.93^C±0.13	2.62^B±0.10	2.86^B±0.59	3.09^C±0.72	2.62^A±0.68

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization x years, capital letters for years. ± - standard deviation

Table S3. Nitrogen and sulphur harvest index, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Nitrogen (%)			Sulphur (%)		
	2011	2012	2013	2011	2012	2013
S0	62.6 ^k ±0.74	54.2 ^e ±0.52	58.3 ^{gh} ±0.77	38.9 ^q ±0.40	32.89 ^o ±0.49	37.0 ^P ±0.80
S1Pd	62.9 ^k ±0.38	67.4 ^m ±0.49	54.6 ^e ±0.82	31.8 ^{no} ±0.23	30.3 ^{lm} ±0.70	32.6 ^{no} ±0.89
S2Pd	59.7 ⁱ ±0.19	51.5 ^d ±0.51	60.2 ^l ±0.63	27.1 ^{hij} ±0.51	29.9 ^{ll} ±0.59	29.1 ^{klt} ±1.11
S3Pd	63.5 ^{kl} ±0.39	57.7 ^{gh} ±0.47	53.9 ^e ±0.75	22.9 ^{bc} ±0.31	20.0 ^a ±0.58	36.1 ^P ±0.91
S1D	56.1 ^f ±0.38	50.8 ^{de} ±0.49	56.4 ^f ±0.75	24.4 ^{de} ±0.41	29.3 ^{kll} ±0.62	35.9 ^P ±0.91

S2D	60.2 ^{ij} ±0.20	50.1 ^{bc} ±0.26	65.3 ⁱ ±0.80	26.0 ^{gh} ±0.32	21.9 ^b ±0.72	26.7 ^{ghi} ±1.33
S3D	61.0 ^j ±0.27	46.6 ^a ±0.41	46.5 ^a ±0.97	24.2 ^{cd} ±0.36	24.6 ^{def} ±0.67	36.4 ^p ±1.05
S1PdD	61.0 ^j ±0.30	58.5 ^h ±0.39	49.1 ^b ±0.83	30.3 ^{lm} ±0.20	25.6 ^{efg} ±0.56	28.2 ^{jk} ±1.17
S2PdD	59.6 ⁱ ±0.23	69.6 ⁿ ±0.70	46.7 ^a ±1.04	25.8 ^{fg} ±0.25	31.6 ^{mn} ±0.76	27.4 ^{ij} ±1.03
S3PdD	64.2 ^{kl} ±0.24	64.8 ^t ±0.45	57.3 ^{fg} ±1.00	29.0 ^{kl} ±0.73	24.2 ^{cd} ±0.45	22.8 ^b ±1.20
Średnio	61.1 ^c ±2.96	57.1 ^B ±7.49	54.8 ^A ±6.22	28.0 ^B ±5.08	27.0 ^A ±4.04	31.2 ^C ±5.32

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization \times years, capital letters for years. \pm - standard deviation

Table S4. Phosphorus content in rape seeds and straw, mean values for interaction variants of sulphur fertilization \times years.

Variants of sulphur fertilization	Phosphorus – seeds (g kg ⁻¹ d.w.)			Phosphorus – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	6.07 ^b ±0.07	5.77 ^a ±0.06	5.77 ^a ±0.07	1.42 ^e ±0.02	1.72 ^h ±0.02	1.22 ^c ±0.02
S1Pd	6.48 ^{def} ±0.07	6.18 ^{bc} ±0.07	6.38 ^{cde} ±0.08	1.22 ^c ±0.01	1.22 ^c ±0.01	1.14 ^b ±0.01
S2Pd	6.48 ^{def} ±0.07	6.48 ^{def} ±0.07	6.68 ^{fg} ±0.08	1.21 ^c ±0.01	2.63 ⁿ ±0.03	2.81 ⁱ ±0.01
S3Pd	6.58 ^b ±0.07	6.07 ^b ±0.07	6.18 ^{bc} ±0.08	2.13 ^k ±0.02	1.82 ⁱ ±0.02	1.52 ^f ±0.02
S1D	6.28 ^{bcd} ±0.07	6.38 ^{cde} ±0.07	6.38 ^{def} ±0.08	2.23 ^l ±0.02	2.63 ⁿ ±0.03	2.33 ⁱ ±0.03
S2D	6.28 ^{bcd} ±0.07	6.78 ^g ±0.07	6.38 ^{def} ±0.09	2.43 ^m ±0.03	1.82 ⁱ ±0.02	1.61 ^g ±0.01
S3D	6.58 ^{efg} ±0.07	6.48 ^{def} ±0.07	6.48 ^{def} ±0.08	1.92 ^j ±0.02	2.83 ^p ±0.03	2.73 ^o ±0.03
S1PdD	6.07 ^b ±0.07	6.18 ^{bc} ±0.07	6.38 ^{cde} ±0.08	1.82 ⁱ ±0.02	1.32 ^d ±0.01	2.94 ^q ±0.04
S2PdD	6.68 ^{fg} ±0.07	6.58 ^{efg} ±0.07	6.28 ^{bcd} ±0.08	1.14 ^b ±0.01	1.01 ^a ±0.01	1.62 ^g ±0.02
S3PdD	6.38 ^{cde} ±0.08	6.28 ^{bcd} ±0.07	6.28 ^{bcd} ±0.09	1.21 ^c ±0.02	1.42 ^e ±0.02	1.52 ^f ±0.02
Mean	6.39 ^A ±0.22	6.32 ^A ±0.28	6.32 ^A ±0.23	1.67 ^A ±0.47	1.85 ^B ±0.62	1.84 ^B ±0.75

Means followed by the same letter are not statistically different \pm at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization \times years, capital letters for years. \pm - standard deviation

Table S5. Potassium content in rape seeds and straw, mean values for interaction variants of sulphur fertilization \times years.

Variants of sulphur fertilization	Potassium – seeds (g kg ⁻¹ d.w.)			Potassium – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	5.67 ^{ab} ±0.06	7.39 ^f ±0.08	5.77 ^{abc} ±0.07	7.19 ^d ±0.08	7.39 ^{de} ±0.08	8.40 ^{fg} ±0.10
S1Pd	5.77 ^{abc} ±0.06	7.80 ^g ±0.08	5.87 ^{bc} ±0.07	9.72 ^j ±0.10	9.21 ⁱ ±0.10	7.39 ^{de} ±0.09
S2Pd	5.87 ^{bc} ±0.06	7.79 ^g ±0.08	5.97 ^{cd} ±0.07	10.53 ^l ±0.11	7.19 ^d ±0.08	5.26 ^b ±0.07
S3Pd	5.77 ^{abc} ±0.06	8.20 ^h ±0.09	5.87 ^{bc} ±0.08	12.35 ^m ±0.13	13.36 ⁿ ±0.14	10.12 ^k ±0.13
S1D	5.77 ^{abc} ±0.06	7.80 ^g ±0.08	6.28 ^e ±0.08	12.05 ^m ±0.13	9.01 ^{hi} ±0.10	7.29 ^d ±0.09
S2D	5.57 ^a ±0.06	7.79 ^g ±0.08	5.97 ^{cd} ±0.07	14.38 ^o ±0.16	13.36 ⁿ ±0.14	4.25 ^a ±0.05
S3D	5.87 ^{bc} ±0.07	7.59 ^{fg} ±0.08	5.87 ^{bc} ±0.07	10.53 ^l ±0.11	11.03 ⁱ ±0.12	5.87 ^c ±0.07
S1PdD	5.87 ^{bc} ±0.06	7.59 ^{fg} ±0.09	5.97 ^{cd} ±0.07	13.26 ⁿ ±0.14	13.16 ⁿ ±0.14	7.69 ^e ±0.10
S2PdD	6.28 ^e ±0.07	7.59 ^{fg} ±0.08	6.17 ^{de} ±0.08	8.71 ^{gh} ±0.09	8.2 ^f ±0.09	5.77 ^c ±0.07
S3PdD	5.97 ^{cd} ±0.07	7.80 ^g ±0.09	6.18 ^{de} ±0.08	7.69 ^e ±0.10	10.02 ^{jk} ±0.11	7.39 ^{de} ±0.09
Mean	5.84 ^A ±0.19	7.73 ^C ±0.22	5.99 ^B ±0.17	10.64 ^C ±2.26	10.19 ^B ±2.30	6.94 ^A ±1.61

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization \times years, capital letters for years. \pm - standard deviation

Table S6. Phosphorus and potassium harvest index, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Phosphorus (%)			Potassium (%)		
	2011	2012	2013	2011	2012	2013
S0	66.7 ^{jk} ±0.37	56.8 ^f ±0.55	65.7 ^j ±0.89	27.1 ^{lmn} ±0.51	28.4 ^{mno} ±0.26	21.9 ⁱ ±0.45
S1Pd	69.8 ^{lm} ±0.13	69.2 ^{lm} ±0.65	68.7 ^{ll} ±1.07	20.2 ^h ±0.19	27.4 ^{lmm} ±0.27	23.5 ^j ±0.49
S2Pd	67.2 ^{kl} ±0.42	53.4 ^{de} ±0.56	56.8 ^f ±0.95	17.8 ^{de} ±0.48	33.8 ^q ±0.41	28.6 ^{no} ±0.39
S3Pd	56.7 ^f ±0.27	54.3 ^e ±0.71	62.4 ⁱ ±1.01	17.9 ^{def} ±0.63	17.8 ^{de} ±0.61	19.6 ^g ±0.74
S1D	50.0 ^{bc} ±0.60	42.6 ^a ±0.72	52.8 ^d ±0.94	14.5 ^{ab} ±0.52	21.2 ^{hi} ±0.67	26.0 ^{kl} ±0.47
S2D	50.3 ^{bc} ±0.21	54.2 ^{de} ±0.74	62.9 ⁱ ±1.00	13.3 ^a ±0.68	15.8 ^c ±0.61	37.4 ^r ±0.20
S3D	59.1 ^g ±0.18	43.7 ^a ±0.64	50.8 ^c ±1.06	19.1 ^{efg} ±0.41	19.2 ^{fg} ±0.86	30.4 ^p ±0.46
S1PdD	60.7 ^h ±0.18	59.7 ^{gh} ±0.47	48.9 ^b ±0.94	17.3 ^d ±0.54	15.3 ^{bc} ±0.45	25.5 ^k ±0.28
S2PdD	70.2 ^{mn} ±0.30	73.1 ^{kl} ±0.53	59.2 ^g ±0.98	22.08 ⁱ ±0.42	27.8 ^{lo} ±0.29	28.8 ^o ±0.09
S3PdD	71.3 ⁿ ±0.23	67.6 ^o ±0.41	62.6 ⁱ ±0.75	27.42 ^{lmn} ±0.51	26.9 ^{hl} ±0.42	25.2 ^k ±0.47
Mean	62.1^C±4.81	57.4^A±9.42	59.1^B±9.90	19.65^A±1.43	23.4^B±6.00	26.7^C±6.85

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization x years, capital letters for years. ± - standard deviation

Table S7. Calcium content in rape seeds and straw, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Calcium – seeds (g kg ⁻¹ d.w.)			Calcium – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	3.44 ^a ±0.04	3.54 ^{ab} ±0.04	3.54 ^{ab} ±0.04	12.25 ^{cde} ±0.13	15.59 ^g ±0.17	21.66 ^m ±0.27
S1Pd	3.44 ^a ±0.04	4.56 ^g ±0.04	3.54 ^{ab} ±0.04	12.45 ^{c-f} ±0.13	16.40 ^{ij} ±0.18	19.23 ⁱ ±0.24
S2Pd	3.64 ^{bc} ±0.04	3.95 ^{ef} ±0.05	3.75 ^{cd} ±0.05	12.55 ^{c-f} ±0.14	15.79 ^{ghi} ±0.17	32.09 ^o ±0.40
S3Pd	3.54 ^{ab} ±0.04	3.85 ^{de} ±0.04	3.54 ^{ab} ±0.05	12.76 ^{ef} ±0.14	15.69 ^{gh} ±0.17	16.60 ^j ±0.21
S1D	3.64 ^{bc} ±0.04	3.75 ^{cd} ±0.06	3.95 ^{ef} ±0.05	12.05 ^{bc} ±0.13	12.15 ^{kl} ±0.19	21.66 ^m ±0.27
S2D	3.64 ^{bc} ±0.05	3.95 ^{ef} ±0.04	3.64 ^{bc} ±0.05	12.65 ^{def} ±0.14	16.30 ^{ji} ±0.18	16.20 ^{hij} ±0.20
S3D	3.44 ^a ±0.05	4.05 ^f ±0.04	4.05 ^f ±0.05	12.96 ^f ±0.14	19.03 ⁱ ±0.21	26.32 ⁿ ±0.33
S1PdD	3.85 ^{de} ±0.04	3.95 ^{ef} ±0.04	3.85 ^{de} ±0.06	11.54 ^b ±0.12	12.15 ^{cd} ±0.13	15.29 ^g ±0.19
S2PdD	3.85 ^{de} ±0.04	3.85 ^{de} ±0.04	3.64 ^{bc} ±0.05	10.43 ^a ±0.11	17.41 ^k ±0.19	18.12 ^l ±0.23
S3PdD	3.64 ^{bc} ±0.05	3.75 ^{cd} ±0.05	3.95 ^{ef} ±0.05	12.55 ^{c-f} ±0.16	21.77 ^m ±0.24	19.54 ^h ±0.24
Mean	3.61^A±0.15	3.92^C±0.26	3.75^B±0.19	12.22^A±0.72	16.81^B±2.39	20.67^C±4.90

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization x years, capital letters for years. ± - standard deviation

Table S8. Magnesium content in rape seeds and straw, mean values for interaction variants of sulphur fertilization x years.

Variants of sulphur fertilization	Magnesium – seeds (g kg ⁻¹ d.w.)			Magnesium – straw (g kg ⁻¹ d.w.)		
	2011	2012	2013	2011	2012	2013
S0	2.43 ^a ±0.03	3.34 ^{de} ±0.04	3.04 ^c ±0.04	0.81 ^a ±0.01	1.42 ^d ±0.02	2.02 ^h ±0.03
S1Pd	2.63 ^b ±0.03	3.23 ^d ±0.03	3.24 ^d ±0.04	0.81 ^a ±0.01	1.42 ^d ±0.02	1.62 ^e ±0.02
S2Pd	2.63 ^b ±0.04	3.24 ^d ±0.03	3.24 ^d ±0.04	0.81 ^a ±0.02	2.02 ^h ±0.02	1.21 ^c ±0.02
S3Pd	2.42 ^a ±0.04	3.24 ^d ±0.03	3.24 ^d ±0.03	0.81 ^a ±0.02	1.42 ^d ±0.02	1.82 ^f ±0.02
S1D	2.63 ^b ±0.03	3.25 ^d ±0.03	3.44 ^e ±0.03	0.81 ^a ±0.01	1.61 ^d ±0.01	1.82 ^f ±0.02

S2D	2.64 ^b ±0.03	3.24 ^d ±0.04	3.24 ^d ±0.04	0.81 ^a ±0.01	1.82 ^f ±0.01	1.01 ^b ±0.01
S3D	2.63 ^b ±0.03	3.24 ^d ±0.03	3.24 ^d ±0.04	1.01 ^b ±0.01	2.23 ⁱ ±0.02	2.23 ⁱ ±0.03
S1PdD	2.63 ^b ±0.03	3.24 ^d ±0.03	3.24 ^d ±0.04	0.81 ^a ±0.01	1.62 ^d ±0.02	2.02 ^h ±0.03
S2PdD	2.53 ^{ab} ±0.04	3.44 ^e ±0.04	3.24 ^d ±0.03	0.81 ^a ±0.01	1.21 ^c ±0.01	1.82 ^f ±0.02
S3PdD	2.63 ^b ±0.03	3.64 ^f ±0.04	3.24 ^d ±0.04	0.81 ^a ±0.02	1.92 ^g ±0.02	1.62 ^e ±0.02
Mean	2.58 ^A ±0.09	3.31 ^C ±0.13	3.24 ^B ±0.10	0.83 ^A ±0.06	1.67 ^B ±0.31	1.72 ^C ±0.35

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization \times years, capital letters for years. ± - standard deviation

Table S9. Calcium and magnesium harvest index, mean values for interaction variants of sulphur fertilization \times years.

Variants of sulphur fertilization	Calcium (%)			Magnesium (%)		
	2011	2012	2013	2011	2012	2013
S0	11.7 ^{lm} ±0.17	8.3 ^{d-h} ±0.53	6.3 ^b ±0.81	58.6 ^p ±0.40	47.8 ^{jk} ±0.59	37.8 ^c ±1.39
S1Pd	10.4 ^{ijk} ±0.18	11.3 ^{kl} ±0.44	6.7 ^{bc} ±0.77	57.6 ^p ±0.39	50.3 ^l ±0.74	43.5 ^{gh} ±1.22
S2Pd	10.2 ^{ijk} ±0.24	10.7 ^{kl} ±0.51	41 ^a ±0.58	55.5 ^o ±0.92	43.3 ^g ±0.62	48.6 ^k ±0.69
S3Pd	11.3 ^{kh} ±0.18	8.2 ^{d-g} ±0.60	8.0 ^{def} ±0.86	58.2 ^p ±0.43	44.8 ⁱ ±0.26	42.3 ^{ef} ±1.21
S1D	9.4 ^{hi} ±0.11	6.1 ^b ±0.67	6.9 ^{bc} ±0.83	53.3 ^{lm} ±0.43	38.1 ^c ±0.55	43.5 ^g ±1.54
S2D	10.3 ^{ijk} ±0.14	7.1 ^{bcd} ±0.53	8.8 ^{fgh} ±0.80	55.9 ^o ±0.21	33.4 ^b ±0.50	57.9 ^p ±1.41
S3D	10.1 ^{ij} ±0.14	6.5 ^{bc} ±0.52	6.4 ^{bc} ±0.79	52.2 ⁱ ±0.31	33.0 ^a ±0.63	38.8 ^c ±0.99
S1PdD	13.4 ⁿ ±0.21	9.3 ^{ghi} ±0.67	10.4 ^{ijk} ±0.78	60.2 ^q ±0.52	38.3 ^c ±0.53	41.3 ^{de} ±1.47
S2PdD	12.6 ^m ±0.09	8.3 ^{e-h} ±0.76	7.2 ^{bcd} ±0.82	55.2 ^{no} ±0.46	54.1 ^{m-n} ±0.69	40.1 ^d ±1.16
S3PdD	12.3 ^{lm} ±0.20	7.3 ^{a-d} ±0.47	7.4 ^{cde} ±0.78	61.3 ^q ±0.47	47.1 ^j ±0.57	44.7 ^{hi} ±1.05
Mean	11.2 ^C ±1.43	8.3 ^B ±1.66	7.2 ^A ±1.67	56.8 ^C ±2.97	43.3 ^A ±6.41	43.8 ^B ±5.62

Means followed by the same letter are not statistically different at the $\alpha = 0.05$ level. Small letters for variants of sulphur fertilization \times years, capital letters for years. ± - standard deviation