

Review

# The Endocannabinoid System and PPARs: Focus on Their Signalling Crosstalk, Action and Transcriptional Regulation

Fabio Arturo Iannotti \*  and Rosa Maria Vitale \* 

Institute of Biomolecular Chemistry, National Research Council (ICB-CNR), Via Campi Flegrei 34, 80078 Pozzuoli (NA), Italy

\* Correspondence: fabio.iannotti@icb.cnr.it (F.A.I.); rmvitale@icb.cnr.it (R.M.V.)

**Abstract:** Peroxisome proliferator-activated receptors (PPARs) are a family of nuclear receptors including PPAR $\alpha$ , PPAR $\gamma$ , and PPAR $\beta/\delta$ , acting as transcription factors to regulate the expression of a plethora of target genes involved in metabolism, immune reaction, cell differentiation, and a variety of other cellular changes and adaptive responses. PPARs are activated by a large number of both endogenous and exogenous lipid molecules, including phyto- and endo-cannabinoids, as well as endocannabinoid-like compounds. In this view, they can be considered an extension of the endocannabinoid system. Besides being directly activated by cannabinoids, PPARs are also indirectly modulated by receptors and enzymes regulating the activity and metabolism of endocannabinoids, and, vice versa, the expression of these receptors and enzymes may be regulated by PPARs. In this review, we provide an overview of the crosstalk between cannabinoids and PPARs, and the importance of their reciprocal regulation and modulation by common ligands, including those belonging to the extended endocannabinoid system (or “endocannabinoidome”) in the control of major physiological and pathophysiological functions.

**Keywords:** peroxisome proliferator-activated receptors (PPARs); endocannabinoid system (ECS); endocannabinoidome; plant cannabinoids; cannabinoid receptors; metabolism; neuroprotection



**Citation:** Iannotti, F.A.; Vitale, R.M. The Endocannabinoid System and PPARs: Focus on Their Signalling Crosstalk, Action and Transcriptional Regulation. *Cells* **2021**, *10*, 586. <https://doi.org/10.3390/cells10030586>

Academic Editor: Hiroshi Miyamoto

Received: 4 February 2021

Accepted: 3 March 2021

Published: 7 March 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. PPAR Receptors Classification, Distribution, and Function

Peroxisome proliferator-activated receptors (PPARs) are a family of ligand-activated receptors/transcriptional factors composed by three distinct isoforms called PPAR $\alpha$  (nuclear receptor subfamily 1 group C, NR1C1), PPAR $\beta/\delta$  (NR1C2), and PPAR $\gamma$  (NR1C3), each of which is encoded by independent genes in rodents and humans. Although co-expressed in several different types of organs and tissues, each isoform shows distinctive functional characteristics and ligand specificity [1,2]. Since their identification, PPARs have been recognized as sensing receptors for a variety of endogenous lipids (such as unsaturated, monounsaturated, and poly-unsaturated fatty acids) and natural exogenous compounds.

### 1.1. PPAR $\alpha$

PPAR $\alpha$  is highly expressed in organs and tissues characterized by a high rate of fatty acid catabolism for energy production including liver, brown adipose tissue, endocrine tissues, gastrointestinal tract, cardiac and skeletal muscle. To a lesser extent, PPAR $\alpha$  is present in the kidney, adrenal tissues, endothelial and immune (i.e. macrophages, monocytes, and lymphocytes) cells [3,4]. Additionally, PPAR $\alpha$  has been found in specific brain areas where it exerts anti-inflammatory and neuroprotective actions (see next section) [5–9].

#### 1.1.1. Role of PPAR $\alpha$ in Metabolism

PPAR $\alpha$  plays a major role in metabolic homeostasis regulating lipid metabolism. Specifically, during the fed-to-fasted transition, PPAR $\alpha$  drives the production of enzymes responsible for fatty acid oxidation (FAO) and the synthesis of ketone bodies from fatty

acids in the liver. Thereby, PPAR $\alpha$  works as a hub that integrates multiple metabolic signals to orchestrate the switch from glucose to fatty acid utilization for energy production [10,11]. Additionally, PPAR $\alpha$  governs hepatic amino acid metabolism [12]. Studies on PPAR $\alpha$  in other tissues including the heart, small intestine, skeletal muscle and brain have indicated that the role of PPAR $\alpha$  in metabolic homeostasis is well conserved between different cell types [12–15]. Interestingly, *Ppara* KO mice, under a normal dietary regimen, do not display pronounced anomalies in their phenotype. However, under fasting conditions or a high-fat diet, *Ppara* KO mice develop hypoglycemia and dyslipidemia characterized by excessive production of triacylglycerols [13,16,17]. Moreover, the lack of PPAR $\alpha$  causes cardiac metabolic and contractile dysfunction, morphological and functional alterations in the brain in association with micro-/macro- vasculature dysfunctions [15,18,19]. In general, although the *Ppara* KO mouse model still shows unresolved aspects that are most likely attributable to the compensatory role of the other two PPAR isotypes, it allowed a deeper understanding of the role of PPAR $\alpha$  in energy metabolism.

#### 1.1.2. Neuroprotective and Anti-Inflammatory Role of PPAR $\alpha$

Although all three PPAR isoforms are expressed in the nervous system during embryogenesis, only PPAR $\beta/\delta$  expression remains high in the brain, whereas PPAR $\alpha$  and PPAR $\gamma$  expression decreases postnatally and remain restricted to the specific brain areas [20]. In particular, PPAR $\alpha$  is expressed in basal ganglia, the reticular formation, some thalamic, mesencephalic and cranial motor nuclei and the large motoneurons of the spinal cord [5]. PPAR $\alpha$  is expressed in dopamine neurons of the substantia nigra and spiny neurons of the dorsal striatum where it decreases dopaminergic transmission thus exerting an important control in behavioural responses [8,21,22]. The expression of PPAR $\alpha$  is also reported in different subfields of the hippocampus of rodents, where it is also involved in the control of neuronal excitability and synaptic plasticity via cyclic AMP response element-binding protein (CREB) [23]. Moreover, PPAR $\alpha$  is present in oligodendrocytes, microglia and astrocytes [24–26]. The studies reported above as well many others suggest that activation of PPAR $\alpha$  in the brain initiates anti-oxidative and anti-inflammatory processes that confer neuroprotection. An effect also exerted by preserving the microvasculature activity [27]. The neuroprotective role of PPAR $\alpha$  has been documented in numerous studies using several genetic and/or pharmacological models of neurodegenerative disorders including stroke, Alzheimer's disease, Parkinson's disease, traumatic brain injury, diabetic peripheral neuropathy, and retinopathy [7,9,22,28,29]. Besides its neuroprotective role, PPAR $\alpha$  exerts important anti-inflammatory effects also in peripheral organs and tissues [3,30]. In this regard, Luisa et al., 2009, using an experimental model of inflammatory bowel disease, showed that *Ppara* KO mice compared to wild-type (WT) ones, have a deficient anti-inflammatory response [31]. Additionally, PPAR $\alpha$  agonists have been reported to exert anti-inflammatory, antipyretic, anti-atherogenic and analgesic effects as demonstrated in experimental models of spinal cord trauma [28,32], neuropathic pain [33–35] and high-fat diet (HFD) induced atherosclerosis [36].

#### 1.2. PPAR $\gamma$

PPAR $\gamma$  is considered a master regulator of adipogenesis and is abundantly expressed in adipose tissue where it is primarily involved in fat and carbohydrate metabolism. Additionally, PPAR $\gamma$  exerts anti-inflammatory effects in several types of tissues and organs by repressing the expression and function of pro-inflammatory factors such as NF- $\kappa$ B and Nrf2/CREB and it is implicated in cancer and atherosclerosis [32]. The expression of PPAR $\gamma$  was also reported in other organs and tissues including the liver, skeletal muscle, spleen, heart, placenta, lung, ovary, and also brain (glial cells and neurons) [33,34].

##### 1.2.1. Role of PPAR $\gamma$ in Metabolism

In both rodents and humans, PPAR $\gamma$  is a master regulator of adipocytes differentiation as well as glucose and lipid metabolism [37]. Additional actions of PPAR $\gamma$  include the

regulation of adipokines and inflammatory mediators expression, M1/M2 macrophage polarization, atherosclerosis and bone formation [38,39]. Genetic ablation of PPAR $\gamma$  in mice is lethal [40]. However, in an elegant study conducted by Gavrilova et al. [41], it was demonstrated that young mice with a PPAR $\gamma$ -deficient liver show a similar profile of wild-type (wt) mice. However, with ageing, LPPAR $\gamma$  Knock-Out (*Lppar $\gamma$ -KO*) mice develop fat intolerance, increased adiposity, hyperlipidemia, and insulin resistance. Interestingly, when fed with a lipogenic diet even young LPPAR $\gamma$ -KO mice developed obesity. Other studies showed that mutant PPAR $\gamma$  mice show a complex metabolic phenotype including increased lean mass with organomegaly, hypermetabolism, hyperphagia, and lipoatrophy [42–44]. Moreover, Lüdtke et al. reported that PPAR $\gamma$  mutations lead to a familial form of lipodystrophy [45]. Therefore, these findings provide evidence that PPAR $\gamma$  is a key target to protect the liver as well as other organs and tissues from fat accumulation, insulin resistance, and uncontrolled inflammatory responses. Interestingly, the PPAR $\gamma$  gene is characterized by distinct mRNA isoforms due to alternative splicing of five exons at the 5'-terminal regions (A1, A2, B, C, and D). In particular, among the seven distinct isoforms identified so far, the most known are PPAR- $\gamma$ 1, - $\gamma$ 2, and - $\gamma$ 3. The PPAR $\gamma$ 1 mRNA isoform is expressed in a wide range of organs and tissues including those aforementioned and also in the immune cells (e.g., monocytes/macrophages). On the contrary, the expression of PPAR $\gamma$ 2 mRNA seems restricted to adipose tissue, and PPAR $\gamma$ 3 is localized in macrophages, colon, and adipose tissue [37,46–49]. Of note, in 2004 Zhang et al. showed that PPAR $\gamma$ 2 deficient mice (*PPAR $\gamma$ <sup>-/-</sup>*) have an impaired fat metabolism and insulin resistance, thus demonstrating the central role of PPAR $\gamma$ 2 in adipogenesis [14]. It is important to recall that different PPAR $\gamma$  isoforms may be responsible for unique tissue-specific biological effects in response or not to endogenous and/or exogenous ligands.

### 1.2.2. Neuroprotective and Anti-Inflammatory Role of PPAR $\gamma$

PPAR $\gamma$ , similarly to PPAR $\alpha$ , in the brain exerts anti-inflammatory and neuroprotective effects through the inhibition of NF $\kappa$ B, AP-1, STATs, and iNOS [50]. This finding has been confirmed in several animal models of Alzheimer's and Parkinson's disease. Moreover, the synthetic PPAR $\gamma$  agonist pioglitazone was shown to extend the survival of SOD1-G93A, a mouse model of Amyotrophic lateral sclerosis (ALS), by delaying the onset of the disease and preventing the death of motor neuron cells [51]. Activation of PPAR $\gamma$  has been shown to exert neuroprotective effects also by preventing astro- and microglial activation [52]. Additionally, PPAR $\gamma$  has been shown to downregulate the expression of pro-inflammatory factors and free radical production in various intestinal disorders including colon cancer, gastritis and irritable bowel syndrome, and to also exert anti-nociceptive effects against somatic pain, allergic and skin diseases [53–55].


### 1.3. PPAR $\beta/\delta$

PPAR $\beta/\delta$  is ubiquitously expressed and its biological functions mainly overlap with those of the other two PPAR isoforms. Recently, the expression of PPAR $\beta/\delta$  was demonstrated in neurons, astrocytes, oligodendrocytes, and also, in microglial cells [56,57]. Several studies demonstrate that PPAR $\beta/\delta$  plays an important role in inflammatory processes and, due to its proangiogenic and anti-/pro-carcinogenic properties, it is considered a therapeutic target for treating metabolic syndrome, dyslipidemia, diabetes while the role of PPAR $\beta/\delta$  on cancerogenesis is still debated [58,59]. In the brain, PPAR $\beta/\delta$  could act as a protective factor for counteracting inflammation and promoting antioxidant mechanisms [60]. Additionally, PPAR $\beta/\delta$  is known to control cell cycle, proliferation, differentiation and also inhibit cell death by promoting the expression of ILK and PDK1 in many types of cells [61,62]. However, compared to the other two isoforms, the role of PPAR $\beta/\delta$  in inflammation still needs to be fully elucidated. For example, at peripheral level, stimulation of PPAR $\beta/\delta$  was shown to promote the inhibition of inflammatory response in streptozotocin-induced diabetic nephropathy and vasculopathy [63,64]. In contrast, other studies report that PPAR $\beta/\delta$  appears to promote inflammation in other con-


texts. This latter evidence was observed in murine models of skin disorders (i.e. psoriasis) and arthritis [65].

## 2. Structure of PPARs

Like other nuclear receptor superfamily members, PPAR structure is organized in four domains, named A/B, C, D, and E/F: (i) the N-terminal A/B domain contains the ligand-independent activation function 1 (AF1) responsible for the transcriptional activation; (ii) the C domain consists in the DNA-binding domain (DBD), formed by two zinc-finger motifs responsible for the binding to peroxisome proliferator response elements (PPREs) within the promoter regions of target genes; (iii) the D domain (hinge domain) modulates the ability of the receptor to bind DNA and it is involved in corepressors binding; (iv) C-terminal E/F domains correspond to the ligand-binding domain (LBD) which contains the ligand-binding pocket (LBP), the ligand-dependent activation function (AF-2) and regions important for the heterodimerization with the retinoid X receptor (RXR) [66,67] (Figure 1A). The PPAR-RXR heterodimer is a prerequisite for PPARs to bind the PPREs in the promoter region of target genes. They act as permissive heterodimers, being activated either by PPAR or RXR ligands. However, the simultaneous presence of both ligands gives rise to a synergistic response [68]. When PPAR-RXR heterodimers are not bound to a ligand, they act as repressors through association with corepressor complexes (Figure 1B,C; see also Figure 2). The ligand binding to LBP induces conformational changes in the AF2 region, facilitating the recruitment of coactivators and the release of corepressors [69]. In detail, canonical agonists within the LBP form a network of hydrogen bonds stabilizing the conformation of a short helix, the helix12 (H12), which in turn allows the interaction of coactivators with the AF-2 surface. However, PPAR ligands can also bind to distinct sub-regions or allosteric sites and activate these receptors through H12-independent mechanisms to elicit a partial agonist activity [70]. Besides the ligand-dependent mechanism, other mechanisms are known to regulate the transactivation and trans-repression activity of PPARs. In particular, the PPAR activity is governed by various post-translational modifications including phosphorylation, SUMOylation, ubiquitination, acetylation, and O-GlcNAcylation [71].




**Figure 1.** PPAR domains and 3D model of their interaction with RXR $\alpha$  and target DNA (A) Schematic representation of the PPAR domain organization and (B,C) orthogonal views of the x-ray structure of PPAR $\gamma$ /RXR $\alpha$  heterodimer in complex with DNA and coactivator peptides (PDB id: 3DZY). PPAR $\gamma$  is colored in slate blue with helix H12 in sky blue, RXR $\alpha$  in gray, the coactivator peptides in green and salmon and DNA in tan. The ligands rosiglitazone and 9-cis-retinoic acid are shown in rose brown and khaki, respectively, while the Zn(II) ions are in gold. Nitrogen, oxygen and sulfur atoms are colored in blue, red and yellow, respectively. This figure has been realized with the Chimera v.1.14 program.



**Figure 2.** Representative illustration of PPARs activating pathways. (A) In absence of ligand binding, the nuclear corepressor complex prevents PPAR/RXR binding to their DNA binding element, called PPAR response element (PPRE). (B) In presence of a ligand, PPAR/RXR dimer binds to the PPRE of target genes and regulates the transcription of target genes. Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 $\alpha$ ) is a transcriptional coactivator of the PPAR/RXR complex, known to be activated by intracellular factors such as AMPK, SIRT and MAP kinases.

### 3. The Endocannabinoidome, Including and beyond the Endocannabinoid System

The endocannabinoid system (ECS) refers to a large group of lipid mediators, enzymes, and receptors largely distributed throughout the mammalian body. The major component of ECS are 1) two main lipid signaling molecules, the eCBs anandamide (*N*-arachidonoyl-ethanolamine, AEA) and 2-arachidonoyl-glycerol (2-AG); 2) a large number of biosynthesizing and inactivating enzymes of which the most studied ones are ABDH4, GDE-1, NAPE-PLD and PTPN22 for the biosynthesis of AEA from *N*-arachidonoyl-phosphatidylethanolamine; FAAH for its degradation to arachidonic acid (AA) and ethanolamine; DAGL $\alpha$  and DAGL $\beta$  for the biosynthesis of 2-AG from sn-2-AA-containing diacylglycerols, and ABDH6, ABDH12 and MAGL for its degradation to AA and glycerol; 3) two main metabotropic receptors responsive to AEA and 2-AG, named cannabinoid receptor of type-1 (CB1) and type 2 (CB2) [72–74] (Figure 3). Beyond its classical definition, numerous studies revealed that many other *N*-acyl-ethanolamines (NAEs) including (i) *N*-palmitoylethanolamine (PEA), (ii) *N*-oleylethanolamine (OEA), (iii) *N*-linoleylethanolamine (LEA), (iv) *N*-stearoylethanolamine (SEA), DHEA and v) *N*-docosahexaenoyl-ethanolamine; monoacylglycerols (2-LG, 2-linoleoyl glycerol; 2-OG, 2-oleoyl glycerol); *N*-acyldopamines/taurines/serotonins together with G-protein coupled receptors (GPR55, GPR119, GPR18), ion channels (TRPV1 and TRPV2), metabolic enzymes (i.e. lipoxygenase LOX, cyclooxygenase COX, and cytochrome P450) as well as their related metabolites (i.e. eicosanoids) are structurally and functionally related to the ECS activity. Moreover, 2-AG was found to act as a direct modulator of the GABAA receptor [75]. Therefore, these discoveries led to expanding our view of the ECS and to look at it as the ‘endocannabinoidome’ (Figure 2).



**Figure 3.** Synthesis, inactivation, receptors and main metabolic functions of endocannabinoidome mediators. Thick arrows denote the biochemical reactions and functional connections underlying endocannabinoidome mediator action. Not shown, for the sake of clarity, are the negative effects exerted by almost all unsaturated *N*-acyl-amides tested so far on T-type ( $\text{Ca}_v3$ )  $\text{Ca}^{2+}$  channels. Blunted arrows denote inhibition. Abbreviations: 2-AG, 2-arachidonoylglycerol; 2-LG, 2-linoleoyl glycerol; 2-OG, 2-oleoyl glycerol; 5/12/15-LOX, 5/12/15-lipoxygenase; 15 HAEA, 15(S)-HETE Ethanolamide; AA, arachidonic acid; ABH4/6/12,  $\alpha\beta$ -hydrolase 4/6/12; AEA, Anandamide; CB1/2, cannabinoid receptor 1/2; COX2, cyclooxygenase 2; DAG, diacylglycerols; DHEA, *N*-docosahexaenoyl-ethanolamine; EET-EA, epoxyeicosatrienoic acid ethanolamide; FA, fatty acid; FAAH, fatty acid amide hydrolase; FA, free fatty acids; GDE1, glycerophosphodiester phosphodiesterase 1; GPR, G-protein-coupled receptor; LEA, *N*-linoleoyl-ethanolamine; MAGL, monoacylglycerol lipase; MAG, monoacylglycerols NAAA, *N*-acylethanolamine-hydrolysing acid amidase, NAPE, *N*-acyl-phosphatidylethanolamine; NAPEPLD, *N*-acyl-phosphatidylethanolamine-specific phospholipase D; NAT, *N*-acyltransferase; OEA, *N*-oleoyl-ethanolamine; P450, cytochrome p450 oxygenases; PEA, *N*-palmitoylethanolamine; PGE2, Prostaglandin E2; PLC $\beta$ , phospholipase C $\beta$ ; PPAR $\gamma$ , peroxisome proliferator-activated receptor  $\gamma$ ; PPAR $\alpha$ , peroxisome proliferator-activated receptor  $\alpha$ ; SEA, *N*-stearoyl-ethanolamine; TRPV1, transient receptor potential vanilloid type-1 channel; TRPV2, transient receptor potential vanilloid type-2 channel.


#### 4. PPARs Ligands

After the discovery of all three PPARs isoforms in the early 90s, a large number of natural and synthetic ligands have been identified. The best known synthetic ligands of PPAR $\alpha$  are fibrates (e.g., clofibrate, fenofibrate, and bezafibrate), a class of drugs used to reduce the risk of cardiovascular disease in patients with dyslipidemia due to their ability to lower plasma triglyceride levels and elevate HDL cholesterol. PPAR $\gamma$  is the molecular target of glitazones such as pioglitazone and rosiglitazone, which are approved drugs for diabetes. Moreover, many other synthetic dual and/or pan- PPARs agonists (e.g., like muraglitazar and tesaglitazar) have been tested in the treatment of obesity, dyslipidemias and type 2 diabetes or for the treatment of hypertension [76–80]. To date,

several endogenous PPAR ligands have been reported so far, including fatty acids (e.g., docosahexaenoic and eicosapentaenoic acid), eicosanoids (e.g., leukotriene B<sub>4</sub> stimulates PPAR $\alpha$ , and prostaglandin PGJ<sub>2</sub> activates PPAR $\gamma$ ). In this context, we focus on the role of (endo)cannabinoid molecules as PPAR ligands and the functional interaction between PPARs and enzymes and receptors related to the endocannabinoid system.

#### *Cannabinoids and Cannabinoid-Like Molecules as PPARs Modulators*

It has been demonstrated that PPARs, mainly the isoforms  $\alpha$  and  $\gamma$ , concur in mediating the metabolic and anti-inflammatory effects of (endo)cannabinoid molecules, towards which they exhibit a different selectivity profile. In this regard, it has been demonstrated that synthetic and plant-derived cannabinoids attenuate neuroinflammation and neurodegeneration in animal models of acute or chronic neurodegenerative disorders through activation of cannabinoid receptors and PPAR $\gamma$  pathway [81–84]. While natural and synthetic phytocannabinoids including  $\Delta^9$ -tetrahydrocannabinol ( $\Delta^9$ -THC), Cannabidiol (CBD),  $\Delta^9$ -THC acid, ajulemic acid, quinone derivatives [85–89], and the recently identified cannabimovone [90] are PPAR $\gamma$  agonists, the acidic derivatives cannabigerolic and cannabidiolic acids exhibit a dual agonist profile [91] (Figure 4). The endocannabinoid anandamide (AEA) activates both PPAR $\alpha$  [92] and PPAR $\gamma$  [93], albeit its efficacy and potency toward PPAR $\alpha$  are higher in comparison to PPAR $\gamma$ . The endocannabinoid-like-compounds OEA and PEA, structurally related to AEA, do not bind endocannabinoid receptors with high affinity but exert their metabolic and anti-inflammatory effects mainly through PPAR $\alpha$  activation [94–96]. A comparative study carried out on a series of oleoylethanolamines, such as PEA, SEA, OEA, AEA, LEA, DHEA and eicosapentaenoyl-ethanolamide (EPEA) showed that all the NAEs activate PPAR $\alpha$ , albeit with different potency, and OEA and LEA were the most potent [97]. The endocannabinoid 2-arachidonoyl glyceryl ether (2-AGE, noladin ether) and virodhamine (O-arachidonoyl ethanolamine; O-AEA) bind to PPAR $\alpha$  with a rank order noladin ether>anandamide>virodhamine (EC<sub>50</sub> values 10–30  $\mu$ M), although they show less efficacy than anandamide at 10  $\mu$ M [98]. 2-AG and its non-hydrolyzable analogue, 2-AGE, were shown to activate PPAR $\gamma$ , through which 2-AG promotes the suppression of interleukin (IL-2) [99]. However, the dependence on COX-2 metabolism of 2-AG for the suppression of IL-2 suggested the active role of a 2-AG metabolite in PPAR $\gamma$  activation, identified by Raman et al. [100] as the 15d-PGJ<sub>2</sub>-glycerol ester, while Kozak et al. [101] showed that lipoxygenase metabolism of 2-AG resulted in the release of the PPAR $\alpha$  agonist 15-hydroxyeicosatetraenoic acid glyceryl ester (15-HETE-G). The 12/15-LOX arachidonic acid metabolites 12-HETE and 15-HETE increase the transcriptional activity of PPAR $\gamma$  in primary cortical neurons, eliciting neuroprotection through the inhibition of the inducible NF- $\kappa$ B, NO synthase, and COX-2, a suppressive effect reversed by the PPAR $\gamma$  antagonist GW9662 [102]. Yu et al. showed that arachidonic acid (AA), but not its precursor AEA, activates PPAR $\beta/\delta$ , unveiling a dual role for the fatty acid-binding protein FABP5 in AA-mediated activation of PPAR $\beta/\delta$ , since it promotes both the hydrolysis of AEA by FAAH and the translocation of the enzymatic product arachidonic acid to the nucleus, thus enabling PPAR $\beta/\delta$  activation by AA [103]. The cannabinoid-like oleamide also was shown to activate PPAR $\beta/\delta$ , along with PPAR $\alpha$  and PPAR $\gamma$  [104]. Further details on cannabinoid molecules as PPAR ligands is beyond the scope of the present review since this topic has been extensively reviewed by Pistis & O'Sullivan [105].



**Figure 4.** Representative illustration showing the different ability of some plant cannabinoids, ajulemic acid and quinone derivatives to activate PPAR $\alpha$  and/or PPAR $\gamma$ .

### 5. Cross-Talk between PPARs and Metabolizing Enzymes Related to Endocannabinoids

A pharmacological strategy to promote the activity of PPARs is to elevate the level of endogenous modulators. In this view, Mazzola et al. [106], using a passive-avoidance task in rats, demonstrated that the pharmacological inhibition of FAAH enzyme by URB597 recapitulates the effects of selective PPAR $\alpha$  agonist administration on memory enhancement, by increasing the level of the endogenous PPAR $\alpha$  agonists *N*-acylethanolamines (NAEs) such as AEA and OEA. A similar strategy has been pursued with NAAA inhibitors to increase the endogenous levels of PEA [107], which, besides directly activating PPAR $\alpha$ , enhances the endogenous tone of *N*-acylethanolamines (NAEs) through the down-regulation of FAAH expression and activity [108], thus explaining, at least in part, its entourage effect. Besides the indirect effects of FAA and NAAA inhibitors, PPARs have been shown to directly regulate the expression level of some endocannabinoid metabolizing enzymes such as cyclooxygenases and lipoxygenases. Cyclooxygenase-2 (COX-2) catalyzes the biosynthesis of prostaglandins (PGs) from arachidonic acid. Among PGs, the PGD<sub>2</sub> metabolite 15-deoxy-D<sup>12,14</sup> PGJ<sub>2</sub> (15d-PGJ<sub>2</sub>) has been identified as a potent agonist PPAR $\gamma$ . Several reports have shown that COX-2 is down-regulated upon PPAR $\gamma$  [109] stimulation, while the epidermal growth factor (EGF), a potent activator of COX-2, induces an enhanced COX-2 expression and a decrease of PPAR $\gamma$  mRNA levels. This down-regulation is linked to COX-2 signalling since it is reversed by NS-398, a selective COX-2 inhibitor [109]. The ability of PPAR $\gamma$  agonists to down-regulate COX-2 expression has been observed in breast cancer cells, macrophages, and cervical cancer cell lines. However, the findings that PPAR $\gamma$  ligands can also up-regulate COX-2 expression in several cell lines [110] is suggestive of a tissue-specific cross-regulation. Moreover, the positive or negative modulation of COX-2 expression by PPAR $\gamma$  ligands could be either PPAR $\gamma$ -dependent or independent [109]. Lipoxygenases (LOXs) are enzymes that catalyze the conversion of polyunsaturated fatty acids, mainly arachidonic and linoleic acids, to hydroxy fatty acid and are classified according to the carbon atom they peroxygenase on a particular substrate. In humans, two forms of 15-LOXs, namely 15-LOX-1 and 15-LOX-2, occur, leading to 13-S-hydroxyoctadecadienoic acid (13-S-HODE), respectively, which activate PPAR $\gamma$ . It was found [111] that an inverse relationship exists between 15-LOX-2 and PPAR $\gamma$  expression levels in normal vs tumour cells, suggestive of the occurrence of a feedback mechanism regulating the expression levels of these proteins. Indeed, the overexpression of PPAR $\gamma$ 1 and PPAR $\gamma$ 2 was shown to dose-dependently downregulate 15-LOX-2 expression in prostate epithelial cells, effect increased by treatment with the 15-S-HETE [48]. Similarly, the overexpression of

PPAR $\gamma$ 1 downregulates 15-LOX-2 in normal epithelial cells from breast or lung tissues, effect enhanced by 15-S-HETE. The feedback regulation of PPAR $\gamma$  by 15-LOX-2 was determined [111] by overexpressing 15-LOX-2 in tumor cells with high levels of PPAR $\gamma$  and observing a concentration-dependent downregulation of PPAR $\gamma$  with increasing amounts of 15-LOX-2, an effect enhanced by coincubation with 15-S-HETE.

## 6. Cross-Talk between PPARs and GPCRs


PPARs have been shown to modulate the expression level of G-protein coupled receptors (GPCRs) linked to cell metabolism, inflammation and cancer. In turn, the activation of certain GPCRs was shown to modulate the expression levels of PPARs. GPR120 is the cognate receptor for omega 3 long-chain fatty acids and it is involved in insulin sensitization and inflammation. It is highly expressed in mature adipocytes, where it stimulates adipocyte differentiation [112–115]. Omega 3 fatty acids activate PPAR $\gamma$  through both direct binding [116,117] and the GPR120-mediated PI3K pathway [118]. Silencing or overexpression of GPR120 in adipocytes was shown to inhibit or enhance the expression of PPAR $\gamma$ , respectively [119]. Silencing of PPAR $\gamma$  inhibits the induction of microRNA-143 by GPR-120 in adipocytes, suggesting the involvement of PPAR $\gamma$  in its GPR120-mediated transcriptional activation [119]. The transcriptome analysis in human adipocytes treated with the PPAR $\gamma$  agonist rosiglitazone showed that it induces the expression of the anti-lipolytic GPR81 and GPR109A in human and murine adipocytes as well as the human-specific GPR109B in differentiated adipocytes. The reduced expression of PPAR $\gamma$  by siRNA-mediated knockdown or the use of the PPAR $\gamma$ -specific antagonist GW9662 was shown to reduce the levels of both GPR81 and GPR109A, indicating that PPAR $\gamma$  directly regulate the expression of these receptors, although responsive PPRE elements were found only within the promoter region of *Gpr81* gene. Thus, it remains unclear whether PPAR $\gamma$  controls the transcriptional activity of the *GPR109A* gene by direct mechanisms [120]. Evidence of cross-talk between GPR109A and PPAR $\gamma$  was also reported by Knowles et al. [121], who showed that Niacin, a GPR109A agonist, induces transcriptional activity and activation of PPAR $\gamma$  in macrophages through the stimulation GPR109A-mediated of the prostaglandin synthesis pathway. Additionally, PPAR $\gamma$  upregulates the expression of both the short-chain free fatty acid receptor GPR43, another anti-lipolytic G-protein coupled receptor expressed in adipocytes, and the long-chain fatty acid receptor GPR40 in pancreatic  $\beta$ -cells [122]. Besides the transcriptional effects on GPRs involved in cell metabolism, PPAR $\gamma$  was shown to downregulate the chemokine receptor CXCR4, which is highly expressed in cancer cell metastasis [123].

## 7. Crosstalk between PPARs and Cannabinoids Receptors

### 7.1. CB1R and PPAR $\alpha$

Azar et al. in a recent paper disclosed the existence of a molecular link between CB1R and p53/miR-22/SIRT1/PPAR $\alpha$  signaling in hepatocytes [124]. Using unbiased bioinformatics techniques and combined in vitro and in vivo experiments, they found that PPAR $\alpha$  is evolutionarily coevolved with CB1R and is negatively modulated by CB1R in hepatocytes since ACEA can downregulate both mRNA and protein levels of PPAR $\alpha$  as well as its targeted genes. These effects were reversed by the CB1R antagonist AM6545, which conversely induces the hepatic expression of PPAR $\alpha$  and upregulates its target genes in WT diet-induced obese mice (DIO) after 7-day treatment. The authors also provided evidence that AM6545 is unable to directly interact with PPAR $\alpha$ , further corroborating the finding that hepatic PPAR $\alpha$  is controlled by the CB1R. In DIO mice treated with AM6545, a robust upregulation of the eCBs PPAR $\alpha$  ligands AEA, AA, OEA, and PEA was observed at hepatic level, probably due to an increased synthesis rather than the degradation rate since AM6545 is ineffective toward FAAH or MAGL. The increased level of these endogenous ligands is supportive of the correlation between the CB1R blockage and the increased hepatic activation of PPAR $\alpha$ . Next, the authors found that the expression and function of SIRT1, a key modulator of hepatic PPAR $\alpha$  activity, is regulated by CB1R

since its activation reduces both the expression and the deacetylase activity of SIRT1 in hepatocytes, effect reversed by AM6545 (Figure 5). Since the abnormal expression of microRNA (miR-22) is reportedly involved in the regulation of both SIRT1 and PPARs expression or activity in various tissues, a miR sequence profiling of liver tissues from the DIO mice treated with AM6545 or vehicle was performed. The authors found that the expression level of miR-22, which downregulates SIRT1 and PPAR $\alpha$  in hepatocytes, is significantly reduced by AM6545 or CB1R ablation (Figure 5). Moreover, the effect of ACEA in reducing the expression of SIRT1 and PPAR $\alpha$  is impaired by a miR-22 inhibitor, while AM6545 failed to reverse their CB1R-induced downregulation in presence of a mimic-miR-22. Finally, the transcription factor p53, known to induce a change in miR expression and to be regulated by CB1R, increases its activity in hepatocytes after ACEA treatment, an effect abolished by AM6545 or genetic ablation of the CB1R. The link between p53 and miR-22 was demonstrated by using DOX, a known activator of p53, which was found to increase the expression of miR-22 while the involvement of p53 in CB1R-induced miR-22 activation was confirmed by using a reversible transcriptional inhibitor.



**Figure 5.** Representative mechanism of PPAR $\alpha$  gene regulation by CB1 receptors. (A) In absence of CB1 ligands, p53 and miRNA22 suppress the activity of SIRT1 which positively regulates PPAR $\alpha$ . (B) In presence of CB1 agonists, p53 and miRNA22 are inhibited and SIRT1 activates PPAR $\alpha$  which forms a heterodimer with RXR, to bind PPARE elements of target genes.

### 7.2. CB1R and PPAR $\gamma$

The endocannabinoid system has a well-established role in the regulation of energy homeostasis through the activation of CB1R [72,73,125]. Numerous studies show that the ECS is overactive in human obesity as well as animal models of genetic and diet-induced obesity [74,126] and thus the pharmacological blockade or genetic ablation of CB1 receptor prevent the risk of obesity and diabetes [127,128]. CB1R is not expressed in undifferentiated preadipocytes but became up-regulated in the early stage of differentiation, while CB2R is expressed in preadipocytes but is down-regulated at an undetectable level upon differentiation [129]. When the cells are differentiated in presence of the PPAR $\gamma$  agonist rosiglitazone, a down-regulation in CB1R and up-regulation in FAAH expression is observed in differentiated adipocytes, suggestive of a reduction in endocannabinoid tone, in

agreement with the ciglitazone-induced decrease of 2-AG levels in partially differentiated adipocytes, observed by Matias et al [130]. On the contrary, the stimulation of preadipocytes in the early phase of adipocyte differentiation with the non-selective cannabinoid agonist WIN55,212 induces a significant expression of PPAR $\gamma$ , an effect no longer observed in fully differentiated cells [129]. The same authors also showed an insulin-mimetic action CB1R-mediated of endocannabinoids on glucose uptake, involving PI3-kinase and the intracellular calcium influx. The increasing of PPAR $\gamma$  expression using WIN55,212 was also observed by Fakhfoury et al. [82], investigating the protective effect of this compound in the beta-amyloid-induced neurodegeneration in rat hippocampus. They observed that the effect on the transcriptional activity of WIN55,212 was partially inhibited with the CB1R antagonist AM251 or the PPAR $\gamma$  antagonist GW9662 but not with the CB2R antagonist SR144528, suggestive of both direct and CB1R-mediated effect on PPAR $\gamma$  signaling [82]. The lipoaminoacid *N*-Oleoylglycine (Olgly) was found to stimulate 3T3-L1 adipogenesis and to increase the expression level of PPAR $\gamma$  through the activation of the CB1 receptor and enhancement of insulin-mediated Akt signaling. Interestingly, Olgly also increases the expression level of CB1R mRNA, whereas the inhibition of CB1R by the antagonist SR141716 abolishes the effects of Olgly on lipid accumulation and PPAR $\gamma$  expression as well as the increment Olgly-dependent of p-Akt/Akt and p-FoxO1/FoxO1 ratio [131], confirming the link between CB1R activation and Akt signaling. Du et al. [132] showed that the 2-AG induced resolution of neuroinflammation in response to pro-inflammatory insults is mediated by the CB1R-dependent expression of PPAR $\gamma$ . In particular, the suppression of NF-kB-p65 phosphorylation, COX-2 expression, and excitatory synaptic transmission in response to proinflammatory IL-1 $\beta$  and LPS were inhibited by the selective PPAR $\gamma$  antagonist GW9662, in hippocampal neurons cultures and the effects of 2-AG were mimicked by the PPAR $\gamma$  agonists 15d-PGJ2 and rosiglitazone. Conversely, Lin et al. [133] demonstrated that in streptozotocin-induced diabetic rats, hyperglycemia causes glomerular hypertrophy and fibrosis with the concomitant increased expression level of inflammatory cytokines and CB1R and reduced PPAR $\gamma$ 2 signaling. Knockdown of PPAR $\gamma$ 2 mimicked the promotional effects of high glucose on CB1R signaling, suggesting that CB1R negatively regulates PPAR $\gamma$ 2 signaling in mesangial cell cultures. Indeed, PPAR $\gamma$ 2 signaling was restored by using CB1R antisense oligonucleotides or inverse agonist AM251. The same effect is obtained with the PPAR $\gamma$  agonist rosiglitazone, which increased PPAR $\gamma$ 2 expression level and counteracted the hyperglycemia-induced enhancement of CB1 expression, inflammation, and glomerular fibrosis in diabetic animals.


### 7.3. CB2R and PPAR $\gamma$ Coactivator 1 $\alpha$ (PGC-1 $\alpha$ )/PPAR $\gamma$

The transcriptional coactivator PGC-1 $\alpha$  enhances the ability of PPARs and other nuclear receptors to promote gene transcription [134] in particular genes linked to fatty acid oxidation, metabolism, and inflammation, and it is upregulated by NAD $^{+}$ -dependent histone deacetylase sirtuin 1 (SIRT1). SIRT1 activity is increased through different pathways, such as glucose restriction, AMP-activated protein kinase (AMPK) activation, and cAMP response element-binding protein (CREB) activation [135,136]. Zheng et al. [137] showed that the selective CB2R agonist trans-caryophyllene, a bicyclic sesquiterpene isolated from cannabis also known as beta-caryophyllene (BCP), stimulates the activity of SIRT1 in skeletal muscle cells by increasing the phosphorylation of CREB, which leads to an enhanced level of PGC-1 $\alpha$  deacetylation and in turn to a higher expression of the genes linked to fatty acid oxidation. The effect of BCP is mediated by CB2R since it is lost when the expression of CB2R in C2C12 myotubes is inhibited with CB2R siRNA. Thus, CB2R contributes to lipid homeostasis by increasing the rate of fatty acid oxidation through the activation of the SIRT1/PGC-1 $\alpha$  pathway. It has been also reported that BCP improves anxiety, memory, and depression by modulating PGC-1 $\alpha$ /BDNF pathway in a CB2R-dependent manner [138]. BCP has been shown to ameliorate arthritis by increasing PGC-1 $\alpha$  and PPAR $\gamma$  expression in human articular chondrocytes and this effect is reverted by the CB2R antagonist AM630 [139]. The modulation of PGC-1 $\alpha$  CB2R-mediated it has been also re-

ported for the CB2R agonist AM1241, which was shown to reduce microglial inflammation by promoting PGC-1 $\alpha$ -induced mitochondrial biogenesis in N9 microglial cells [140].

#### 7.4. CB2R and PPAR $\alpha$

Using a combined pharmacological, biochemical, and bioinformatics approach, it was shown that PEA, as well as the PPAR $\alpha$  selective agonist GW7647, enhances CB2R expression via PPAR $\alpha$  activation in both rat microglia and human macrophage cells [26] (Figure 6). The involvement of PPAR $\alpha$  was demonstrated using either the PPAR $\alpha$  antagonist GW6471, or PPAR $\alpha$  silenced cells, where the incubation with PEA or GW7647 failed to increase CB2R expression. Moreover, it was shown that after stimulation with PEA, PPAR $\alpha$  regulates the transcription of the gene encoding for CB2R *Cnr2* through the high-affinity binding to a specific region in the *Cnr2* gene, identified by a bioinformatic approach.




**Figure 6.** Representative mechanism of PEA and canonical PPAR $\alpha$  agonists in regulating the transcriptional activity of CB2 gene. In presence of a ligand, PPAR $\alpha$  forms a heterodimer with RXR, which binds to the PPRE of the CB2 gene, leading to an increased expression of CB2 protein.

## 8. Crosstalk between PPARs and TRP Channels

### 8.1. PPAR $\alpha$ and TRPV1

Ambrosino et al. [141] have shown that PEA, as well as the canonical PPAR $\alpha$  agonists clofibrate (CLO) and GW7647, induce a PPAR $\alpha$ -dependent activation and desensitization of TRPV1 currents in TRPV1-transfected CHO cells, where PPAR $\alpha$  is endogenously present (Figure 7). While TRPV1 currents induced by capsaicin were unaffected by the PPAR $\alpha$  antagonist GW-6471, those evoked by the PPAR $\alpha$  canonical agonists and PEA were greatly inhibited. The involvement of the TRPV1 channel in the elevation PPAR $\alpha$ -dependent of  $[Ca^{2+}]_i$  was confirmed by the ability of the TRPV1 antagonist capsazepine to counteract the calcium signaling induced by PEA and the other PPAR $\alpha$  agonists in sensory neurons. Later, the same authors confirmed this result using CLO, WY14643 or GW7647, differing each other in structural features, potency and selectivity for PPAR $\alpha$  over the other two PPAR isoforms  $\gamma$  and  $\beta/\delta$ . All these compounds were shown to act as partial TRPV1 agonists in TRPV1-transfected CHO cells, eliciting an efficacy of about half of that of capsaicin [142] and different potency, with CLO and WY14643 resulting about 100-times weaker than GW7647. This rank of potency for TRPV1 activation in vitro is similar to that observed for their analgesic activity in vivo [35]. Moreover, the involvement of PPAR $\alpha$  in TRPV1 activation


by these compounds was shown by down-regulating or up-regulating the levels PPAR $\alpha$  expression: while capsaicin was unaffected by this modulation, a reduction/potentialization of CLO-induced TRPV1 currents was observed in PPAR $\alpha$ -silenced CHO cells and PPAR $\alpha$  over-expressing cells, respectively [142]. Moreover, all the tested compounds desensitized TRPV1 to a greater extent than capsaicin. Evidence of a close association between PPAR $\alpha$  and TRPV1 arises from co-immunoprecipitation experiments performed on total protein lysates from CHO cells co-expressing TRPV1 channels and enhanced GFP (EGFP)-tagged PPAR $\alpha$  receptors, which showed a significant fraction of PPAR $\alpha$  in the anti-TRPV1 purified fraction. Since phosphorylation enhances the function of TRPV1 [143], it is possible to speculate that PPAR $\alpha$  activates and desensitizes TRPV1 by promoting its phosphorylation, similarly to what occurs with the regulation PPAR $\alpha$ -mediated of the nicotine acetylcholine receptors [144].



**Figure 7.** Representative mechanism of TRPV1 receptors regulation by PPAR $\alpha$ . (A) TRPV1 channels stimulation induced by agonists. Phosphorylation of TRPV1 (indicated as P in yellow circles) is known to be a cellular event promoting the channel activity (B) The activation of PPAR $\alpha$  by PEA or canonical agonists, suppresses TRPV1 activity, making the channels refractory to stimuli.

### 8.2. PPAR $\gamma$ and TRPV1

Baskaran et al. demonstrated that dietary capsaicin enhances the expression levels of PPAR $\gamma$  and PGC-1 $\alpha$  (Figure 8) in wt but not in *Trpv1* KO mice (Figure 8). Moreover, it stimulates the SIRT-1-dependent deacetylation of PPAR $\gamma$ , which triggers the browning of white adipose tissue (WAT) and prevents the obesity induced by a high-fat diet in mice [145]. The browning of WAT is a mechanism by which energy expenditure is promoted via thermogenesis. Dietary administration of capsaicin increases the transcription factor PRDM-16 and facilitates its interaction with PPAR $\gamma$  (Figure 8), further contributing to inducing browning of WAT.



**Figure 8.** Representative mechanism of regulation PPAR $\gamma$  by TRPV1. Following the activation by agonists, TRPV1 promotes the activity of PRDM16 which in turn induces the activation of PGC1 $\alpha$ , a major co-activator of the PPAR/RXR complex.

### 8.3. PPAR $\delta$ and TRPV1

The study carried out by Li et al. [146] showed that TRPV1 activation reduces lipid accumulation in hepatocytes by stimulating lipolysis without affecting lipogenesis. This effect occurs through the activation of PPAR $\delta$  since chronic administration of capsaicin promotes the up-regulation of the hepatic PPAR $\delta$  expression level and the enhancement of PPAR $\delta$ -mediated hepatocyte autophagy in wt but not in TRPV1 KO mice, thus disclosing the beneficial role of chronic dietary capsaicin in the prevention of hepatic steatosis, inflammatory responses, and body weight gain. The cross-talk between PPAR $\delta$  and TRPV1 also emerged in a study carried out by Gao et al. [147] where the administration of chronic dietary capsaicin attenuates cardiac hypertrophy and fibrosis associated to with a high-salt diet through the TRPV1-mediated up-regulation of PPAR $\delta$  expression in cardiomyocytes.

## 9. Conclusions

PPARs represent relevant therapeutic targets for neurodegenerative, metabolic, and inflammation-related disorders. On the other hand, several reports have highlighted the therapeutic potential of (endo)cannabinoid molecules for the same diseases. Thus, the ability of such lipid molecules to either directly activate PPARs or modulate them through the cross-talk between PPARs and cannabinoid receptors opens new avenues for the development of synergistic therapeutic approaches. Additionally, there is fascinating evidence that perturbation of the symbiotic gut -microbial community may have a great impact on the host health either directly or indirectly by regulating PPARs activity through new endogenous, endocannabinoid-like modulators. In this context, it has been recently demonstrated that daily ingestion of *Akkermansia muciniphila*, a mucin-degrading bacterium commonly found in the human gut, reduces the risk of developing diet-induced obesity and concomitantly insulin resistance by increasing the plasma levels of mono-palmitoyl-glycerol, an endocannabinoid lipid acting as PPAR $\alpha$  agonist [148]. Conversely, deletion of the gut microbiota in germ-free mice is accompanied by a strong increase in PPAR $\alpha$  expression and PPAR $\alpha$  NAE ligands in the small intestine [149]. Therefore, the identification of

novel modulators and mechanism regulating PPARs activity could pave the way towards future healthcare strategies and therapeutic solutions.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Acknowledgments:** F.A.I. is the recipient of a Duchenne Parent Project NL.

**Conflicts of Interest:** The authors declare that no conflict of interest could be perceived as prejudicing the impartiality of this review.

## References

1. Kliewer, S.A.; Forman, B.M.; Blumberg, B.; Ong, E.S.; Borgmeyer, U.; Mangelsdorf, D.J.; Umesono, K.; Evans, R.M. Differential Expression and Activation of a Family of Murine Peroxisome Proliferator-Activated Receptors. *PNAS* **1994**, *91*, 7355–7359. [[CrossRef](#)] [[PubMed](#)]
2. Zoete, V.; Grosdidier, A.; Michielin, O. Peroxisome Proliferator-Activated Receptor Structures: Ligand Specificity, Molecular Switch and Interactions with Regulators. *Biochim. Biophys. Acta* **2007**, *1771*, 915–925. [[CrossRef](#)] [[PubMed](#)]
3. Kersten, S.; Desvergne, B.; Wahli, W. Roles of PPARs in Health and Disease. *Nature* **2000**, *405*, 421–424. [[CrossRef](#)] [[PubMed](#)]
4. Takeyama, K.; Kodera, Y.; Suzawa, M.; Kato, S. [Peroxisome proliferator-activated receptor(PPAR)–structure, function, tissue distribution, gene expression]. *Nihon Rinsho* **2000**, *58*, 357–363.
5. Moreno, S.; Farioli-Vecchioli, S.; Cerù, M.P. Immunolocalization of Peroxisome Proliferator-Activated Receptors and Retinoid X Receptors in the Adult Rat CNS. *Neuroscience* **2004**, *123*, 131–145. [[CrossRef](#)]
6. Kapadia, R.; Yi, J.-H.; Vemuganti, R. Mechanisms of Anti-Inflammatory and Neuroprotective Actions of PPAR-Gamma Agonists. *Front. Biosci.* **2008**, *13*, 1813–1826. [[CrossRef](#)] [[PubMed](#)]
7. Melis, M.; Carta, S.; Fattore, L.; Tolu, S.; Yasar, S.; Goldberg, S.R.; Fratta, W.; Maskos, U.; Pistis, M. Peroxisome Proliferator-Activated Receptors-Alpha Modulate Dopamine Cell Activity through Nicotinic Receptors. *Biol. Psychiatry* **2010**, *68*, 256–264. [[CrossRef](#)] [[PubMed](#)]
8. Melis, M.; Scheggi, S.; Carta, G.; Madeddu, C.; Lecca, S.; Luchicchi, A.; Cadeddu, F.; Frau, R.; Fattore, L.; Fadda, P.; et al. PPAR $\alpha$  Regulates Cholinergic-Driven Activity of Midbrain Dopamine Neurons via a Novel Mechanism Involving A7 Nicotinic Acetylcholine Receptors. *J. Neurosci.* **2013**, *33*, 6203–6211. [[CrossRef](#)]
9. Wójtowicz, S.; Strosznajder, A.K.; Jeżyna, M.; Strosznajder, J.B. The Novel Role of PPAR Alpha in the Brain: Promising Target in Therapy of Alzheimer’s Disease and Other Neurodegenerative Disorders. *Neurochem. Res.* **2020**, *45*, 972–988. [[CrossRef](#)]
10. Patsouris, D.; Mandard, S.; Voshol, P.J.; Escher, P.; Tan, N.S.; Havekes, L.M.; Koenig, W.; März, W.; Tafuri, S.; Wahli, W.; et al. PPAR $\alpha$  Governs Glycerol Metabolism. *J. Clin. Investig.* **2004**, *114*, 94–103. [[CrossRef](#)] [[PubMed](#)]
11. Peeters, A.; Baes, M. Role of PPAR $\alpha$  in Hepatic Carbohydrate Metabolism. *PPAR Res.* **2010**, *2010*. [[CrossRef](#)]
12. Kersten, S. Integrated Physiology and Systems Biology of PPAR $\alpha$ . *Mol. Metab.* **2014**, *3*, 354–371. [[CrossRef](#)] [[PubMed](#)]
13. Peters, J.M.; Hennuyer, N.; Staels, B.; Fruchart, J.C.; Fievet, C.; Gonzalez, F.J.; Auwerx, J. Alterations in Lipoprotein Metabolism in Peroxisome Proliferator-Activated Receptor Alpha-Deficient Mice. *J. Biol. Chem.* **1997**, *272*, 27307–27312. [[CrossRef](#)]
14. Zhang, J.; Fu, M.; Cui, T.; Xiong, C.; Xu, K.; Zhong, W.; Xiao, Y.; Floyd, D.; Liang, J.; Li, E.; et al. Selective Disruption of PPARgamma 2 Impairs the Development of Adipose Tissue and Insulin Sensitivity. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 10703–10708. [[CrossRef](#)]
15. Guellich, A.; Damy, T.; Lecarpentier, Y.; Conti, M.; Claes, V.; Samuel, J.-L.; Quillard, J.; Hébert, J.-L.; Pineau, T.; Coirault, C. Role of Oxidative Stress in Cardiac Dysfunction of PPARalpha-/- Mice. *Am. J. Physiol. Heart Circ. Physiol.* **2007**, *293*, H93–H102. [[CrossRef](#)] [[PubMed](#)]
16. Lee, S.S.; Pineau, T.; Drago, J.; Lee, E.J.; Owens, J.W.; Kroetz, D.L.; Fernandez-Salguero, P.M.; Westphal, H.; Gonzalez, F.J. Targeted Disruption of the Alpha Isoform of the Peroxisome Proliferator-Activated Receptor Gene in Mice Results in Abolishment of the Pleiotropic Effects of Peroxisome Proliferators. *Mol. Cell Biol.* **1995**, *15*, 3012–3022. [[CrossRef](#)]
17. Leone, T.C.; Weinheimer, C.J.; Kelly, D.P. A Critical Role for the Peroxisome Proliferator-Activated Receptor  $\alpha$  (PPAR $\alpha$ ) in the Cellular Fasting Response: The PPAR $\alpha$ -Null Mouse as a Model of Fatty Acid Oxidation Disorders. *Proc. Natl. Acad. Sci. USA* **1999**, *96*, 7473–7478. [[CrossRef](#)]
18. Hu, Y.; Chen, Y.; Ding, L.; He, X.; Takahashi, Y.; Gao, Y.; Shen, W.; Cheng, R.; Chen, Q.; Qi, X.; et al. Pathogenic Role of Diabetes-Induced PPAR- $\alpha$  down-Regulation in Microvascular Dysfunction. *Proc. Natl. Acad. Sci. USA* **2013**, *110*, 15401–15406. [[CrossRef](#)]
19. Pérez-Martín, M.; Rivera, P.; Blanco, E.; Loreface, C.; Decara, J.; Pavón, F.J.; Serrano, A.; Rodríguez de Fonseca, F.; Suárez, J. Environmental Enrichment, Age, and PPAR $\alpha$  Interact to Regulate Proliferation in Neurogenic Niches. *Front. Neurosci.* **2016**, *10*, 89. [[CrossRef](#)] [[PubMed](#)]
20. Braissant, O.; Fougelle, F.; Scotto, C.; Dauça, M.; Wahli, W. Differential Expression of Peroxisome Proliferator-Activated Receptors (PPARs): Tissue Distribution of PPAR-Alpha, -Beta, and -Gamma in the Adult Rat. *Endocrinology* **1996**, *137*, 354–366. [[CrossRef](#)]

21. Esposito, E.; Impellizzeri, D.; Mazzon, E.; Paterniti, I.; Cuzzocrea, S. Neuroprotective Activities of Palmitoylethanolamide in an Animal Model of Parkinson's Disease. *PLoS ONE* **2012**, *7*. [[CrossRef](#)]
22. D'Agostino, G.; Cristiano, C.; Lyons, D.J.; Citraro, R.; Russo, E.; Avagliano, C.; Russo, R.; Raso, G.M.; Meli, R.; De Sarro, G.; et al. Peroxisome Proliferator-Activated Receptor Alpha Plays a Crucial Role in Behavioral Repetition and Cognitive Flexibility in Mice. *Mol. Metab.* **2015**, *4*, 528–536. [[CrossRef](#)] [[PubMed](#)]
23. Roy, A.; Jana, M.; Corbett, G.T.; Ramaswamy, S.; Kordower, J.H.; Gonzalez, F.J.; Pahan, K. Regulation of Cyclic AMP Response Element Binding and Hippocampal Plasticity-Related Genes by Peroxisome Proliferator-Activated Receptor  $\alpha$ . *Cell Rep.* **2013**, *4*, 724–737. [[CrossRef](#)]
24. Xu, J.; Storer, P.D.; Chavis, J.A.; Racke, M.K.; Drew, P.D. Agonists for the Peroxisome Proliferator-Activated Receptor-Alpha and the Retinoid X Receptor Inhibit Inflammatory Responses of Microglia. *J. Neurosci. Res.* **2005**, *81*, 403–411. [[CrossRef](#)]
25. Bernardo, A.; Bianchi, D.; Magnaghi, V.; Minghetti, L. Peroxisome Proliferator-Activated Receptor-Gamma Agonists Promote Differentiation and Antioxidant Defenses of Oligodendrocyte Progenitor Cells. *J. Neuropathol. Exp. Neurol.* **2009**, *68*, 797–808. [[CrossRef](#)] [[PubMed](#)]
26. Guida, F.; Luongo, L.; Boccella, S.; Giordano, M.E.; Romano, R.; Bellini, G.; Manzo, I.; Furiano, A.; Rizzo, A.; Imperatore, R.; et al. Palmitoylethanolamide Induces Microglia Changes Associated with Increased Migration and Phagocytic Activity: Involvement of the CB2 Receptor. *Sci. Rep.* **2017**, *7*, 375. [[CrossRef](#)] [[PubMed](#)]
27. Moran, E.P.; Ma, J. Therapeutic Effects of PPAR $\alpha$  on Neuronal Death and Microvascular Impairment. Available online: <https://www.hindawi.com/journals/ppar/2015/595426/> (accessed on 24 January 2021).
28. Esposito, E.; Rinaldi, B.; Mazzon, E.; Donniacuo, M.; Impellizzeri, D.; Paterniti, I.; Capuano, A.; Bramanti, P.; Cuzzocrea, S. Anti-Inflammatory Effect of Simvastatin in an Experimental Model of Spinal Cord Trauma: Involvement of PPAR- $\alpha$ . *J. Neuroinflammation* **2012**, *9*, 81. [[CrossRef](#)]
29. D'Orio, B.; Fracassi, A.; Ceru, M.P.; Moreno, S. Targeting PPARalpha in Alzheimer's Disease. *Curr. Alzheimer Res.* **2018**, *15*, 345–354. [[CrossRef](#)]
30. Mirza, A.Z.; Althagafi, I.I.; Shamshad, H. Role of PPAR Receptor in Different Diseases and Their Ligands: Physiological Importance and Clinical Implications. *Eur. J. Med. Chem.* **2019**, *166*, 502–513. [[CrossRef](#)] [[PubMed](#)]
31. Riccardi, L.; Mazzon, E.; Bruscoli, S.; Esposito, E.; Crisafulli, C.; Di Paola, R.; Caminiti, R.; Riccardi, C.; Cuzzocrea, S. Peroxisome Proliferator-Activated Receptor-Alpha Modulates the Anti-Inflammatory Effect of Glucocorticoids in a Model of Inflammatory Bowel Disease in Mice. *Shock* **2009**, *31*, 308–316. [[CrossRef](#)]
32. Genovese, T.; Esposito, E.; Mazzon, E.; Crisafulli, C.; Paterniti, I.; Di Paola, R.; Galuppo, M.; Bramanti, P.; Cuzzocrea, S. PPAR-Alpha Modulate the Anti-Inflammatory Effect of Glucocorticoids in the Secondary Damage in Experimental Spinal Cord Trauma. *Pharmacol. Res.* **2009**, *59*, 338–350. [[CrossRef](#)]
33. Wen, W.; Wang, J.; Zhang, B.; Wang, J. PPAR $\alpha$  Agonist WY-14643 Relieves Neuropathic Pain through SIRT1-Mediated Deacetylation of NF-KB. *PPAR Res.* **2020**, *2020*, 6661642. [[CrossRef](#)]
34. Di Cesare Mannelli, L.; D'Agostino, G.; Pacini, A.; Russo, R.; Zanardelli, M.; Ghelardini, C.; Calignano, A. Palmitoylethanolamide Is a Disease-Modifying Agent in Peripheral Neuropathy: Pain Relief and Neuroprotection Share a PPAR-Alpha-Mediated Mechanism. *Med. Inflamm.* **2013**, *2013*, 328797. [[CrossRef](#)]
35. LoVerme, J.; Russo, R.; La Rana, G.; Fu, J.; Farthing, J.; Mattace-Raso, G.; Meli, R.; Hohmann, A.; Calignano, A.; Piomelli, D. Rapid Broad-Spectrum Analgesia through Activation of Peroxisome Proliferator-Activated Receptor-Alpha. *J. Pharmacol. Exp. Ther.* **2006**, *319*, 1051–1061. [[CrossRef](#)] [[PubMed](#)]
36. Devi, S.; Rangra, N.K.; Rawat, R.; Alrobaian, M.M.; Alam, A.; Singh, R.; Singh, A. Anti-Atherogenic Effect of Nepitrin-7-O-Glucoside: A Flavonoid Isolated from Nepeta Hindostana via Acting on PPAR- $\alpha$  Receptor. *Steroids* **2021**, *165*, 108770. [[CrossRef](#)]
37. Tontonoz, P.; Spiegelman, B.M. Fat and beyond: The Diverse Biology of PPARgamma. *Annu. Rev. Biochem.* **2008**, *77*, 289–312. [[CrossRef](#)] [[PubMed](#)]
38. Nguyen, M.T.A.; Chen, A.; Lu, W.J.; Fan, W.; Li, P.-P.; Oh, D.Y.; Patsouris, D. Regulation of Chemokine and Chemokine Receptor Expression by PPAR $\gamma$  in Adipocytes and Macrophages. *PLoS ONE* **2012**, *7*, e34976. [[CrossRef](#)] [[PubMed](#)]
39. Yang, C.-C.; Wu, C.-H.; Lin, T.-C.; Cheng, Y.-N.; Chang, C.-S.; Lee, K.-T.; Tsai, P.-J.; Tsai, Y.-S. Inhibitory Effect of PPAR $\gamma$  on NLRP3 Inflammasome Activation. *Theranostics* **2021**, *11*, 2424–2441. [[CrossRef](#)]
40. Gray, S.L.; Dalla Nora, E.; Vidal-Puig, A.J. Mouse Models of PPAR-Gamma Deficiency: Dissecting PPAR-Gamma's Role in Metabolic Homeostasis. *Biochem. Soc. Trans.* **2005**, *33*, 1053–1058. [[CrossRef](#)] [[PubMed](#)]
41. Gavrilova, O.; Haluzik, M.; Matsusue, K.; Cutson, J.J.; Johnson, L.; Dietz, K.R.; Nicol, C.J.; Vinson, C.; Gonzalez, F.J.; Reitman, M.L. Liver Peroxisome Proliferator-Activated Receptor  $\gamma$  Contributes to Hepatic Steatosis, Triglyceride Clearance, and Regulation of Body Fat Mass. *J. Biol. Chem.* **2003**, *278*, 34268–34276. [[CrossRef](#)]
42. Kintscher, U.; Law, R.E. PPARgamma-Mediated Insulin Sensitization: The Importance of Fat versus Muscle. *Am. J. Physiol. Endocrinol. Metab.* **2005**, *288*, E287–E291. [[CrossRef](#)]
43. Amin, R.H.; Mathews, S.T.; Camp, H.S.; Ding, L.; Leff, T. Selective Activation of PPARgamma in Skeletal Muscle Induces Endogenous Production of Adiponectin and Protects Mice from Diet-Induced Insulin Resistance. *Am. J. Physiol. Endocrinol. Metab.* **2010**, *298*, E28–E37. [[CrossRef](#)]

44. Gilardi, F.; Winkler, C.; Quignodon, L.; Diserens, J.-G.; Toffoli, B.; Schiffrin, M.; Sardella, C.; Preitner, F.; Desvergne, B. Systemic PPAR $\gamma$  Deletion in Mice Provokes Lipoatrophy, Organomegaly, Severe Type 2 Diabetes and Metabolic Inflexibility. *Metabolism* **2019**, *95*, 8–20. [[CrossRef](#)]
45. Lüdtke, A.; Buettner, J.; Schmidt, H.H.; Worman, H.J. New PPAR $\gamma$  Mutation Leads to Lipodystrophy and Loss of Protein Function That Is Partially Restored by a Synthetic Ligand. *J. Med. Genet.* **2007**, *44*, e88. [[CrossRef](#)]
46. Fajas, L.; Auboeuf, D.; Raspé, E.; Schoonjans, K.; Lefebvre, A.-M.; Saladin, R.; Najib, J.; Laville, M.; Fruchart, J.-C.; Deeb, S.; et al. The Organization, Promoter Analysis, and Expression of the Human PPAR $\gamma$  Gene. *J. Biol. Chem.* **1997**, *272*, 18779–18789. [[CrossRef](#)] [[PubMed](#)]
47. Zhou, J.; Wilson, K.M.; Medh, J.D. Genetic Analysis of Four Novel Peroxisome Proliferator Activated Receptor-Gamma Splice Variants in Monkey Macrophages. *Biochem. Biophys. Res. Commun.* **2002**, *293*, 274–283. [[CrossRef](#)]
48. Subbarayan, V.; Sabichi, A.L.; Kim, J.; Llansa, N.; Logothetis, C.J.; Lippman, S.M.; Menter, D.G. Differential Peroxisome Proliferator-Activated Receptor-Gamma Isoform Expression and Agonist Effects in Normal and Malignant Prostate Cells. *Cancer Epidemiol. Biomark. Prev.* **2004**, *13*, 1710–1716. [[PubMed](#)]
49. Chen, Y.; Jimenez, A.R.; Medh, J.D. Identification and Regulation of Novel PPAR-Gamma Splice Variants in Human THP-1 Macrophages. *Biochim. Biophys. Acta* **2006**, *1759*, 32–43. [[CrossRef](#)] [[PubMed](#)]
50. Daynes, R.A.; Jones, D.C. Emerging Roles of PPARs in Inflammation and Immunity. *Nat. Rev. Immunol.* **2002**, *2*, 748–759. [[CrossRef](#)]
51. Kiaei, M.; Kipiani, K.; Chen, J.; Calingasan, N.Y.; Beal, M.F. Peroxisome Proliferator-Activated Receptor-Gamma Agonist Extends Survival in Transgenic Mouse Model of Amyotrophic Lateral Sclerosis. *Exp. Neurol.* **2005**, *191*, 331–336. [[CrossRef](#)]
52. Heneka, M.T.; Landreth, G.E. PPARs in the Brain. *Biochim. Biophys. Acta (BBA) Mol. Cell Biol. Lipids* **2007**, *1771*, 1031–1045. [[CrossRef](#)]
53. Roudsari, N.M.; Lashgari, N.-A.; Zandi, N.; Pazoki, B.; Momtaz, S.; Sahebkar, A.; Abdolghaffari, A.H. PPAR $\gamma$ : A Turning Point for Irritable Bowel Syndrome Treatment. *Life Sci.* **2020**, *257*, 118103. [[CrossRef](#)] [[PubMed](#)]
54. Tachibana, M.; Wada, K.; Katayama, K.; Kamisaki, Y.; Maeyama, K.; Kadowaki, T.; Blumberg, R.S.; Nakajima, A. Activation of Peroxisome Proliferator-Activated Receptor Gamma Suppresses Mast Cell Maturation Involved in Allergic Diseases. *Allergy* **2008**, *63*, 1136–1147. [[CrossRef](#)] [[PubMed](#)]
55. Ramot, Y.; Mastrofrancesco, A.; Camera, E.; Desreumaux, P.; Paus, R.; Picardo, M. The Role of PPAR $\gamma$ -Mediated Signalling in Skin Biology and Pathology: New Targets and Opportunities for Clinical Dermatology. *Exp. Dermatol.* **2015**, *24*, 245–251. [[CrossRef](#)] [[PubMed](#)]
56. Warden, A.; Truitt, J.; Merriman, M.; Ponomareva, O.; Jameson, K.; Ferguson, L.B.; Mayfield, R.D.; Harris, R.A. Localization of PPAR Isotypes in the Adult Mouse and Human Brain. *Sci. Rep.* **2016**, *6*. [[CrossRef](#)]
57. Strosznajder, A.K.; Wójtowicz, S.; Jeżyna, M.J.; Sun, G.Y.; Strosznajder, J.B. Recent Insights on the Role of PPAR- $\beta/\delta$  in Neuroinflammation and Neurodegeneration, and Its Potential Target for Therapy. *Neuromol. Med.* **2020**. [[CrossRef](#)]
58. Luquet, S.; Gaudel, C.; Holst, D.; Lopez-Soriano, J.; Jehl-Pietri, C.; Fredenrich, A.; Grimaldi, P.A. Roles of PPAR Delta in Lipid Absorption and Metabolism: A New Target for the Treatment of Type 2 Diabetes. *Biochim. Biophys. Acta (BBA) Mol. Cell Biol. Lipids* **2005**, *1740*, 313–317. [[CrossRef](#)] [[PubMed](#)]
59. Wagner, N.; Wagner, K.-D. PPAR Beta/Delta and the Hallmarks of Cancer. *Cells* **2020**, *9*, 1133. [[CrossRef](#)] [[PubMed](#)]
60. Hall, M.G.; Quignodon, L.; Desvergne, B. Peroxisome Proliferator-Activated Receptor Beta/Delta in the Brain: Facts and Hypothesis. *PPAR Res.* **2008**, *2008*, 780452. [[CrossRef](#)] [[PubMed](#)]
61. Müller, R.; Rieck, M.; Müller-Brüsselbach, S. Regulation of Cell Proliferation and Differentiation by PPAR $\beta/\delta$ . *PPAR Res.* **2008**, *2008*. [[CrossRef](#)]
62. Barz, T.; Spengler, D. Peroxisome Proliferator-Activated Receptor. In *Encyclopedia of Cancer*; Schwab, M., Ed.; Springer: Berlin, Heidelberg, 2016; pp. 1–5. ISBN 978-3-642-27841-9.
63. Matsushita, Y.; Ogawa, D.; Wada, J.; Yamamoto, N.; Shikata, K.; Sato, C.; Tachibana, H.; Toyota, N.; Makino, H. Activation of Peroxisome Proliferator-Activated Receptor Delta Inhibits Streptozotocin-Induced Diabetic Nephropathy through Anti-Inflammatory Mechanisms in Mice. *Diabetes* **2011**, *60*, 960–968. [[CrossRef](#)] [[PubMed](#)]
64. Cheang, W.S.; Wong, W.T.; Zhao, L.; Xu, J.; Wang, L.; Lau, C.W.; Chen, Z.Y.; Ma, R.C.W.; Xu, A.; Wang, N.; et al. PPAR $\delta$  Is Required for Exercise to Attenuate Endoplasmic Reticulum Stress and Endothelial Dysfunction in Diabetic Mice. *Diabetes* **2017**, *66*, 519–528. [[CrossRef](#)] [[PubMed](#)]
65. Liu, Y.; Colby, J.K.; Zuo, X.; Jaoude, J.; Wei, D.; Shureiqi, I. The Role of PPAR- $\delta$  in Metabolism, Inflammation, and Cancer: Many Characters of a Critical Transcription Factor. *Int. J. Mol. Sci.* **2018**, *19*, 3339. [[CrossRef](#)] [[PubMed](#)]
66. O’Sullivan, S.E. Cannabinoids Go Nuclear: Evidence for Activation of Peroxisome Proliferator-Activated Receptors. *Br. J. Pharmacol.* **2007**, *152*, 576–582. [[CrossRef](#)] [[PubMed](#)]
67. Kroker, A.J.; Bruning, J.B. Review of the Structural and Dynamic Mechanisms of PPAR $\gamma$  Partial Agonism. *PPAR Res.* **2015**, *2015*, 816856. [[CrossRef](#)]
68. Evans, R.M.; Mangelsdorf, D.J. Nuclear Receptors, RXR, and the Big Bang. *Cell* **2014**, *157*, 255–266. [[CrossRef](#)]
69. Viswakarma, N.; Jia, Y.; Bai, L.; Vluggens, A.; Borensztajn, J.; Xu, J.; Reddy, J.K. Coactivators in PPAR-Regulated Gene Expression. *PPAR Res.* **2010**, *2010*. [[CrossRef](#)]

70. Bruning, J.B.; Chalmers, M.J.; Prasad, S.; Busby, S.A.; Kamenecka, T.M.; He, Y.; Nettles, K.W.; Griffin, P.R. Partial Agonists Activate PPAR $\gamma$  Using a Helix 12 Independent Mechanism. *Structure* **2007**, *15*, 1258–1271. [[CrossRef](#)]
71. Brunmeir, R.; Xu, F. Functional Regulation of PPARs through Post-Translational Modifications. *Int. J. Mol. Sci.* **2018**, *19*, 1738. [[CrossRef](#)]
72. Di Marzo, V. Endocannabinoids: Synthesis and Degradation. *Rev. Physiol. Biochem. Pharmacol.* **2008**, *160*, 1–24. [[CrossRef](#)]
73. Iannotti, F.A.; Di Marzo, V.; Petrosino, S. Endocannabinoids and Endocannabinoid-Related Mediators: Targets, Metabolism and Role in Neurological Disorders. *Prog. Lipid Res.* **2016**, *62*, 107–128. [[CrossRef](#)]
74. Iannotti, F.A.; Di Marzo, V. The Gut Microbiome, Endocannabinoids and Metabolic Disorders. *J Endocrinol* **2020**. [[CrossRef](#)] [[PubMed](#)]
75. Sigel, E.; Baur, R.; Rácz, I.; Marazzi, J.; Smart, T.G.; Zimmer, A.; Gertsch, J. The Major Central Endocannabinoid Directly Acts at GABA(A) Receptors. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 18150–18155. [[CrossRef](#)]
76. Tenenbaum, A.; Motro, M.; Fisman, E.Z. Dual and Pan-Peroxisome Proliferator-Activated Receptors (PPAR) Co-Agonism: The Bezafibrate Lessons. *Cardiovasc. Diabetol.* **2005**, *4*, 14. [[CrossRef](#)]
77. Adeghate, E.; Adem, A.; Hasan, M.Y.; Tekes, K.; Kalasz, H. Medicinal Chemistry and Actions of Dual and Pan PPAR Modulators. *Open Med. Chem. J.* **2011**, *5*, 93–98. [[CrossRef](#)] [[PubMed](#)]
78. Wallenius, K.; Kjellstedt, A.; Thalén, P.; Löfgren, L.; Oakes, N.D. The PPAR $\alpha/\gamma$  Agonist, Tesaglitazar, Improves Insulin Mediated Switching of Tissue Glucose and Free Fatty Acid Utilization In Vivo in the Obese Zucker Rat. *PPAR Res.* **2013**, *2013*. [[CrossRef](#)] [[PubMed](#)]
79. Mansour, M. Chapter Seven—The Roles of Peroxisome Proliferator-Activated Receptors in the Metabolic Syndrome. In *Progress in Molecular Biology and Translational Science; Glucose Homeostasis and the Pathogenesis of Diabetes Mellitus*; Tao, Y.-X., Ed.; Academic Press; Volume 121, pp. 217–266.
80. Yu, X.-H.; Zheng, X.-L.; Tang, C.-K. Peroxisome Proliferator-Activated Receptor  $\alpha$  in Lipid Metabolism and Atherosclerosis. *Adv. Clin. Chem.* **2015**, *71*, 171–203. [[CrossRef](#)]
81. Esposito, E.; Cuzzocrea, S. Targeting the Peroxisome Proliferator-Activated Receptors (PPARs) in Spinal Cord Injury. *Expert Opin. Ther. Targets* **2011**, *15*, 943–959. [[CrossRef](#)]
82. Fakhfour, G.; Ahmadiani, A.; Rahimian, R.; Grolla, A.A.; Moradi, F.; Haeri, A. WIN55212-2 Attenuates Amyloid-Beta-Induced Neuroinflammation in Rats through Activation of Cannabinoid Receptors and PPAR- $\gamma$  Pathway. *Neuropharmacology* **2012**, *63*, 653–666. [[CrossRef](#)]
83. González-Mateo, G.T.; Aroeira, L.S.; López-Cabrera, M.; Ruiz-Ortega, M.; Ortiz, A.; Selgas, R. Pharmacological Modulation of Peritoneal Injury Induced by Dialysis Fluids: Is It an Option? *Nephrol. Dial. Trans.* **2012**, *27*, 478–481. [[CrossRef](#)]
84. Scuderi, C.; Steardo, L.; Esposito, G. Cannabidiol Promotes Amyloid Precursor Protein Ubiquitination and Reduction of Beta Amyloid Expression in SHSY5YAPP+ Cells through PPAR $\gamma$  Involvement. *Phytother. Res.* **2014**, *28*, 1007–1013. [[CrossRef](#)] [[PubMed](#)]
85. Liu, J.; Li, H.; Burstein, S.H.; Zurier, R.B.; Chen, J.D. Activation and Binding of Peroxisome Proliferator-Activated Receptor Gamma by Synthetic Cannabinoid Ajulemic Acid. *Mol. Pharmacol.* **2003**, *63*, 983–992. [[CrossRef](#)] [[PubMed](#)]
86. Granja, A.G.; Carrillo-Salinas, F.; Pagani, A.; Gómez-Cañas, M.; Negri, R.; Navarrete, C.; Mecha, M.; Mestre, L.; Fiebich, B.L.; Cantarero, I.; et al. A Cannabigerol Quinone Alleviates Neuroinflammation in a Chronic Model of Multiple Sclerosis. *J. Neuroimmune Pharmacol.* **2012**, *7*, 1002–1016. [[CrossRef](#)]
87. O’Sullivan, S.E. An Update on PPAR Activation by Cannabinoids. *Br. J. Pharmacol.* **2016**, *173*, 1899–1910. [[CrossRef](#)]
88. Burgaz, S.; García, C.; Gómez-Cañas, M.; Navarrete, C.; García-Martín, A.; Rolland, A.; Del Río, C.; Casarejos, M.J.; Muñoz, E.; Gonzalo-Consuegra, C.; et al. Neuroprotection with the Cannabigerol Quinone Derivative VCE-003.2 and Its Analogs CBGA-Q and CBGA-Q-Salt in Parkinson’s Disease Using 6-Hydroxydopamine-Lesioned Mice. *Mol. Cell Neurosci.* **2020**, *110*, 103583. [[CrossRef](#)] [[PubMed](#)]
89. Caprioglio, D.; Mattoteia, D.; Pollastro, F.; Negri, R.; Lopatriello, A.; Chianese, G.; Minassi, A.; Collado, J.A.; Munoz, E.; Tagliatela-Scafati, O.; et al. The Oxidation of Phytocannabinoids to Cannabinoquinoids. *J. Nat. Prod.* **2020**, *83*, 1711–1715. [[CrossRef](#)] [[PubMed](#)]
90. Iannotti, F.A.; De Maio, F.; Panza, E.; Appendino, G.; Tagliatela-Scafati, O.; De Petrocellis, L.; Amodeo, P.; Vitale, R.M. Identification and Characterization of Cannabimovone, a Cannabinoid from Cannabis Sativa, as a Novel PPAR $\gamma$  Agonist via a Combined Computational and Functional Study. *Molecules* **2020**, *25*, 1119. [[CrossRef](#)]
91. D’Aniello, E.; Fellous, T.; Iannotti, F.A.; Gentile, A.; Allarà, M.; Balestrieri, F.; Gray, R.; Amodeo, P.; Vitale, R.M.; Di Marzo, V. Identification and Characterization of Phytocannabinoids as Novel Dual PPAR $\alpha/\gamma$  Agonists by a Computational and in Vitro Experimental Approach. *Biochim. Biophys. Acta Gen. Subj.* **2019**, *1863*, 586–597. [[CrossRef](#)] [[PubMed](#)]
92. Sun, Y.; Bennett, A. Cannabinoids: A New Group of Agonists of PPARs. *PPAR Res.* **2007**, *2007*, 23513. [[CrossRef](#)]
93. Bouaboula, M.; Hilairet, S.; Marchand, J.; Fajas, L.; Le Fur, G.; Casellas, P. Anandamide Induced PPAR $\gamma$  Transcriptional Activation and 3T3-L1 Preadipocyte Differentiation. *Eur. J. Pharmacol.* **2005**, *517*, 174–181. [[CrossRef](#)]
94. Fu, J.; Gaetani, S.; Oveisi, F.; Lo Verme, J.; Serrano, A.; Rodríguez de Fonseca, F.; Rosengarth, A.; Luecke, H.; Di Giacomo, B.; Tarzia, G.; et al. Oleyethanolamide Regulates Feeding and Body Weight through Activation of the Nuclear Receptor PPAR- $\alpha$ . *Nature* **2003**, *425*, 90–93. [[CrossRef](#)]

95. Guzmán, M.; Lo Verme, J.; Fu, J.; Oveisi, F.; Blázquez, C.; Piomelli, D. Oleoylethanolamide Stimulates Lipolysis by Activating the Nuclear Receptor Peroxisome Proliferator-Activated Receptor Alpha (PPAR-Alpha). *J. Biol. Chem.* **2004**, *279*, 27849–27854. [[CrossRef](#)] [[PubMed](#)]
96. LoVerme, J.; La Rana, G.; Russo, R.; Calignano, A.; Piomelli, D. The Search for the Palmitoylethanolamide Receptor. *Life Sci.* **2005**, *77*, 1685–1698. [[CrossRef](#)] [[PubMed](#)]
97. Artmann, A.; Petersen, G.; Hellgren, L.I.; Boberg, J.; Skonberg, C.; Nellemann, C.; Hansen, S.H.; Hansen, H.S. Influence of Dietary Fatty Acids on Endocannabinoid and N-Acylethanolamine Levels in Rat Brain, Liver and Small Intestine. *Biochim. Biophys. Acta* **2008**, *1781*, 200–212. [[CrossRef](#)] [[PubMed](#)]
98. Y, S.; Sp, A.; Mj, G.; Cl, G.; K, H.; Sp, M.; Da, K.; Aj, B. Cannabinoid Activation of PPAR Alpha; a Novel Neuroprotective Mechanism. *Br. J. Pharmacol.* **2007**, *152*, 734–743. [[CrossRef](#)]
99. Rockwell, C.E.; Snider, N.T.; Thompson, J.T.; Vanden Heuvel, J.P.; Kaminski, N.E. Interleukin-2 Suppression by 2-Arachidonyl Glycerol Is Mediated through Peroxisome Proliferator-Activated Receptor Gamma Independently of Cannabinoid Receptors 1 and 2. *Mol. Pharmacol.* **2006**, *70*, 101–111. [[CrossRef](#)]
100. Raman, P.; Kaplan, B.L.F.; Thompson, J.T.; Vanden Heuvel, J.P.; Kaminski, N.E. 15-Deoxy-Delta12,14-Prostaglandin J2-Glycerol Ester, a Putative Metabolite of 2-Arachidonyl Glycerol, Activates Peroxisome Proliferator Activated Receptor Gamma. *Mol. Pharmacol.* **2011**, *80*, 201–209. [[CrossRef](#)]
101. Kozak, K.R.; Gupta, R.A.; Moody, J.S.; Ji, C.; Boeglin, W.E.; DuBois, R.N.; Brash, A.R.; Marnett, L.J. 15-Lipoxygenase Metabolism of 2-Arachidonylglycerol. Generation of a Peroxisome Proliferator-Activated Receptor Alpha Agonist. *J. Biol. Chem.* **2002**, *277*, 23278–23286. [[CrossRef](#)]
102. Sun, L.; Xu, Y.-W.; Han, J.; Liang, H.; Wang, N.; Cheng, Y. 12/15-Lipoxygenase Metabolites of Arachidonic Acid Activate PPAR $\gamma$ : A Possible Neuroprotective Effect in Ischemic Brain. *J. Lipid Res.* **2015**, *56*, 502–514. [[CrossRef](#)]
103. Yu, S.; Levi, L.; Casadesus, G.; Kunos, G.; Noy, N. Fatty Acid-Binding Protein 5 (FABP5) Regulates Cognitive Function Both by Decreasing Anandamide Levels and by Activating the Nuclear Receptor Peroxisome Proliferator-Activated Receptor  $\beta/\delta$  (PPAR $\beta/\delta$ ) in the Brain. *J. Biol. Chem.* **2014**, *289*, 12748–12758. [[CrossRef](#)]
104. Dionisi, M.; Alexander, S.P.; Bennett, A.J. Oleamide Activates Peroxisome Proliferator-Activated Receptor Gamma (PPAR $\gamma$ ) in Vitro. *Lipids Health Dis.* **2012**, *11*, 51. [[CrossRef](#)]
105. Pistis, M.; O'Sullivan, S.E. The Role of Nuclear Hormone Receptors in Cannabinoid Function. *Adv. Pharmacol.* **2017**, *80*, 291–328. [[CrossRef](#)]
106. Mazzola, C.; Medalie, J.; Scherma, M.; Panlilio, L.V.; Solinas, M.; Tanda, G.; Drago, F.; Cadet, J.L.; Goldberg, S.R.; Yasar, S. Fatty Acid Amide Hydrolase (FAAH) Inhibition Enhances Memory Acquisition through Activation of PPAR-Alpha Nuclear Receptors. *Learn. Mem.* **2009**, *16*, 332–337. [[CrossRef](#)] [[PubMed](#)]
107. Taylor, B.K. N-Acylethanolamine Acid Amidase (NAAA), a New Path to Unleash PPAR-Mediated Analgesia. *Pain* **2013**, *154*, 326–327. [[CrossRef](#)] [[PubMed](#)]
108. Di Marzo, V.; Melck, D.; Orlando, P.; Bisogno, T.; Zagoory, O.; Bifulco, M.; Vogel, Z.; De Petrocellis, L. Palmitoylethanolamide Inhibits the Expression of Fatty Acid Amide Hydrolase and Enhances the Anti-Proliferative Effect of Anandamide in Human Breast Cancer Cells. *Biochem. J.* **2001**, *358*, 249–255. [[CrossRef](#)]
109. Han, S.; Inoue, H.; Flowers, L.C.; Sidell, N. Control of COX-2 Gene Expression through Peroxisome Proliferator-Activated Receptor Gamma in Human Cervical Cancer Cells. *Clin. Cancer Res.* **2003**, *9*, 4627–4635.
110. Knopfová, L.; Šmarda, J. The Use of Cox-2 and PPAR $\gamma$  Signaling in Anti-Cancer Therapies (Review). *Exp. Ther. Med.* **2010**, *1*, 257–264. [[CrossRef](#)] [[PubMed](#)]
111. Subbarayan, V.; Xu, X.C.; Kim, J.; Yang, P.; Hoque, A.; Sabichi, A.L.; Llansa, N.; Mendoza, G.; Logothetis, C.J.; Newman, R.A.; et al. Inverse Relationship between 15-Lipoxygenase-2 and PPAR- $\gamma$  Gene Expression in Normal Epithelia Compared with Tumor Epithelia. *Neoplasia* **2005**, *7*, 280–293. [[CrossRef](#)]
112. Gotoh, C.; Hong, Y.-H.; Iga, T.; Hishikawa, D.; Suzuki, Y.; Song, S.-H.; Choi, K.-C.; Adachi, T.; Hirasawa, A.; Tsujimoto, G.; et al. The Regulation of Adipogenesis through GPR120. *Biochem. Biophys. Res. Commun.* **2007**, *354*, 591–597. [[CrossRef](#)]
113. Ichimura, A.; Hara, T.; Hirasawa, A. Regulation of Energy Homeostasis via GPR120. *Front. Endocrinol.* **2014**, *5*, 111. [[CrossRef](#)]
114. Moniri, N.H. Free-Fatty Acid Receptor-4 (GPR120): Cellular and Molecular Function and Its Role in Metabolic Disorders. *Biochem. Pharmacol.* **2016**, *110–111*, 1–15. [[CrossRef](#)]
115. Song, T.; Yang, Y.; Zhou, Y.; Wei, H.; Peng, J. GPR120: A Critical Role in Adipogenesis, Inflammation, and Energy Metabolism in Adipose Tissue. *Cell. Mol. Life Sci.* **2017**, *74*, 2723–2733. [[CrossRef](#)] [[PubMed](#)]
116. Deckelbaum, R.J.; Worgall, T.S.; Seo, T. N-3 Fatty Acids and Gene Expression. *Am. J. Clin. Nutr.* **2006**, *83*, 1520S–1525S. [[CrossRef](#)] [[PubMed](#)]
117. Grygiel-Górniak, B. Peroxisome Proliferator-Activated Receptors and Their Ligands: Nutritional and Clinical Implications—A Review. *Nutr. J.* **2014**, *13*, 17. [[CrossRef](#)] [[PubMed](#)]
118. Hasan, A.U.; Ohmori, K.; Konishi, K.; Igarashi, J.; Hashimoto, T.; Kamitori, K.; Yamaguchi, F.; Tsukamoto, I.; Uyama, T.; Ishihara, Y.; et al. Eicosapentaenoic Acid Upregulates VEGF-A through Both GPR120 and PPAR $\gamma$  Mediated Pathways in 3T3-L1 Adipocytes. *Mol. Cell Endocrinol.* **2015**, *406*, 10–18. [[CrossRef](#)]
119. Bae, I.-S.; Park, P.J.; Lee, J.H.; Cho, E.-G.; Lee, T.R.; Kim, S.H. PPAR $\gamma$ -Mediated G-Protein Coupled Receptor 120 Signaling Pathway Promotes Transcriptional Activation of MiR-143 in Adipocytes. *Gene* **2017**, *626*, 64–69. [[CrossRef](#)] [[PubMed](#)]

120. Jeninga, E.H.; Bugge, A.; Nielsen, R.; Kersten, S.; Hamers, N.; Dani, C.; Wabitsch, M.; Berger, R.; Stunnenberg, H.G.; Mandrup, S.; et al. Peroxisome Proliferator-Activated Receptor  $\gamma$  Regulates Expression of the Anti-Lipolytic G-Protein-Coupled Receptor 81 (GPR81/Gpr81). *J. Biol. Chem.* **2009**, *284*, 26385–26393. [CrossRef]
121. Knowles, H.J.; te Poele, R.H.; Te Poole, R.; Workman, P.; Harris, A.L. Niacin Induces PPAR $\gamma$  Expression and Transcriptional Activation in Macrophages via HM74 and HM74a-Mediated Induction of Prostaglandin Synthesis Pathways. *Biochem. Pharmacol.* **2006**, *71*, 646–656. [CrossRef]
122. Kim, H.-S.; Hwang, Y.-C.; Koo, S.-H.; Park, K.S.; Lee, M.-S.; Kim, K.-W.; Lee, M.-K. PPAR- $\gamma$  Activation Increases Insulin Secretion through the Up-Regulation of the Free Fatty Acid Receptor GPR40 in Pancreatic  $\beta$ -Cells. *PLoS ONE* **2013**, *8*, e50128. [CrossRef]
123. Richard, C.L.; Blay, J. CXCR4 in Cancer and Its Regulation by PPAR $\gamma$ . Available online: <https://www.hindawi.com/journals/ppar/2008/769413/> (accessed on 12 January 2021).
124. Azar, S.; Udi, S.; Drori, A.; Hadar, R.; Nemirovski, A.; Vemuri, K.V.; Miller, M.; Sherill-Rofe, D.; Arad, Y.; Gur-Wahnon, D.; et al. Reversal of Diet-Induced Hepatic Steatosis by Peripheral CB1 Receptor Blockade in Mice Is P53/MiRNA-22/SIRT1/PPAR $\alpha$  Dependent. *Mol. Metab.* **2020**, *42*, 101087. [CrossRef]
125. Di Marzo, V. New Approaches and Challenges to Targeting the Endocannabinoid System. *Nat. Rev. Drug Discov.* **2018**, *17*, 623–639. [CrossRef] [PubMed]
126. Silvestri, C.; Di Marzo, V. The Endocannabinoid System in Energy Homeostasis and the Etiopathology of Metabolic Disorders. *Cell Metab.* **2013**, *17*, 475–490. [CrossRef] [PubMed]
127. Ravinet Trillou, C.; Delgorge, C.; Menet, C.; Arnone, M.; Soubrié, P. CB1 Cannabinoid Receptor Knockout in Mice Leads to Leanness, Resistance to Diet-Induced Obesity and Enhanced Leptin Sensitivity. *Int. J. Obes. Relat. Metab. Disord.* **2004**, *28*, 640–648. [CrossRef]
128. Després, J.-P.; Golay, A.; Sjöström, L.; Rimonabant in Obesity-Lipids Study Group. Effects of Rimonabant on Metabolic Risk Factors in Overweight Patients with Dyslipidemia. *N. Engl. J. Med.* **2005**, *353*, 2121–2134. [CrossRef]
129. Pagano, C.; Pilon, C.; Calcagno, A.; Urbanet, R.; Rossato, M.; Milan, G.; Bianchi, K.; Rizzuto, R.; Bernante, P.; Federspil, G.; et al. The Endogenous Cannabinoid System Stimulates Glucose Uptake in Human Fat Cells via Phosphatidylinositol 3-Kinase and Calcium-Dependent Mechanisms. *J. Clin. Endocrinol. Metab.* **2007**, *92*, 4810–4819. [CrossRef]
130. Matias, I.; Gonthier, M.-P.; Orlando, P.; Martiadis, V.; De Petrocellis, L.; Cervino, C.; Petrosino, S.; Hoareau, L.; Festy, F.; Pasquali, R.; et al. Regulation, Function, and Dysregulation of Endocannabinoids in Models of Adipose and Beta-Pancreatic Cells and in Obesity and Hyperglycemia. *J. Clin. Endocrinol. Metab.* **2006**, *91*, 3171–3180. [CrossRef]
131. Wang, S.; Xu, Q.; Shu, G.; Wang, L.; Gao, P.; Xi, Q.; Zhang, Y.; Jiang, Q.; Zhu, X. N-Oleoyl Glycine, a Lipoamino Acid, Stimulates Adipogenesis Associated with Activation of CB1 Receptor and Akt Signaling Pathway in 3T3-L1 Adipocyte. *Biochem. Biophys. Res. Commun.* **2015**, *466*, 438–443. [CrossRef] [PubMed]
132. Du, H.; Chen, X.; Zhang, J.; Chen, C. Inhibition of COX-2 Expression by Endocannabinoid 2-Arachidonoylglycerol Is Mediated via PPAR- $\gamma$ . *Br. J. Pharmacol.* **2011**, *163*, 1533–1549. [CrossRef] [PubMed]
133. Lin, C.-L.; Hsu, Y.-C.; Lee, P.-H.; Lei, C.-C.; Wang, J.-Y.; Huang, Y.-T.; Wang, S.-Y.; Wang, F.-S. Cannabinoid Receptor 1 Disturbance of PPAR $\gamma$ 2 Augments Hyperglycemia Induction of Mesangial Inflammation and Fibrosis in Renal Glomeruli. *J. Mol. Med.* **2014**, *92*, 779–792. [CrossRef]
134. Scarpulla, R.C.; Vega, R.B.; Kelly, D.P. Transcriptional Integration of Mitochondrial Biogenesis. *Trends Endocrinol. Metab.* **2012**, *23*, 459–466. [CrossRef] [PubMed]
135. Rodgers, J.T.; Lerin, C.; Haas, W.; Gygi, S.P.; Spiegelman, B.M.; Puigserver, P. Nutrient Control of Glucose Homeostasis through a Complex of PGC-1 $\alpha$  and SIRT1. *Nature* **2005**, *434*, 113–118. [CrossRef]
136. Gerhart-Hines, Z.; Dominy, J.E.; Blättler, S.M.; Jedrychowski, M.P.; Banks, A.S.; Lim, J.-H.; Chim, H.; Gygi, S.P.; Puigserver, P. The CAMP/PKA Pathway Rapidly Activates SIRT1 to Promote Fatty Acid Oxidation Independently of Changes in NAD(+). *Mol. Cell* **2011**, *44*, 851–863. [CrossRef] [PubMed]
137. Zheng, X.; Sun, T.; Wang, X. Activation of Type 2 Cannabinoid Receptors (CB2R) Promotes Fatty Acid Oxidation through the SIRT1/PGC-1 $\alpha$  Pathway. *Biochem. Biophys. Res. Commun.* **2013**, *436*, 377–381. [CrossRef]
138. Youssef, D.A.; El-Fayoumi, H.M.; Mahmoud, M.F. Beta-Caryophyllene Alleviates Diet-Induced Neurobehavioral Changes in Rats: The Role of CB2 and PPAR- $\gamma$  Receptors. *Biomed. Pharmacother.* **2019**, *110*, 145–154. [CrossRef]
139. Irrera, N.; D’Ascola, A.; Pallio, G.; Bitto, A.; Mazzon, E.; Mannino, F.; Squadrito, V.; Arcoraci, V.; Minutoli, L.; Campo, G.M.; et al.  $\beta$ -Caryophyllene Mitigates Collagen Antibody Induced Arthritis (CAIA) in Mice Through a Cross-Talk between CB2 and PPAR- $\gamma$  Receptors. *Biomolecules* **2019**, *9*, 326. [CrossRef] [PubMed]
140. Ma, L.; Niu, W.; Lv, J.; Jia, J.; Zhu, M.; Yang, S. PGC-1 $\alpha$ -Mediated Mitochondrial Biogenesis Is Involved in Cannabinoid Receptor 2 Agonist AM1241-Induced Microglial Phenotype Amelioration. *Cell. Mol. Neurobiol.* **2018**, *38*, 1529–1537. [CrossRef] [PubMed]
141. Ambrosino, P.; Soldovieri, M.V.; Russo, C.; Tagliatalata, M. Activation and Desensitization of TRPV1 Channels in Sensory Neurons by the PPAR $\alpha$  Agonist Palmitoylethanolamide. *Br. J. Pharmacol.* **2013**, *168*, 1430–1444. [CrossRef]
142. Ambrosino, P.; Soldovieri, M.V.; De Maria, M.; Russo, C.; Tagliatalata, M. Functional and Biochemical Interaction between PPAR $\alpha$  Receptors and TRPV1 Channels: Potential Role in PPAR $\alpha$  Agonists-Mediated Analgesia. *Pharmacol. Res.* **2014**, *87*, 113–122. [CrossRef]

143. Iannotti, F.A.; Hill, C.L.; Leo, A.; Alhusaini, A.; Soubrane, C.; Mazzarella, E.; Russo, E.; Whalley, B.J.; Di Marzo, V.; Stephens, G.J. Nonpsychotropic Plant Cannabinoids, Cannabidiol (CBD) and Cannabidiol (CBD), Activate and Desensitize Transient Receptor Potential Vanilloid 1 (TRPV1) Channels in Vitro: Potential for the Treatment of Neuronal Hyperexcitability. *ACS Chem. Neurosci.* **2014**, *5*, 1131–1141. [[CrossRef](#)]
144. Melis, M.; Pistis, M. Targeting the Interaction between Fatty Acid Ethanolamides and Nicotinic Receptors: Therapeutic Perspectives. *Pharmacol. Res.* **2014**, *86*, 42–49. [[CrossRef](#)] [[PubMed](#)]
145. Baskaran, P.; Krishnan, V.; Ren, J.; Thyagarajan, B. Capsaicin Induces Browning of White Adipose Tissue and Counters Obesity by Activating TRPV1 Channel-Dependent Mechanisms. *Br. J. Pharmacol.* **2016**, *173*, 2369–2389. [[CrossRef](#)] [[PubMed](#)]
146. Li, L.; Chen, J.; Ni, Y.; Feng, X.; Zhao, Z.; Wang, P.; Sun, J.; Yu, H.; Yan, Z.; Liu, D.; et al. TRPV1 Activation Prevents Nonalcoholic Fatty Liver through UCP2 Upregulation in Mice. *Pflugers Arch* **2012**, *463*, 727–732. [[CrossRef](#)]
147. Gao, F.; Liang, Y.; Wang, X.; Lu, Z.; Li, L.; Zhu, S.; Liu, D.; Yan, Z.; Zhu, Z. TRPV1 Activation Attenuates High-Salt Diet-Induced Cardiac Hypertrophy and Fibrosis through PPAR- $\delta$  Upregulation. Available online: <https://www.hindawi.com/journals/ppar/2014/491963/> (accessed on 25 February 2021).
148. Depommier, C.; Vitale, R.M.; Iannotti, F.A.; Silvestri, C.; Flamand, N.; Druart, C.; Everard, A.; Pelicaen, R.; Maiter, D.; Thissen, J.-P.; et al. Beneficial Effects of *Akkermansia muciniphila* Are Not Associated with Major Changes in the Circulating Endocannabinoidome but Linked to Higher Mono-Palmitoyl-Glycerol Levels as New PPAR $\alpha$  Agonists. *Cells* **2021**, *10*, 185. [[CrossRef](#)] [[PubMed](#)]
149. Manca, C.; Shen, M.; Boubertakh, B.; Martin, C.; Flamand, N.; Silvestri, C.; Di Marzo, V. Alterations of Brain Endocannabinoidome Signaling in Germ-Free Mice. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **2020**, *1865*, 158786. [[CrossRef](#)] [[PubMed](#)]