

Article

PET Imaging Biomarkers of Anti-EGFR Immunotherapy in Esophageal Squamous Cell Carcinoma Models

Tae Sup Lee^{1,*}, In Ho Song¹, Jong Il Shin¹, Yong Serk Park², Jung Young Kim¹, Kwang Il Kim¹, Yong Jin Lee¹, Joo Hyun Kang¹

¹ Division of RI Application, Korea Institute of Radiological and Medical Sciences (KIRAMS), Seoul, Republic of Korea; nobelcow@kirams.re.kr (T.S.L); inosong@naver.com (I.H.S); shinjongil@kirams.re.kr (J.I.S); ykim@kirams.re.kr (J.Y.K); kikim@kirams.re.kr (K.I.K); yjlee@kirams.re.kr (Y.J.L); kang2325@kirams.re.kr (J.H.K);

² Department of Biomedical Laboratory Science, College of Health Science, Yonsei University, Wonju, Republic of Korea; parkys@yonsei.ac.kr (Y.S.P)

* Correspondence: nobelcow@kirams.re.kr; Tel.: +82-2-970-1351

Supplementary data




Figure S1. Relative bodyweight changes by cetuximab treatment in ESCC tumor model. a and b, comparison of relative body mass of TE-4 (a) and TE-8 (b) xenograft models treated with isotype or cetuximab. No apparent body weight loss was observed in all groups.



© 2018 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).