

Using next generation sequencing to disentangle the diet and incidence of intestinal parasites of Falkland Flightless Steamer Duck *Tachyeres brachypterus* and Patagonian Crested Ducks *Lophonetta specularioides* sharing a South Atlantic island

Juan F. Masello ^{1,*}, Yvonne R. Schumm ¹, Sven Griep ² and Petra Quillfeldt ¹

¹ Department of Animal Ecology & Systematics, Justus Liebig University Giessen, Heinrich-Buff-Ring 26, D-35392 Giessen, Germany;

² Institute for Bioinformatics & Systems, Biology Justus Liebig University Giessen, Heinrich-Buff-Ring 58, D-35392 Giessen, Germany;

* Correspondence: juan.masello@gmx.de



Figure S1. Falkland Flightless Steamer Duck *Tachyeres brachypterus* (front) and Patagonian Crested Ducks *Lophonetta specularioides* (back) sharing a beach environment on New Island. Note the female Falkland Flightless Steamer Duck defecating (front left). Photo credit: Petra Quillfeldt



Figure S2. Female (right) and male (left) Falkland Flightless Steamer Duck *Tachyeres brachydactyla* from New Island. Photo credit: Petra Quillfeldt



Figure S3. Falkland Flightless Steamer Duck *Tachyeres brachydactyla* parents and their chicks foraging in the vicinity of Protector Beach, New Island. Photo credit: Petra Quillfeldt



Figure S4. Falkland Flightless Steamer Duck *Tachyeres brachydactyla* parents and their chicks resting on South Harbour Beach, New Island, where several of the scats were collected. Photo credit: Petra Quillfeldt



Figure S5. Patagonian Crested Duck *Lophonetta specularoides* foraging on New Island. Photo credit: Petra Quillfeldt



Figure S6. Patagonian Crested Duck *Lophonetta specularoides* adult and chicks foraging on Protector Beach, New Island. Photo credit: Petra Quillfeldt



Figure S7. Patagonian Crested Duck *Lophonetta specularioides* adult foraging on Protector Beach, New Island.
Photo credit: Petra Quillfeldt