






Article

Impact of Agricultural Drought on Sunflower Production across Hungary

Endre Harsányi ¹, Bashar Bashir ² , Firas Alsilibe ³, Karam Alsafadi ⁴ , Abdullah Alsalman ², Adrienn Széles ¹, Muhammad Habib ur Rahman ^{5,6} , István Bácskai ¹, Csaba Juhász ¹ , Tamás Ratonyi ¹ and Safwan Mohammed ^{1,*} 

- ¹ Faculty of Agricultural and Food Sciences and Environmental Management, Institute of Land Utilization, Technology and Regional Development, University of Debrecen, 4032 Debrecen, Hungary; harsanyie@agr.unideb.hu (E.H.); szelesa@agr.unideb.hu (A.S.); bacsikai.istvan@agr.unideb.hu (I.B.); juhasz@agr.unideb.hu (C.J.); ratonyi@agr.unideb.hu (T.R.)
- ² Department of Civil Engineering, College of Engineering, King Saud University, Riyadh 11421, Saudi Arabia; bbashir@ksu.edu.sa (B.B.); asalman@ksu.edu.sa (A.A.)
- ³ Department of Transport Infrastructure and Water Resources Engineering, Széchenyi István University, 9026 Győr, Hungary; firas.alsilibe@hallgato.sze.hu
- ⁴ School of Geographical Sciences, Nanjing University of Information Science and Technology, Nanjing 210044, China; karam.alsafadi@alexu.edu.eg
- ⁵ Institute of Crop Science and Resource Conservation (INRES), University of Bonn, 53115 Bonn, Germany; mhabibur@uni-bonn.de
- ⁶ Department of Agronomy, MNS-University of Agriculture, Multan 60000, Pakistan
- * Correspondence: safwan@agr.unideb.hu



Citation: Harsányi, E.; Bashir, B.; Alsilibe, F.; Alsafadi, K.; Alsalman, A.; Széles, A.; Rahman, M.H.u.; Bácskai, I.; Juhász, C.; Ratonyi, T.; et al. Impact of Agricultural Drought on Sunflower Production across Hungary.

Atmosphere **2021**, *12*, 1339. <https://doi.org/10.3390/atmos12101339>

Academic Editor: Ognjen Bonacci

Received: 23 September 2021

Accepted: 8 October 2021

Published: 13 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: In the last few decades, agricultural drought (Ag.D) has seriously affected crop production and food security worldwide. In Hungary, little research has been carried out to assess the impacts of climate change, particularly regarding droughts and crop production, and especially on regional scales. Thus, the main aim of this study was to evaluate the impact of agricultural drought on sunflower production across Hungary. Drought data for the Standardized Precipitation Index (SPI) and the Standardized Precipitation Evapotranspiration Index (SPEI) were collected from the CAR-BATCLIM database (1961–2010), whereas sunflower production was collected from the Hungarian national statistical center (KSH) on regional and national scales. To address the impact of Ag.D on sunflower production, the sequence of standardized yield residuals (SSYR) and yield losses Y_{lossAD} was applied. Additionally, sunflower resilience to Ag.D (SRAG.D) was assessed on a regional scale. The results showed that Ag.D is more severe in the western regions of Hungary, with a significantly positive trend. Interestingly, drought events were more frequent between 1990 and 2010. Moreover, the lowest SSYR values were reported as -3.20 in the Hajdu-Bihar region (2010). In this sense, during the sunflower growing cycle, the relationship between SSYR and Ag.D revealed that the highest correlations were recorded in the central and western regions of Hungary. However, 75% of the regions showed that the plantation of sunflower is not resilient to drought where $SRAG.D_x < 1$. To cope with climate change in Hungary, an urgent mitigation plan should be implemented.

Keywords: crop resilience; food security; poverty; SPI; SPEI; SSYR; Hungary

1. Introduction

Climate change is one of the most challenging and complex issues facing the world today [1]. Over the last few centuries, the burning of fossil fuels and widespread deforestation have resulted in increased atmospheric greenhouse gas (GHG) concentrations, which have led to significant climate shifts across the planet [2]. Since 1750, the concentrations of GHGs such as methane (CH₄), carbon dioxide (CO₂), and nitrous oxide (N₂O) have risen by 156%, 47%, and 23%, respectively [1,3]. Due to anthropogenic activities, the average temperature rose by 0.99 °C, in the first two decades of the 21st century (2001–2020), and according to

predictions, by 2050, the temperature increase is expected to reach 1.5 °C, or perhaps even higher than the average temperature in the preindustrial period (1850–1900) [1]. This accelerated rise in temperature has caused a spike in droughts, floods, irregular precipitation patterns, heat waves, and other extreme climate events around the world [4,5].

According to the annual report of Weather, Climate, and Catastrophe Insight [6], natural disasters alone caused direct losses and damage totaling over USD 268 billion worldwide in 2020; on average, losses due to natural disasters and emergencies have exceeded USD 200 billion per year since 2016. Approximately 95% of these losses are linked to weather-related events, where hurricanes, floods and droughts are the major contributors and have a direct link to climate change [6]. Overall, the impact of climate change is extensive, but its effects are now clearly visible on the agricultural sector [7], on which the world's food production and economy relies. It is also important to highlight that the world population is predicted to hit 9.7 billion by 2050 [8], which would increase the pressure on the agricultural sector to fulfill the growing food demands already impacted by climate change.

Among natural disasters caused by weather and climate change, drought is a unique issue; the global trend towards more or less frequent drought episodes is still a highly discussed topic [9–14]. This is attributed to the complicated characteristics of droughts, which usually develop slowly but can extend for months or even years [15,16]. Furthermore, the consequences can have a variety of effects on several sectors [17–20]. Additionally, the lack of a unified universal definition [21,22] and the wide range of indicators used to assess meteorological, agricultural, hydrological, and socioeconomic droughts [23] enhance the complexity.

Drought impacts all socioeconomic and environmental systems [24], with significant effects on anthropogenic fields including agriculture, forestry, water resource management, energy generation, and health [25]. Drought impacts can be divided into two categories: direct and indirect [25]. Direct effects include decreased crop production [26], increased numbers of forest fires [27], decreased water levels [28], and increased livestock mortality rates [29]. Indirect effects of drought include highly unstable food prices, which may be exacerbated by market effects in the agricultural sector [30]. As a result, estimating total costs and losses at the regional and national levels is difficult. Indirect losses are mostly greater than direct losses [31], although they are more difficult to link to a specific event.

In order to detect, monitor, and characterize drought events, many drought indices have been developed, such as the Palmer Drought Severity Index Standardized (PDSI) [32], Precipitation Index (SPI) [33], and the Standardized Precipitation Evapotranspiration Index (SPEI) [34]. Table 1 presents more details about the most commonly used drought indices in the literature.

Table 1. Common drought indices.

Index	Definition	Factor	Calculation Time Scale
PDSI [32]	Palmer Drought Severity Index	Precipitation, temperature, soil moisture and evapotranspiration	Monthly
CMI [35]	Crop Moisture Index	Mean temperature and precipitation	Weekly
CSDI [36]	Crop Specific Drought Index	Evapotranspiration	Seasonal
RI [37]	National Rainfall Index	Precipitation	Annually and every century
RDI [38]	Reclamation Drought Index	Level of river water, snowfall, stream flows, reservoirs level and temperature	Monthly
EPI [39]	Effective Precipitation Index	Precipitation	Daily

Table 1. *Cont.*

Index	Definition	Factor	Calculation Time Scale
BMDI [40]	Bhalme and Mooley Drought Index	Precipitation	Monthly, annually
SPI [33]	Standardized Precipitation Index	Precipitation	3-, 6-, 12-, 24- and 48-month periods.
SPEI [34]	Standardized Precipitation Evapotranspiration Index	Precipitation, evapotranspiration	Monthly
SRI [41]	Standardized Runoff Index	Precipitation	3-, 6-, 12-, 24- and 48-month periods.

In Europe, the accelerated warming in this region [42], in comparison with many other parts of the world, has intensified drought events for prolonged periods, accompanied by rising temperatures and low rainfall, particularly in the center of the continent (i.e., Hungary) [43,44].

Hungary, similarly to other European countries in the Carpathians, is affected by droughts and climate change [45,46]. Droughts have been common in Hungarian history, leading to reduced crops, animal devastation, and the risk of hunger and illnesses [47]. The frequency of drought events has also increased significantly in Hungary [47]. Generally, every two years, Hungary experiences a moderate drought, and every three years, it experiences a severe drought [48]; these drought conditions are projected to continue for the rest of the 21st century [45,49]. In this sense, little research has been conducted in Hungary to assess the impact of climate change on crop production, especially on a regional scale. Hence, our understanding of the direct impacts of different drought cycles on crop production is still limited. Thus, the main aims of this research were: (1) to track agricultural drought (Ag.D) evolution across Hungarian counties between 1960 and 2010 by using the Standardized Precipitation Index (SPI) and the Standardized Precipitation Evapotranspiration Index (SPEI); (2) to analyze the dynamic interaction between sunflower yield and drought cycles; and (3) to identify the counties most prone to drought. In this study, sunflower crops were chosen due to their economic importance, where it is considered one of the most important oil crops (550,000 ha) across Hungary.

2. Materials and Methods

2.1. Data Collection and Trend Analysis

Agricultural drought (Ag.D) data were collected from the CARBATCLIM database [50]. The CARBATCLIM platform provides researchers with necessary climate data between 1960 and 2010 for the whole Carpathian Region (44° N and 50° N, 17° E and 27° E). This project was supported by the European Commission, and the output is a climate atlas with spatial resolution $0.1^\circ \times 0.1^\circ$ [51,52]. Within the CARBATCLIM database, Hungary is covered by 1045 grid points (Figure 1), where each point contains all climate variables. In this study, we downloaded the data for both the Standardized Precipitation Index (SPI) and the Standardized Precipitation Evapotranspiration Index (SPEI) for six months (i.e., SPI-6, SPEI-6), as a proxy of Ag.D. Notably, homogeneity of the data and their quality were assessed by the CARBATCLIM team. Later on, gridded points were sorted into groups according to their Hungarian counties; then, the average of each group was calculated and adopted as a representative value of Ag.D (Figure 1).




Figure 1. Map of Hungary: (a) Hungarian counties, and (b) distribution of 1045 gridded points across Hungary.

The sunflower database was built using information from the Hungarian national statistical center (KSH) (https://www.ksh.hu/stadat_eng accessed on 6 May 2020). This database contains the planted area (ha), production (thousand tons) and yield (kg/ha), at a national scale (1990–2019), as represented in Figure 2. However, to address the specific study goals, available sunflower yield data were collected on a county scale from 2000 to 2019.

For trend analysis, Mann–Kendall (MK) [53] tests were performed and Sen slopes (β) [54] were calculated for the studied variables. Both the MK test and Sen slope (β) are non-parametric tests; MK is used to indicate trends (i.e., increase or decrease) in the studied time series, whereas the Sen slope computes the value of the change (slope).




Figure 2. Evolution of sunflower cultivation across Hungary at a national scale (1990–2019): (a) harvested sunflower land between 1990 and 2019 (left graph); (b) production in tons across Hungary (middle graph); and (c) sunflower yield between 1990 and 2019 (right graph).

2.2. Agricultural Drought Indices

To quantify drought characteristics (severity, spatial extent, duration, and frequency), several drought indices could be employed, which mainly depend on the research questions and data availability [55–57]. Many ecosystem elements such as rainfall, evapotranspiration, temperature, river discharge, soil moisture, and change in vegetation cover could be used as inputs for modeling and monitoring drought [58–60]. Scientifically, the Ag.D indicates a shortage in soil moisture, which inhibits soil from providing necessary amounts of water to attain optimal crop production requirements [56,61]. In this study, the SPI and SPEI in a six-month time scale were adopted as a proxy for Ag.D (1961–2010).

2.2.1. Standardized Precipitation Index (SPI)

The SPI [33] is one of the most common indices used for tracking and monitoring drought episodes [62], and is recommended by the World Meteorological Organization [55,63]. Monthly rainfall data are the only input for calculating drought in several time scales [33]. Through different transformations, from gamma distribution to normal distribution, the final equations for SPI can be drawn as follows:

$$\text{SPI} = -\left(t - \frac{2.515517 + 0.802853t + 0.010328t^2}{1 + 1.432788t + 0.189269t^2 + 0.001308t^3}\right) \text{ for } 0 < H(x) \leq 0.5 \quad (1)$$

$$\text{SPI} = +\left(t - \frac{2.515517 + 0.802853t + 0.010328t^2}{1 + 1.432788t + 0.189269t^2 + 0.001308t^3}\right) \text{ for } 0 < H(x) < 1 \quad (2)$$

where:

$$H(x) = q + (1 - q)G(x) \quad (3)$$

More details about SPI calculation, (t , q , $G(x)$), was presented in detail by McKee et al. [33]. Drought classifications according to SPI values are presented in Table 2.

Table 2. Drought classes based on SPI and SPEI values.

SPI Value	SPEI Value	Ag.D Class
>0	>0	NA
−0.84–0	−0.84–0	Moderate Ag.D
−1.28–−0.84	−1.28–−0.84	Severe Ag.D
−1.65–−1.28	−1.65–−1.28	Extreme Ag.D
>−1.65	>−1.65	Very extreme Ag.D

2.2.2. SPEI

The SPEI [34] is a developed indicator from the same background as SPI [52]. Monthly rainfall and potential evapotranspiration are the main inputs for calculating the SPEI [34]. The final equation of SPEI could be drawn as follows:

$$\text{SPEI} = W - \frac{C_0 + C_1 W + C_2 W}{1 + d_1 W + d_2 W^2 + d_3 W^3} \quad (4)$$

$$\text{where } W = \begin{cases} \sqrt{-2 \ln(P)} & \text{when } P \leq 0.5 \\ \sqrt{-2 \ln(1-P)} & \text{when } P > 0.5 \end{cases} \quad (5)$$

The equation constant ($C_0, C_1 \dots$); and other mathematical approaches was presented in detail by Vicente-Serrano et al. [34]. Drought classifications according to SPEI values are presented in Table 2.

Despite the fact that both indices have the same background, the SPEI has been proven to be superior to SPI in drought monitoring and climate change assessments [33,52].

2.3. Impact of Agricultural Drought

2.3.1. Sequence of Standardized Yield Residuals (SSYR)

To address the impact of Ag.D on crop yield, the bias attributed to technological factors (i.e., pest control, fertilization, and high-yield varieties) should be removed. To do so, polynomial regression was calculated for observed yield ($Y_{R(x)}$) on the county scale (2000–2019); then, the data were detrended ($Y_{P(x)}$), and the residuals were calculated (i.e., $Y_{R(x)} - Y_{P(x)}$). Finally, the SSYR was computed using the following equation [64,65]:

$$\text{SSYR} = \frac{Y_{P(x)} - \forall_{P(x)}}{\partial_{P(x)}} \quad (6)$$

where $Y_{P(x)}$ is the potential yield (detrended), $\forall_{P(x)}$ is the mean of $Y_{P(x)}$, and $\partial_{P(x)}$ is the standard deviation $Y_{P(x)}$. The classifications of SSYR values are presented in Table 3. Additionally, the correlation coefficients (r) between SSYR and both the SPI and SPEI were calculated (2000–2010) on a monthly scale.

Table 3. Classification of the sequence of standardized yield residual (SSYR) values which ranged from normal conditions to extreme drought impacts.

SSYR Value	Impacts of Ag.D
$-0.5 < \text{SSYR} \leq 0.5$	Normal
$-1.0 < \text{SSYR} \leq -0.5$	Mild
$-1.5 < \text{SSYR} \leq -1.0$	Moderate
$-2.0 < \text{SSYR} \leq -1.5$	High
$\text{SSYR} \leq -2.0$	Extreme

2.3.2. Yield Losses, $Y_{\text{loss Ag.D}}$

The direct impacts of Ag.D on sunflower production and yield losses across the Hungarian counties were calculated using the approach suggested by Tigkas et al. [66]:

$$Y_{\text{loss Ag.D}(x)} = \frac{Y_{R(x)} - Y_{P(x)}}{Y_{P(x)}} \times 100 \quad (7)$$

where $Y_{\text{loss Ag.D}(x)}$ is the yield loss due to agricultural drought, $Y_{R(x)}$ is the recorded yield (observed), $Y_{P(x)}$ is the potential yield (detrended), and x is the year. $Y_{P(x)}$ refers to the development of crop production without any environmental constraints such as water shortages, heat waves, or other environmental factors [66].

2.3.3. Sunflower Resilience to Ag.D (SRAg.D):

The SRAg.D was calculated for each county in Hungary using the equation suggested by Sharma and Goyal [67]:

$$\text{SRAg.D}_x = \frac{Y_{R(x)}^d}{Y_{P(x)}^d} \quad (8)$$

where SRAg.D_x is the sunflower resilience to Ag.D in each county, $Y_{R(x)}^d$ is the recorded yield in the driest year (2000–2010), and $Y_{P(x)}^d$ is the potential yield. The SRAg.D ranged from >1 (resilient) to <0.8 (severely non-resilient). The classification of SRAg.D values are presented in Table 4.

Table 4. Classification of sunflower resilience to Ag.D (SRAg.D) values, which ranged from resilient to severely non-resilient.

SRAg.D Value	SRAg.D
$\text{SRAg.D} > 1$	Resilient
$0.9 < \text{SRAg.D} < 1$	Slightly non-SRAg.D
$0.8 < \text{SRAg.D} < 0.9$	Moderately non-SRAg.D
$\text{SRAg.D} < 0.8$	Severely non-SRAg.D

For simplification, Figure 3 depicts a flowchart of the study steps.




Figure 3. Flowchart of the study process from obtaining the data through to providing conclusions and recommendations. SPI, SPEI: drought indices. Trend: MK test. $Y_{R(x)}$: observed yield (2000–2019), $Y_{P(x)}$: potential yield (detrended). SSYR: sequence of standardized yield residuals, $Y_{loss Ag.D}$: yield losses, SRAg.D_x : sunflower resilience to Ag.D.

3. Results

3.1. Trend in Ag.D across Hungarian Counties:

Hungary experienced Ag.D cycles several times, as can be seen in Figure 4. Figure 4a,d demonstrate that Ag.D episodes were more intense in western and central Hungary (i.e., ZA, VE, VA, and GY) compared with the eastern region of the country (i.e., HB, HE, JN, and SS). Notably, Ag.D cycles have become more frequent since 1990, when more negative values were recorded (Figure 4a,d).

Trend analyses by MK test and Sen slope also indicate that western Hungarian counties are more prone to Ag.D compared with other country regions (Table 5). In terms of SPI-6 (1960–2010), the majority of the counties (14 out of 20) exhibited a positive trend, which indicate less susceptibility to Ag.D. In fact, only four counties, HB, HE, JN, and SS, which are located in eastern Hungary, exhibited a significantly positive trend ($p < 0.05$) (Table 5).

For SPEI, most of counties exhibited a negative trend for SPEI-6 (1960–2010), except for HB, HE, JN, and SS. Nonetheless, significantly negative ($p < 0.05$) trends in SPEI-6 (1960–2010) were recorded in GY, SO, VA, VE, and ZA (western regions) (Table 5). Notably, both drought indices agreed on negative trends in each of the following counties: GY, SO, TO, VA, VE, and ZA, as shown in Figure 4b.

Table 5. Trends in Ag.D indices (SPI-6, SPEI-6) and sunflower production (kg/ha) across Hungary.

County	Code	SPI-6		SPEI-6		Sunflower	
		MK and β	p	MK and β	p	MK and β	p
Bács-Kiskun	BC	0.0005	0.05	-5×10^{-5}	0.83	+55.83	0
Baranya	BA	0.0002	0.32	-2×10^{-4}	0.40	+48.04	0
Békés	BE	0.0005	0.05	6×10^{-5}	0.79	+59.03	<0.0001
Borsod-Abaúj-Zemplén	BO	0.0005	0.06	7×10^{-5}	0.75	+75.19	<0.0001
Budapest	BU	0.0003	0.28	-3×10^{-4}	0.27	+70.28	0
Csongrád-Csanád	CS	0.0003	0.21	-1×10^{-4}	0.55	+32.18	0.01
Fejér	FE	0.0001	0.67	-4×10^{-4}	0.06	+47.99	0.01
Győr-Moson-Sopron	GY	0.0001	0.74	-5×10^{-4}	0.03	+38.33	0.01
Hajdú-Bihar	HB	0.0006	0.01	2×10^{-4}	0.46	+71.6	<0.0001
Heves	HE	0.0005	0.02	1×10^{-4}	0.60	+71.34	0
Jász-Nagykun-Szolnok	JN	0.0007	0.00	2×10^{-4}	0.44	+60	<0.0001
Komárom-Esztergom	KE	0.0003	0.21	-2×10^{-4}	0.26	+58.06	<0.0001
Nógrád	NO	0.0003	0.24	-1×10^{-4}	0.65	+65	0.01
Pest	PE	0.0003	0.16	-2×10^{-4}	0.48	+65.15	0
Somogy	SO	−0.0004	0.07	-8×10^{-4}	0.00	+61.46	0
Szabolcs-Szatmár-Bereg	SS	0.0008	0.00	3×10^{-4}	0.15	+72.58	0
Tolna	TO	0.0000	0.94	-4×10^{-4}	0.06	+53.54	0
Vas	VA	−0.0008	0.00	-1×10^{-3}	<0.0001	+44.77	0
Veszprém	VE	−0.0001	0.69	-6×10^{-4}	0.01	+56.57	0
Zala	ZA	−0.0007	0.01	-1×10^{-3}	<0.0001	+50	<0.0001

* Gray and bold values indicate a significance level of 0.05 ($p < 0.05$). MK, trend; β , Sen slope.

Despite the differences in identifying drought events between SPI-6 and SPEI-6 due to different inputs and mathematical equations, both indices highlighted the following years: 1962, 1968, 1972, 1983, 1986, 1990, 1992–1993, 2000–2003, and 2007 as drastic periods (SPI-6 and SPEI-6 values of less than -1.28) in terms of Ag.D (Figure 4a,d).

The lowest recorded values of SPI-6 and SPEI-6 (i.e., the highest recorded levels of drought) in each county are presented in Figure 4c. For SPI-6, the lowest values were recorded in CS (-3.33), GY (-2.96), VA (-2.95), and TO (-2.8). Unexpectedly, the lowest values of SPEI-6 were recorded in central and southern counties. For instance, the lowest value of SPEI-6 was recorded in Csongrád-Csanád (CS) (-4.71), which is located in southern Hungary, followed by BU (-4.21) (central), then GY (-4.04) (western) (Figure 4c).

As depicted in Figure 4c, the SPEI-6 (red line) represents higher Ag.D values than the SPI-6 (blue line); this can mainly be explained by the fact that the SPEI-6 employs both evapotranspiration and rainfall for drought computation, whereas the SPI-6 depends only on the monthly changes in rainfall. Notably, the conjunction of evapotranspiration and rainfall inflated the drought values in the study area.




Figure 4. Ag.D evolution across Hungary (1960–2010): (a) Evolution of SPI-6 across Hungarian counties (1960–2010); (b) trends in SPI-6 and SPEI-6 based on MK tests and Sen slopes; (c) lowest recorded values of SPI-6 and SPEI-6 across Hungarian counties (1960–2010); (d) Evolution of SPEI-6 across Hungarian counties (1960–2010).

3.2. Sequence of Standardized Yield Residuals (SSYR)

According to the MK tests and Sen slopes, sunflower yield (kg/ha) exhibited a positive trend across Hungarian counties between 2000 and 2019 (Table 5, Figure 5). The greatest increase in sunflower yield was observed in BO (+75.19 kg/ha, $p < 0.05$), followed by SS (+72.58 kg/ha, $p < 0.05$), then HB (+71.6 kg/ha, $p < 0.05$). Notably, these counties are located in eastern Hungary, and experienced positive but non-significant trends for both SPI-6 and SPEI-6 (Table 5, Figure 5).

Thus, to isolate the positive impact of human intervention which include, but are not limited to, the selection of new varieties, pest control, advance field irrigation technology, and precision agriculture, the sequence of standardized yield residuals (SSYRs) was applied.




Figure 5. Sunflower yield in the studied Hungarian counties (2000–2019). (a) BA, (b) BC, (c) BE, (d) BO, (e) BU, (f) CS, (g) FE, (h) GY, (i) HB, (j) HE, (k) JN, (l) KE, (m) NO, (n) PE, (o) SO, (p) SS, (q) TO, (r) VA, (s) VE, (t) ZA.

The output of the SSYR analysis highlighted the yeEars 2000, 2010, and 2012 as the years most affected by Ag.D ($SSYR \leq -1$) in terms of sunflower production, as shown in Figure 6. The lowest SSYR was recorded in HB (−3.2, extreme impact in 2010), followed by TO (−2.41, extreme impact in 2012), and VA (−2.34, extreme impact in 2010) (Figure 6).




Figure 6. Evolution of SSYR in the studied Hungarian counties (2000–2019). (a) BA, (b) BC, (c) BE, (d) BO, (e) BU, (f) CS, (g) FE, (h) GY, (i) HB, (j) HE, (k) JN, (l) KE, (m) NO, (n) PE, (o) SO, (p) SS, (q) TO, (r) VA, (s) VE, (t) ZA.

During the growing cycle (April–October), a high correlation between SSYR and Ag.D was observed, especially in summer seasons (Figures 7 and 8). For SPI-6, the highest correlation (Max. r SSYR vs. SPI-6) was recorded in VA (r SSYR vs. SPI-6 = 0.6, July, western Hungary) and BC (r SSYR vs. SPI-6 = 0.55, August, central Hungary) (Figure 7). Similarly, central and western counties exhibited the highest r values between SSYR and SPEI-6. For instance, the Max. r SSYR vs. SPEI-6 was obtained in PE (r SSYR vs. SPEI-6 = 0.63, July, central Hungary), BC (r SSYR vs. SPEI-6 = 0.59, July, central Hungary), and VE (r SSYR vs. SPEI-6 = 0.51, July, western Hungary) (Figure 8). However, both Ag.D indices indicate that drought cycles affected the sunflower yield during the summer months, where the western counties were classified as the most affected by drought.




Figure 7. Correlation values between SSYR and SPI during the sunflower growing cycle (2000–2010) (right table), and max correlation (Max. r) between the SSYR and SPI (left figure).




Figure 8. Correlation values between the SSYR and SPEI during the sunflower growing cycle (2000–2010) (right table), and max correlation (Max. r) between the SSYR and SPEI (left figure).

3.3. Yield Losses $Y_{loss Ag.D}$

Between 2000 and 2010, the loss of sunflower yield was distinguished in two different years, as depicted in Figure 9. In 2004, the highest $Y_{loss Ag.D}$ was recorded in VE ($Y_{loss Ag.D} = -38.32\%$, western Hungary), followed by HE ($Y_{loss Ag.D} = -33.95\%$, eastern Hungary) and then BE ($Y_{loss Ag.D} = -32.89\%$, central Hungary) (Figure 9). However, in 2008, all counties across Hungary experienced yield losses, where the highest losses were

observed in NO ($Y_{loss\ Ag.D} = -42.03\%$, northern Hungary), JN ($Y_{loss\ Ag.D} = -33.49\%$, central Hungary), and SS ($Y_{loss\ Ag.D} = -29.37\%$, eastern Hungary).




Figure 9. Sunflower resilience to Ag.D (SRAg.D) (right), and max yield losses (left).

By calculating the SRAg.D, the resistance of sunflower crops to drought events could be distinguished across Hungarian counties (Figure 10). The results showed that the crop yields in central and western regions were more prone to loss due to drought events. The SRAg.D value reached 0.74 (severely non-SRAg.D) in HE, and 0.79 (also severely non-SRAg.D) in PE. In contrast, the highest value was recorded in eastern Hungary (BE; 1.11).




Figure 10. Resilience of sunflower production to Ag.D across Hungarian counties (2000–2010).

4. Discussion

In the last few decades, drought cycles have become more intense and frequent all over the world due to rapid climate change [9,68–70]; for instance, in Brazil [71], Syria [63], Hungary [52], China [72], and southern Europe [73]. Subsequently, many industries have been affected either directly or indirectly, such as the agricultural sector [74–79], hydrology [80–83], the economy [76,84], human health [85–87], and tourism [88–90]. In this sense, our research shows a negative trend in agricultural drought episodes across Hungarian counties (Figure 4; Table 5). Additionally, the results emphasize the direct

impact of Ag.D on sunflower production (Figure 6), where the yield losses reached 40% in some counties (Figure 9).

Many studies have been carried out across Europe for monitoring and assessing drought evolution, identifying a positive trend, especially in southern and central regions [73,91–93]. Hungary, which is in the center of the Pannonian basin (central Europe), suffers from drought episodes. For instance, Alsafadi et al. [52] reported an increase in drought trends in western Hungary comparing with the east; however, drought frequency was more intense in central Hungary. Mohammed et al. [49] indicated a positive strong correlation between Ag.D (SPI-3, SPI-6) and NDVI (Normalized Difference Vegetation Index). However, future climate projections indicate an increase in drought cycles due to changing future rainfall patterns (2071–2100) [45]. At the regional scale (county scale), drought evolution has been more intense in central and western Hungary than the east. Similar results were reported by Szabó et al. [94], where western Hungary was identified as being more susceptible to climate change. This issue could be explained by the fact that the central region receives less rainfall than other areas [52]; thus, the SPI and SPEI values are decreased, which directly lead to the evolution of drought. In this case, soil moisture will be decreased, and less water will be available for use on agricultural crops (i.e., sunflowers), which will lead to water stress and yield losses (Figures 6, 9 and 10) due to the inhibition of physiological functioning in some crops [69]. Interestingly, this phenomenon could affect all crops, especially maize, which is very sensitive [79]. In this sense, Adrienn and Janos [95] reported that agricultural drought is the main constraint for crop production across Hungary.

Both SPI and SPEI have drawbacks, which generates some uncertainty in the results. For SPI, using only monthly rainfall data without considering their temporal distribution could affect the interpretation of results, along with failure in predicting the exact times of the drought evolution cycle (i.e., start and end) [34,96]. For SPEI, heat waves can be misinterpreted as droughts in some areas [97]. Additionally, calculating evapotranspiration based only on temperature is not sufficient [98]. However, both indices are widely used all over the world, and their outputs can be used for monitoring drought [72,99–101].

The interaction between crop yield and drought indices during the growing cycle (April–October) could be linked to increasing crop evapotranspiration (crop-ET₀) and temperature. In this sense, Stagge et al. [73] reported an increase in crop-ET₀ and temperature. However, the resistance of sunflower to drought was tracked across Hungarian counties. The results also showed that crop yield in central and eastern Hungary was less resilient to drought, where the SRAg.D value reached 0.74 (severely non-SRAg.D) in Heves and 0.79 (severely non-SRAg.D) in Pest. However, the highest value was recorded in eastern Hungary (Bekes, 1.11). In fact, climate change has affected crop production for both irrigated and rainfed agricultural systems, where drought has significantly reduced crop yield [79,102,103]. Globally, temperature and rainfall directly influence crop production, where they behave nonlinearly [103–106]. Thus, any changes in these climate variables or even the interaction between them will affect agricultural production, not only for sunflowers, but for all crop types. Shortage of rainfall is linked to abnormal atmospheric circulation (at high pressure) that prevents clouds and precipitation from forming [68,107], or a change in the rain belt [68]. However, less cloud and precipitation lead to drier conditions, increased temperature, decreased humidity, and increased evapotranspiration demand, which amplify drought conditions [68]. Scientifically, drought causes a shortage in soil water content and leads to a water deficit which directly affects crop yields [65,79,108]. Unfortunately, most agricultural land across Hungary is cultivated as a rainfed agricultural system; thus, climate change (precipitation patterns), and especially drought, will negatively affect the agricultural sector and crop production [49].

There were some limitations to this study, such as drought trends being assessed based on the average of gridded point data that represent each county, instead of using the whole gridded points. Additionally, only 10 years were available as a record of sunflower yield across Hungarian counties, which used to calculate the direct impact of Ag.D on

sunflower yields; more years would have been better for interpreting the relationship between drought and crop production. On the other hand, both SPI and SPEI have their own limitations in identifying drought cycles. In this context, for calculating drought by using SPI, we only need monthly rainfall data; other climate variables are neglected, which could affect the credibility of the SPI [52]. Even though SPEI involves more climate variables (rainfall and potential evapotranspiration (PET)), and is based on the Thorn Thwaite equation, other climate variables are not considered [52]. However, the output from our analysis is enough to draw the attention of decision makers to the evolution of drought and climate change in central Europe.

5. Conclusions

Drought is a multifaceted and complicated natural hazard, which slowly evolves in the ecosystem and is affected by a multitude of physical and biological factors. Hungary, which is located in central Europe (Pannonian basin), frequently suffers from drought events and climate change. However, little research has been conducted in Hungary to assess the impacts of climate change on crop production, especially on a regional scale. In this context, this study was designed to track the evolution of agricultural drought (SPI-6, SPEI-6) across Hungary (1960–2010), and to investigate the dynamic interactions between sunflower yield and drought cycles. The results showed that agricultural drought episodes were more intense in western and central Hungary (i.e., ZA, VE, VA, and GY) compared with the eastern region of the country. Nonetheless, the Ag.D cycles had become more frequent since 1990, where more negative values of Ag.D were recorded.

Sunflower production was badly affected by agricultural drought in 2000, 2010, and 2012, where $SSYR \leq -1$. The lowest SSYR values were recorded in HB (−3.2, extreme impact in 2010), followed by TO (−2.41, extreme impact in 2012), and VA (−2.34, extreme impact in 2010). Notably, yield losses (Yloss Ag.D) were experienced in all Hungarian counties in 2004 and 2008 due to agricultural drought. The results also reveal that crop yield in central and western Hungary were more prone to yield loss due to drought events.

The outputs of this study will be of great interest to stakeholders and decision makers for formulating climate mitigation and adaptation plans for the agricultural sector in Hungary. However, more research to develop new varieties of sunflowers, which are tolerant to drought and other extreme climate events, is highly recommended. In our future research, we will involve other drought indices such as the Crop Moisture Index and Crop Specific Drought Index to highlight the impacts of drought on sunflowers and other strategic crops in Hungary.

Author Contributions: Conceptualization, S.M.; Data curation, A.S.; Formal analysis, K.A.; Funding acquisition, B.B. and A.A.; Investigation, S.M.; Methodology, K.A. and S.M.; Supervision, E.H.; Visualization, S.M.; Writing—original draft preparation, F.A.; Writing—review and editing, E.H., B.B., A.A., A.S., M.H.u.R., I.B., C.J., T.R., and S.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research was supported by RESEARCHERS SUPPORTING PROJECT, grant number RSP-2021/296, King Saud University, Riyadh, Saudi Arabia.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: <http://www.carpatclim-eu.org/pages/download/> (accessed on 6 October 2021).

Acknowledgments: The authors are grateful to CARBATCLIM team for providing data used in this study. Authors would like to thank Debrecen University (Hungary) and King Saud University for their unlimited support.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Zhu, Z.; Lu, L.; Zhang, W.; Liu, W. *AR6 Climate Change 2021: The Physical Science Basis*; IPCC: Geneva, Switzerland, 2021.
2. Shukla, P.R.; Skea, J.; Calvo Buendia, E.; Masson-Delmotte, V.; Pörtner, H.O.; Roberts, D.C.; Zhai, P.; Slade, R.; Ferrat, M.; Neogi, S.; et al. *Climate Change and Land. IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*; IPCC: Geneva, Switzerland, 2019.
3. Mohammed, S.; Gill, A.R.; Alsafadi, K.; Hijazi, O.; Yadav, K.K.; Khan, A.H.; Islam, S.; Cabral-Pinto, M.M.S.; Harsanyi, E. An overview of greenhouse gases emissions in Hungary. *J. Clean. Prod.* **2021**, 127865. [\[CrossRef\]](#)
4. Pachauri, R.K.; Allen, M.R.; Barros, V.R.; Broome, J.; Cramer, W.; Christ, R.; Church, J.A.; Clarke, L.; Dahe, Q.; Dasgupta, P.; et al. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; IPCC: Geneva, Switzerland, 2014.
5. Boris, O.; Seneviratne, S.I. Global changes in extreme events: Regional and seasonal dimension. *Clim. Chang.* **2012**, *110*, 669–696.
6. Podlaha, A.; Löhrinc, M.; Srivastava, G.; Bowen, S.; Kerschner, B. *2020 Annual Report. Weather, Climate & Catastrophe Insight*; AON: London, UK, 2020.
7. Gurdeep Singh, M.; Kaur, M.; Kaushik, P. Impact of climate change on agriculture and its mitigation strategies: A review. *Sustainability* **2021**, *13*, 1318.
8. United Nations. *World Population Prospects 2019: Highlights*; United Nations: New York, USA, 2019.
9. Dai, A. Drought under global warming: A review. *Wiley Interdiscip. Rev. Clim. Chang.* **2011**, *1*, 45–65. [\[CrossRef\]](#)
10. Dai, A. Increasing drought under global warming in observations and models. *Nat. Clim. Chang.* **2012**, *3*, 52–58. [\[CrossRef\]](#)
11. Sheffield, J.; Wood, E.; Roderick, M. Little change in global drought over the past 60 years. *Nature* **2012**, *491*, 435–438. [\[CrossRef\]](#) [\[PubMed\]](#)
12. Jonathan, S.; Naumann, G.; Carrao, H.; Barbosa, P.; Vogt, J. World drought frequency, duration, and severity for 1951–2010. *Int. J. Climatol.* **2014**, *34*, 2792–2804.
13. Trenberth, K.E.; Dai, A.; Van Der Schrier, G.; Jones, P.; Barichivich, J.; Briffa, K.; Sheffield, J. Global warming and changes in drought. *Nat. Clim. Chang.* **2013**, *4*, 17–22. [\[CrossRef\]](#)
14. Siegfried, D.S.; Stewart, R.E.; Wang, H.; Barlow, M.; Berbery, E.H.; Cai, W.; Hoerling, M.P.; Kanikicharla, K.K.; Koster, R.D.; Lyon, B.; et al. Global meteorological drought: A synthesis of current understanding with a focus on SST drivers of precipitation deficits. *J. Clim.* **2016**, *29*, 3989–4019.
15. Jürgen, V.; Somma, F. (Eds.) *Drought and Drought Mitigation in Europe*; Springer Science & Business Media: Berlin, Germany, 2013; p. 14.
16. Raza, A.; Hussain, I.; Ali, Z.; Faisal, M.; Elashkar, E.E.; Shoukry, A.M.; Al-Deek, F.F.; Gani, S. A seasonally blended and regionally integrated drought index using Bayesian network theory. *Meteorol. Appl.* **2021**, *28*, e1992. [\[CrossRef\]](#)
17. Roger, C.S. Agricultural drought indices in current use in Australia: Strengths, weaknesses, and limitations. In *Proceedings of the Agricultural Drought Indices Proceedings of an Expert Meeting, Murcia, Spain, 2–4 June 2011*; p. 85.
18. Ding, Y.; Hayes, M.J.; Widhalm, M. Measuring economic impacts of drought: A review and discussion. *Disaster Prev. Manag. Int. J.* **2011**, *20*, 434–446. [\[CrossRef\]](#)
19. Blauhut, V.; Gudmundsson, L.; Stahl, K. Towards pan-European drought risk maps: Quantifying the link between drought indices and reported drought impacts. *Environ. Res. Lett.* **2015**, *10*. [\[CrossRef\]](#)
20. Blauhut, V.; Stahl, K.; Stagge, J.H.; Tallaksen, L.M.; Stefano, L.D.; Vogt, J. Estimating drought risk across Europe from reported drought impacts, drought indices, and vulnerability factors. *Hydrol. Earth Syst. Sci.* **2016**, *20*, 2779–2800. [\[CrossRef\]](#)
21. Mishra, A.K.; Singh, V.P. A review of drought concepts. *J. Hydrol.* **2010**, *391*, 202–216. [\[CrossRef\]](#)
22. Lloyd-Hughes, B. The impracticality of a universal drought definition. *Theor. Appl. Clim.* **2013**, *117*, 607–611. [\[CrossRef\]](#)
23. Heim Richard, R., Jr. A review of twentieth-century drought indices used in the United States. *Bull. Am. Meteorol. Soc.* **2002**, *83*, 1149–1166. [\[CrossRef\]](#)
24. Vicente-Serrano, S.M.; Lopez-Moreno, I.; Beguería, S.; Lorenzo-Lacruz, J.; Sanchez-Lorenzo, A.; García-Ruiz, J.M.; Azorin-Molina, C.; Morán-Tejada, E.; Revuelto, J.; Trigo, R.; et al. Evidence of increasing drought severity caused by temperature rise in southern Europe. *Environ. Res. Lett.* **2014**, *9*, 044001. [\[CrossRef\]](#)
25. Stahl, K.; Kohn, I.; Blauhut, V.; Urquijo, J.; De Stefano, L.; Acácio, V.; Dias, S.; Stagge, J.H.; Tallaksen, L.M.; Kampragou, E.; et al. Impacts of European drought events: Insights from an international database of text-based reports. *Nat. Hazards Earth Syst. Sci.* **2016**, *16*, 801–819. [\[CrossRef\]](#)
26. Tigchelaar, M.; Battisti, D.S.; Naylor, R.L.; Ray, D. Future warming increases probability of globally synchronized maize production shocks. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 6644–6649. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Allen, C.D.; Breshears, D.D.; McDowell, N.G. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **2015**, *6*, 1–55. [\[CrossRef\]](#)
28. Payus, C.; Huey, L.A.; Adnan, F.; Rimba, A.B.; Mohan, G.; Chapagain, S.K.; Roder, G.; Gasparatos, A.; Fukushi, K. Impact of Extreme Drought Climate on Water Security in North Borneo: Case Study of Sabah. *Water* **2020**, *12*, 1135. [\[CrossRef\]](#)
29. Murray-Tortarolo, G.N.; Jaramillo, V.J. The impact of extreme weather events on livestock populations: The case of the 2011 drought in Mexico. *Clim. Chang.* **2019**, *153*, 79–89. [\[CrossRef\]](#)
30. Auerbach, R.; Piek, H.; Battersby, J.; Devereux, S.; Olivier, N. *The Likely Impact of the 2015–2018 Drought in South Africa: Lessons from the 2008 Food Price Crisis and Future Implications*; Cabi: Wallingford, UK, 2020; pp. 81–99.

31. Wilhite, D.A.; Svoboda, M.D.; Hayes, M.J. Understanding the complex impacts of drought: A key to enhancing drought mitigation and preparedness. *Water Resour. Manag.* **2007**, *21*, 763–774. [\[CrossRef\]](#)
32. Alley, W.M. The Palmer drought severity index: Limitations and assumptions. *J. Appl. Meteorol. Climatol.* **1984**, *23*, 1100–1109. [\[CrossRef\]](#)
33. McKee Thomas, B.; Doesken, N.J.; Kleist, J. The relationship of drought frequency and duration to time scales. In Proceedings of the 8th Conference on Applied Climatology, Anaheim, VA, USA, 17–22 January 1993; Volume 17, pp. 179–183.
34. Vicente-Serrano, S.M.; Beguería, S.; Lopez-Moreno, I. A Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation Evapotranspiration Index. *J. Clim.* **2010**, *23*, 1696–1718. [\[CrossRef\]](#)
35. Palmer, W.C. Keeping Track of Crop Moisture Conditions, Nationwide: The New Crop Moisture Index. *Weatherwise* **1968**, *21*, 156–161. [\[CrossRef\]](#)
36. Meyer, S.J.; Hubbard, K.; Wilhite, D.A. A Crop-Specific Drought Index for Corn: II. Application in Drought Monitoring and Assessment. *Agron. J.* **1993**, *85*, 396–399. [\[CrossRef\]](#)
37. Gommers, R.; Petrassi, F. *Rainfall Variability and Drought in Sub-Saharan Africa*; Food and agriculture organization of the United Nations: Rome, Italy, 1996.
38. Weghorst, K.M. *The Reclamation Drought Index: Guidelines and Practical Applications*; Bureau of Reclamation: Denver, CO, USA, 1996.
39. Byun, H.-R.; Wilhite, D.A. Objective quantification of drought severity and duration. *J. Clim.* **1999**, *12*, 2747–2756. [\[CrossRef\]](#)
40. Bhalme, H.N.; Mooley, D.A. Large-scale droughts/floods and monsoon circulation. *Mon. Weather Rev. Am. Meteorol. Soc.* **1980**, *108*, 8. [\[CrossRef\]](#)
41. Shukla, S.; Wood, A. Use of a standardized runoff index for characterizing hydrologic drought. *Geophys. Res. Lett.* **2008**, *35*. [\[CrossRef\]](#)
42. Hong, X.W.; Lu, R.Y.; Li, S.L. Amplified summer warming in Europe–West Asia and Northeast Asia after the mid-1990s. *Environ. Res. Lett.* **2017**, *12*, 9. [\[CrossRef\]](#)
43. Hanel, M.; Rakovec, O.; Markonis, Y.; Máca, P.; Samaniego, L.; Kysely, J.; Kumar, R. Revisiting the recent European droughts from a long-term perspective. *Sci. Rep.* **2018**, *8*, 9499. [\[CrossRef\]](#) [\[PubMed\]](#)
44. Spinoni, J.; Vogt, J.V.; Naumann, G.; Barbosa, P.; Dosio, A. Will drought events become more frequent and severe in Europe? *Int. J. Climatol.* **2018**, *38*, 1718–1736. [\[CrossRef\]](#)
45. Bartholy, J.; Pongracz, R.; Hollósi, B. Analysis of projected drought hazards for Hungary. *Adv. Geosci.* **2013**, *35*, 61–66. [\[CrossRef\]](#)
46. Mezösi, G.; Bata, T.; Meyer, B.C.; Blanka, V.; Ladányi, Z. Climate Change Impacts on Environmental Hazards on the Great Hungarian Plain, Carpathian Basin. *Int. J. Disaster Risk Sci.* **2014**, *5*, 136–146. [\[CrossRef\]](#)
47. Ceglar, A.; Croitoru, A.-E.; Cuxart, J.; Djurdjevic, V.; Güttler, I.; Ivančan-Picek, B.; Jug, D.; Lakatos, M.; Weidinger, T. PannEx: The pannonian basin experiment. *Clim. Serv.* **2018**, *11*, 78–85. [\[CrossRef\]](#)
48. Gálos, B.; Lorenz, P.; Jacob, D. Will dry events occur more often in Hungary in the future? *Environ. Res. Lett.* **2007**, *2*. [\[CrossRef\]](#)
49. Mohammed, S.; Alsafadi, K.; Daher, H.; Gombos, B.; Mahmood, S.; Harsányi, E. Precipitation pattern changes and response of vegetation to drought variability in the eastern Hungary. *Bull. Natl. Res. Cent.* **2020**, *44*, 1–10. [\[CrossRef\]](#)
50. CARBATCLIM. Available online: <http://www.carpatclim-eu.org/pages/about/> (accessed on 9 September 2021).
51. Szalai, S.; Vogt, J. CARPATCLIM—high resolution gridded database of the Carpathian Region and calculation of drought indices as a contribution to the European Drought Observatory. In Proceedings of the WRCF Conference, Denver, CO, USA, 24–28 October 2011; pp. 24–28.
52. Alsafadi, K.; Mohammed, S.A.; Ayugi, B.; Sharaf, M.; Harsányi, E. Spatial–temporal evolution of drought characteristics over Hungary between 1961 and 2010. *Pure Appl. Geophys.* **2020**, *177*, 3961–3978. [\[CrossRef\]](#)
53. Mann, H.B. Nonparametric tests against trend. *Econom. J. Econom. Soc.* **1945**, *1*, 245–259. [\[CrossRef\]](#)
54. Sen, P.K. Estimates of the regression coefficient based on Kendall’s tau. *J. Am. Stat. Assoc.* **1968**, *1*, 1379–1389. [\[CrossRef\]](#)
55. Stagge, J.H.; Tallaksen, L.M.; Gudmundsson, L.; Van Loon, A.F.; Stahl, K. Candidate distributions for climatological drought indices (SPI and SPEI). *Int. J. Climatol.* **2015**, *35*, 4027–4040. [\[CrossRef\]](#)
56. Yang, J.; Chang, J.X.; Wang, Y.M.; Li, Y.Y.; Hu, H.; Chen, Y.T.; Huang, Q.; Yao, J. Comprehensive drought characteristics analysis based on a nonlinear multivariate drought index. *J. Hydrol.* **2018**, *557*, 651–667. [\[CrossRef\]](#)
57. Zargar, A.; Sadiq, R.; Naser, G.; Khan, F.I. A review of drought indices. *Environ. Rev.* **2011**, *19*, 333–349. [\[CrossRef\]](#)
58. Chang, K.Y.; Xu, L.; Starr, G. A drought indicator reflecting ecosystem responses to water availability: The normalized ecosystem drought index. *Agric. For. Meteorol.* **2018**, *250*, 102–117. [\[CrossRef\]](#)
59. Sims, A.P.; Niyogi, D.D.S.; Raman, S. Adopting drought indices for estimating soil moisture: A North Carolina case study. *Geophys. Res. Lett.* **2002**, *29*, 24-1–24-4. [\[CrossRef\]](#)
60. Ogunrinde, A.T.; Oguntunde, P.G.; Akinwumiju, A.S.; Fasinmirin, J.T. Analysis of recent changes in rainfall and drought indices in Nigeria, 1981–2015. *Hydrol. Sci. J.* **2019**, *64*, 1755–1768. [\[CrossRef\]](#)
61. Hao, Z.; Singh, V.P. Drought characterization from a multivariate perspective: A review. *J. Hydrol.* **2015**, *527*, 668–678. [\[CrossRef\]](#)
62. Park, J.; Sung, J.H.; Lim, Y.-J.; Kang, H.-S. Introduction and application of non-stationary standardized precipitation index considering probability distribution function and return period. *Theor. Appl. Clim.* **2018**, *136*, 529–542. [\[CrossRef\]](#)
63. Mohammed, S.; Alsafadi, K.; Al-Awadhi, T.; Sherief, Y.; Harsanyie, E.; El Kenawy, A.M. Space and time variability of meteorological drought in Syria. *Acta Geophys.* **2020**, *68*, 1877–1898. [\[CrossRef\]](#)

64. Potopová, V.; Štěpánek, P.; Možný, M.; Türkott, L.; Soukup, J. Performance of the standardised precipitation evapotranspiration index at various lags for agricultural drought risk assessment in the Czech Republic. *Agric. For. Meteorol.* **2015**, *202*, 26–38. [\[CrossRef\]](#)
65. Liu, X.; Pan, Y.; Zhu, X.; Yang, T.; Bai, J.; Sun, Z. Drought evolution and its impact on the crop yield in the North China Plain. *J. Hydrol.* **2018**, *564*, 984–996. [\[CrossRef\]](#)
66. Tigkas, D.; Vangelis, H.; Tsakiris, G. Drought characterisation based on an agriculture-oriented standardised precipitation index. *Theor. Appl. Climatol.* **2019**, *135*, 1435–1447. [\[CrossRef\]](#)
67. Sharma, A.; Goyal, M.K. Assessment of ecosystem resilience to hydroclimatic disturbances in India. *Glob. Chang. Biol.* **2017**, *24*, e432–e441. [\[CrossRef\]](#) [\[PubMed\]](#)
68. Dai, A.; Zhao, T.; Chen, J. Climate Change and Drought: A Precipitation and Evaporation Perspective. *Curr. Clim. Chang. Rep.* **2018**, *4*, 301–312. [\[CrossRef\]](#)
69. Berg, A.; Sheffield, J. Climate Change and Drought: The Soil Moisture Perspective. *Curr. Clim. Chang. Rep.* **2018**, *4*, 180–191. [\[CrossRef\]](#)
70. Cook, B.I.; Mankin, J.S.; Anchukaitis, K.J. Climate Change and Drought: From Past to Future. *Curr. Clim. Chang. Rep.* **2018**, *4*, 164–179. [\[CrossRef\]](#)
71. Wang, Z.; Zhong, R.; Lai, C.; Zeng, Z.; Lian, Y.; Bai, X. Climate change enhances the severity and variability of drought in the Pearl River Basin in South China in the 21st century. *Agric. For. Meteorol.* **2018**, *249*, 149–162. [\[CrossRef\]](#)
72. Pei, Z.; Fang, S.; Wang, L.; Yang, W. Comparative Analysis of Drought Indicated by the SPI and SPEI at Various Timescales in Inner Mongolia, China. *Water* **2020**, *12*, 1925. [\[CrossRef\]](#)
73. Stagge, J.H.; Kingston, D.G.; Tallaksen, L.M.; Hannah, D.M. Observed drought indices show increasing divergence across Europe. *Sci. Rep.* **2017**, *7*, 1–10. [\[CrossRef\]](#)
74. Kuwayama, Y.; Thompson, A.; Bernknopf, R.; Zaitchik, B.; Vail, P. Estimating the Impact of Drought on Agriculture Using the U.S. Drought Monitor. *Am. J. Agric. Econ.* **2018**, *101*, 193–210. [\[CrossRef\]](#)
75. Peña-Gallardo, M.; Vicente-Serrano, S.M.; Domínguez-Castro, F.; Beguería, S. The impact of drought on the productivity of two rainfed crops in Spain. *Nat. Hazards Earth Syst. Sci.* **2019**, *19*, 1215–1234. [\[CrossRef\]](#)
76. Ziolkowska, J.R. Socio-economic implications of drought in the agricultural sector and the state economy. *Economies* **2016**, *4*, 19. [\[CrossRef\]](#)
77. Nath, R.; Nath, D.; Li, Q.; Chen, W.; Cui, X. Impact of drought on agriculture in the Indo-Gangetic Plain, India. *Adv. Atmos. Sci.* **2017**, *34*, 335–346. [\[CrossRef\]](#)
78. Parsons, D.J.; Rey, D.; Tanguy, M.; Holman, I.P. Regional variations in the link between drought indices and reported agricultural impacts of drought. *Agric. Syst.* **2019**, *173*, 119–129. [\[CrossRef\]](#)
79. Ray, R.L.; Fares, A.; Risch, E. Effects of Drought on Crop Production and Cropping Areas in Texas. *Agric. Environ. Lett.* **2018**, *3*, 170037. [\[CrossRef\]](#)
80. Van Lanen, H.A.; Laaha, G.; Kingston, D.G.; Gauster, T.; Ionita, M.; Vidal, J.; Vlnas, R.; Tallaksen, L.M.; Stahl, K.; Hannaford, J.; et al. Hydrology needed to manage droughts: The 2015 European case. *Hydrol. Process.* **2016**, *30*, 3097–3104. [\[CrossRef\]](#)
81. Van Loon, A.F. Hydrological drought explained. *Wiley Interdiscip. Rev. Water* **2015**, *2*, 359–392. [\[CrossRef\]](#)
82. Herrera-Estrada, J.E.; Duffenbaugh, N.S.; Wagner, F.; Craft, A.; Sheffield, J. Response of electricity sector air pollution emissions to drought conditions in the western United States. *Environ. Res. Lett.* **2018**, *13*, 124032. [\[CrossRef\]](#)
83. Myrionidis, D.; Fotakis, D.; Ioannou, K.; Sgouropoulou, K. Comparison of ten notable meteorological drought indices on tracking the effect of drought on streamflow. *Hydrol. Sci. J.* **2018**, *63*, 2005–2019. [\[CrossRef\]](#)
84. Musolino, D.; de Carli, A.; Massarutto, A. Evaluation of socio-economic impact of drought events: The case of Po river basin. *Eur. Countrys.* **2017**, *9*, 163. [\[CrossRef\]](#)
85. Edwards, B.; Gray, M.; Hunter, B. The Impact of Drought on Mental Health in Rural and Regional Australia. *Soc. Indic. Res.* **2014**, *121*, 177–194. [\[CrossRef\]](#)
86. Yusa, A.; Berry, P.J.; Cheng, J.; Ogden, N.; Bonsal, B.; Stewart, R.; Waldick, R. Climate change, drought and human health in Canada. *Int. J. Environ. Res. Public Health* **2015**, *12*, 8359–8412. [\[CrossRef\]](#) [\[PubMed\]](#)
87. Ebi, K.L.; Bowen, K. Extreme events as sources of health vulnerability: Drought as an example. *Weather. Clim. Extrem.* **2016**, *11*, 95–102. [\[CrossRef\]](#)
88. Smith, T.; Fitchett, J.M. Drought challenges for nature tourism in the Sabi Sands Game Reserve in the eastern region of South Africa. *Afr. J. Range Forage Sci.* **2020**, *37*, 107–117. [\[CrossRef\]](#)
89. Lwoga, N.B.; Asubisiye, E. Effects of drought on cultural tourism: Selected cases of Maasai tourism groups surrounding Ta-rangire National Park in Tanzania. *J. Tour. Cult. Chang.* **2018**, *16*, 248–264. [\[CrossRef\]](#)
90. Rosselló, J.; Becken, S.; Gallego, M.S. The effects of natural disasters on international tourism: A global analysis. *Tour. Manag.* **2020**, *79*, 104080. [\[CrossRef\]](#)
91. Bordi, I.; Fraedrich, K.; Sutera, A. Observed drought and wetness trends in Europe: An update. *Hydrol. Earth Syst. Sci.* **2009**, *13*, 1519–1530. [\[CrossRef\]](#)
92. Lloyd-Hughes, B.; Saunders, M.A. A drought climatology for Europe. *Int. J. Climatol. A J. R. Meteorol. Soc.* **2002**, *22*, 1571–1592. [\[CrossRef\]](#)

93. Trnka, M.; Balek, J.; Zahradníček, P.; Eitzinger, J.; Formayer, H.; Turna, M.; Nejedlík, P.; Semerádová, D.; Hlavinka, P.; Brázdil, R. Drought trends over part of Central Europe between 1961 and 2014. *Climate Res.* **2016**, *70*, 143–160. [[CrossRef](#)]
94. Szabó, S.; Szopos, N.M.; Bertalan-Balázs, B.; László, E.; Milošević, D.D.; Conoscenti, C.; Lázár, I. Geospatial analysis of drought tendencies in the Carpathians as reflected in a 50-year time series. *Hung. Geogr. Bull.* **2019**, *68*, 269–282. [[CrossRef](#)]
95. Adrienn, V.S.; Janos, N. Effects of nutrition and water supply on the yield and grain protein content of maize hybrids. *Aust. J. Crop Sci.* **2012**, *6*, 381–390.
96. Paltineanu, C.R.; Mihailescu, I.F.; Prefac, Z.; Dragota, C.; Vasenciuc, F.; Claudia, N. Combining the standardized precipitation index and climatic water deficit in characterizing droughts: A case study in Romania. *Theor. Appl. Climatol.* **2009**, *97*, 219. [[CrossRef](#)]
97. Spinoni, J.; Antofie, T.; Barbosa, P.; Bihari, Z.; Lakatos, M.; Szalai, S.; Szentimrey, T.; Vogt, J. An overview of drought events in the Carpa-thian Region in 1961–2010. *Adv. Sci. Res.* **2013**, *10*, 21–32. [[CrossRef](#)]
98. Wang, W.; Zhu, Y.; Xu, R.; Liu, J. Drought severity change in China during 1961–2012 indicated by SPI and SPEI. *Nat. Hazards* **2014**, *75*, 2437–2451. [[CrossRef](#)]
99. Won, J.; Choi, J.; Lee, O.; Kim, S. Copula-based Joint Drought Index using SPI and EDDI and its application to climate change. *Sci. Total Environ.* **2020**, *744*, 140701. [[CrossRef](#)] [[PubMed](#)]
100. Chen, X.; Li, Y.; Yao, N.; Liu, D.L.; Javed, T.; Liu, C.; Liu, F. Impacts of multi-timescale SPEI and SMDI variations on winter wheat yields. *Agric. Syst.* **2020**, *185*, 102955. [[CrossRef](#)]
101. Ma, B.; Zhang, B.; Jia, L.; Huang, H. Conditional distribution selection for SPEI-daily and its revealed meteorological drought characteristics in China from 1961 to 2017. *Atmos. Res.* **2020**, *246*, 105108. [[CrossRef](#)]
102. Li, Y.; Ye, W.; Wang, M.; Yan, X. Climate change and drought: A risk assessment of crop-yield impacts. *Clim. Res.* **2009**, *39*, 31–46. [[CrossRef](#)]
103. Matiu, M.; Ankerst, D.P.; Menzel, A. Interactions between temperature and drought in global and regional crop yield variability during 1961–2014. *PLoS ONE* **2017**, *12*, 178339. [[CrossRef](#)]
104. Lobell, D.B.; Schlenker, W.; Costa-Roberts, J. Climate Trends and Global Crop Production Since 1980. *Science* **2011**, *333*, 616–620. [[CrossRef](#)] [[PubMed](#)]
105. Lobell, D.; Field, C.B. Global scale climate–crop yield relationships and the impacts of recent warming. *Environ. Res. Lett.* **2007**, *2*. [[CrossRef](#)]
106. Schlenker, W.; Roberts, M.J. Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 15594–15598. [[CrossRef](#)] [[PubMed](#)]
107. Hoerling, M.P.; Eischeid, J.; Kumar, A.; Leung, R.; Mariott, A.; Mo, K.; Schubert, S.; Seager, R. Causes and predictability of the 2012 Great Plains drought. *Bull. Amer. Meteor. Soc.* **2014**, *95*, 269–282. [[CrossRef](#)]
108. Foster, T.; Brozović, N.; Butler, A.P. Why well yield matters for managing agricultural drought risk. *Weather Clim. Extremes* **2015**, *10*, 11–19. [[CrossRef](#)]