

Supplementary

# Three-Year Variations in Criteria Atmospheric Pollutants and Their Relationship with Rainwater Chemistry in Karst Urban Region, Southwest China

Jie Zeng <sup>1</sup>, Xin Ge <sup>2</sup>, Qixin Wu <sup>2,\*</sup> and Shitong Zhang <sup>1</sup>

<sup>1</sup> Institute of Earth Sciences, China University of Geosciences (Beijing), Beijing 100083, China; zengjie@cugb.edu.cn (J.Z.); stongzhang0103@edu.cn (S.Z.)

<sup>2</sup> Key Laboratory of Karst Georesources and Environment, Ministry of Education, College of Resources and Environmental Engineering, Guizhou University, Guiyang 500025, China; april\_gexin@163.com

\* Correspondence: qxwu@gzu.edu.cn

**Table S1.** The annual mean contents of atmospheric pollutants and the Grade II standard values of CAAQS II and CAAQS III.

	Unit	CAAQS II-Grade II standard	CAAQS III-Grade II standard	2013	2014	2015
SO <sub>2</sub>	µg/m <sup>3</sup>	60	60	30	23	17
NO <sub>2</sub>	µg/m <sup>3</sup>	40	40	32	29	26
CO	mg/m <sup>3</sup>	4	4	0.78	0.77	0.70
PM <sub>2.5</sub>	µg/m <sup>3</sup>	-	35	51	46	38
PM <sub>10</sub>	µg/m <sup>3</sup>	100	70	81	70	58