


## Supplementary Material

# Onsite Chlorination of Greywater in a Vertical Flow Constructed Wetland—Significance of Trihalomethane Formation

Anat Bernstein, Hagar Siebner, Andrew Kaufman and Amit Gross \*

Department of Environmental Hydrology and Microbiology (EHM), The Zuckerberg Institute for Water Research (ZIWR), The Jacob Blaustein Institutes for Desert Research, Ben-Gurion University of the Negev, Israel; anatbern@bgu.ac.il (A.B.); siebner@bgu.ac.il (H.S.); andrewka@post.bgu.ac.il (A.K.)

\* Correspondence: amgross@bgu.ac.il



**Figure S1.** A recirculating vertical flow constructed wetland system

**Table S1.** Chlorine yield as THM for system S5 water that was incubated with 5 mg/l chlorine for 1 h. Yield was calculated as the ratio of chlorine recovered in THM versus chlorine consumed

Sample	Circulation time (h)	Residual chlorine after 1 h (mg/l)	Chlorine consumed in 1 h (mg/l)	Chlorine recovered in THM (µg/l)	Chlorine yield as THM (-)
B1	0	0.39	4.61	15.0	0.0033
B2	0.5	0.84	4.16	14.9	0.0036
B3	1	1.61	3.39	8.7	0.0026
B4	3	3.02	1.98	21.5	0.0109
B5	6	2.89	2.11	111.3	0.0527