

## Supplementary Material

# Feasibility study of using excess sludge fermentation broth as a co-metabolic carbon source for 2,4,6-Trichlorophenol degradation

Jianguang Wang <sup>1,2\*</sup>, Zhirong Sun <sup>1\*</sup>, Jun Li <sup>1</sup>

<sup>1</sup> National Engineering Laboratory for Advanced Municipal Wastewater Treatment and Reuse Technology, Beijing University of Technology, Beijing 100124, China

<sup>2</sup> PowerChina Huadong Engineering Corporation Limited, Hangzhou 311122, China

\* Correspondence: wjg909090@163.com (J.W.); zrsun@bjut.edu.cn (Z.S.)

Table S1: The DSBR Influent Trace Elements Concentrations

Number	Trace elements	Concentration(mg/L)
1	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	0.25
2	MgSO <sub>4</sub> ·7H <sub>2</sub> O	12.0
3	FeSO <sub>4</sub> ·7H <sub>2</sub> O	2.0
4	H <sub>3</sub> BO <sub>3</sub>	0.3 μg/L
5	CuSO <sub>4</sub> ·5H <sub>2</sub> O	0.06
6	KI	0.06
7	MnSO <sub>4</sub> ·H <sub>2</sub> O	0.24
8	Na <sub>2</sub> MoO <sub>4</sub> ·2H <sub>2</sub> O	0.12