

Article

On Novel Results about the Algebraic Properties of Symbolic 3-Plithogenic and 4-Plithogenic Real Square Matrices

Hamiyet Merkepçி 

Department of Mathematics, Gaziantep University, Gazinatep 27310, Turkey; hamiyetmerkepc@gmail.com; Tel.: +90-507-355-75-15

Abstract: Symbolic n-plithogenic sets are considered to be modern concepts that carry within their framework both an algebraic and logical structure. The concept of symbolic n-plithogenic algebraic rings is considered to be a novel generalization of classical algebraic rings with many symmetric properties. These structures can be written as linear combinations of many symmetric elements taken from other classical algebraic structures, where the square symbolic k-plithogenic real matrices are square matrices with real symbolic k-plithogenic entries. In this research, we will find easy-to-use algorithms for calculating the determinant of a symbolic 3-plithogenic/4-plithogenic matrix, and for finding its inverse based on its classical components, and even for diagonalizing matrices of these types. On the other hand, we will present a new algorithm for calculating the eigenvalues and eigenvectors associated with matrices of these types. Also, the exponent of symbolic 3-plithogenic and 4-plithogenic real matrices will be presented, with many examples to clarify the novelty of this work.

Keywords: symbolic 3-plithogenic rings; symbolic 3-plithogenic real matrices; symbolic 4-plithogenic rings; symbolic 4-plithogenic real matrices; the diagonalization problem



Citation: Merkepçி, H. On Novel Results about the Algebraic Properties of Symbolic 3-Plithogenic and 4-Plithogenic Real Square Matrices. *Symmetry* **2023**, *15*, 1494. <https://doi.org/10.3390/sym15081494>

Academic Editor: Qing-Wen Wang

Received: 20 June 2023

Revised: 14 July 2023

Accepted: 24 July 2023

Published: 27 July 2023



Copyright: © 2023 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Symbolic n-plithogenic sets were propounded by Smarandache in [1–3] to be a novel generalization of classical sets. These sets play a key role in the generalization of classical algebraic structures to new versions possessing several characteristic properties; in these we can see many symbolic n-plithogenic algebraic structures, such as symbolic 2-plithogenic structures [4–8], and symbolic 3-plithogenic algebraic structures [9–11]. For example, symbolic 2-plithogenic rings were used to generalize vector spaces and modules into symbolic 2-plithogenic spaces and modules. The same work was presented for symbolic 3-plithogenic rings [12,13]. Symbolic 2-plithogenic matrices were defined and studied in [14]; these matrices consist of symbolic 2-plithogenic real entries. These matrices are recognized as a similar structure of refined neutrosophic matrices and structures [15–24]. In matrix theory, it is very important to deal with the exponents of matrices and their related problems, such as how to diagonalize a matrix, and how to compute eigenvalues and eigenvectors.

From this point of view, we continue the previous efforts presented in order to further our understanding of the behavior of symbolic n-plithogenic matrices, and present some algorithms for computing inverses, eigen values and vectors, and the diagonalization of square matrices using symbolic 3-plithogenic entries; we also present some algorithms for computing inverses, eigen values and vectors, and the diagonalization of square matrices using symbolic 4-plithogenic entries.

Our study opens the door to many future applications of this type of matrices, especially those related to the problem of representation through linear transformations, or the representation of algebraic groups by matrices of this type.

2. Preliminaries

Definition 1. The ring of symbolic of 3-plithogenic real numbers is defined as follows:

$$3 - SP_R = \left\{ x + yP_1 + zP_2 + tP_3; P_i \times P_j = P_{\max(i,j)}, P_i^2 = P_i \right\}$$

Addition on 3-SP_R is defined as follows:

$$[x_0 + x_1P_1 + x_2P_2 + x_3P_3] + [y_0 + y_1P_1 + y_2P_2 + y_3P_3] = (x_0 + y_0) + (x_1 + y_1)P_1 + (x_2 + y_2)P_2 + (x_3 + y_3)P_3$$

Multiplication on 3 - SP_R is defined as follows:

$$\begin{aligned} & [x_0 + x_1P_1 + x_2P_2 + x_3P_3][y_0 + y_1P_1 + y_2P_2 + y_3P_3] \\ &= x_0y_0 + (x_0y_1 + x_1y_0 + x_1y_1)P_1 + (x_0y_2 + x_2y_1 + x_2y_2 + x_2y_0 + x_1y_2)P_2 \\ &+ (x_0y_3 + x_1y_3 + x_2y_3 + x_3y_3 + x_3y_0 + x_3y_1 + x_3y_2)P_3 \end{aligned}$$

Remark 1. If we Let $X = x_0 + x_1P_1 + x_2P_2 + x_3P_3 \in 3 - SP_R$, we have the following:

$$\begin{aligned} X \text{ is invertible if, and only if, } & x_0 \neq 0, x_0 + x_1 \neq 0, x_0 + x_1 + x_2 \neq 0, x_0 + x_1 + x_2 + x_3 \neq 0, \\ \text{and } X^{-1} = \frac{1}{X} = \frac{1}{x_0} & + \left[\frac{1}{x_0+x_1} - \frac{1}{x_0} \right] P_1 + \left[\frac{1}{x_0+x_1+x_2} - \frac{1}{x_0+x_1} \right] P_2 + \left[\frac{1}{x_0+x_1+x_2+x_3} - \frac{1}{x_0+x_1+x_2} \right] P_3 \\ \text{For } n \in \mathbb{N}, X^n = x_0^n & + [(x_0 + x_1)^n - x_0^n]P_1 + [(x_0 + x_1 + x_2)^n - (x_0 + x_1)^n]P_2 + \\ & [(x_0 + x_1 + x_2 + x_3)^n - (x_0 + x_1 + x_2)^n]P_3. \end{aligned}$$

Definition 2. The ring of symbolic of 4-plithogenic real numbers is defined as follows:

$$4 - SP_R = \left\{ x + yP_1 + zP_2 + tP_3 + lP_4; P_i \times P_j = P_{\max(i,j)}, P_i^2 = P_i \right\}$$

Addition on 4 - SP_R is defined as follows:

$$[x_0 + x_1P_1 + x_2P_2 + x_3P_3 + x_4P_4] + [y_0 + y_1P_1 + y_2P_2 + y_3P_3 + y_4P_4] = (x_0 + y_0) + (x_1 + y_1)P_1 + (x_2 + y_2)P_2 + \\ (x_3 + y_3)P_3 + (x_4 + y_4)P_4$$

Multiplication on 4-SP_R is defined as follows:

$$\begin{aligned} & [x_0 + x_1P_1 + x_2P_2 + x_3P_3 + x_4P_4][y_0 + y_1P_1 + y_2P_2 + y_3P_3 + y_4P_4] = x_0y_0 + (x_0y_1 + x_1y_0 + x_1y_1)P_1 \\ & + (x_0y_2 + x_2y_1 + x_2y_2 + x_2y_0 + x_1y_2)P_2 + (x_0y_3 + x_1y_3 + x_2y_3 + x_3y_3 + x_3y_0 + x_3y_1 + x_3y_2)P_3 + \\ & (x_0y_4 + x_1y_4 + x_2y_4 + x_3y_4 + x_3y_0 + x_4y_1 + x_4y_2 + x_4y_3 + x_4y_4)P_4. \end{aligned}$$

3. Main Discussion

3.1. Symbolic 3-Plithogenic Matrices

Definition 3. A square symbolic 3-plithogenic matrix is a matrix with symbolic 3-plithogenic entries.

Example 1.

$$A = \begin{pmatrix} 4 + P_1 + P_2 - 2P_3 & 1 + P_3 & 3 - P_1 + 5P_3 \\ 5 - P_1 + P_2 & 2 - P_1 & 7P_1 + 4P_2 - P_3 \\ 6P_2 + 4P_3 & 5 + 11P_2 + P_3 & P_1 + P_3 \end{pmatrix}$$

is a symbolic 3-plithogenic 3×3 real matrix.

Remark 2. Any symbolic 3-plithogenic matrix can be written as follows: $A = A_0 + A_1P_1 + A_2P_2 + A_3P_3$.

For example, the matrix defined in the previous example can be written as follows:

$$A = \begin{pmatrix} 4 & 1 & 3 \\ 5 & 2 & 0 \\ 0 & 5 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 & -1 \\ -1 & -1 & 7 \\ 0 & 0 & 1 \end{pmatrix} P_1 + \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 4 \\ 6 & 11 & 0 \end{pmatrix} P_2 + \begin{pmatrix} -2 & 1 & 5 \\ 0 & 0 & -1 \\ 4 & 1 & 1 \end{pmatrix} P_3$$

We denote the set of all symbolic 3-plithogenic n -square matrices by $(3 - SP_M)$. It is clear that $(3 - SP_M, +, .)$ is a ring.

Theorem 1. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 \in 3 - SP_M$, then the following is true:

1. S is invertible if, and only if, $S_0, S_0 + S_1, S_0 + S_1 + S_2, S_0 + S_1 + S_2 + S_3$ are invertible.
2. $S^{-1} = S_0^{-1} + [(S_0 + S_1)^{-1} - S_0^{-1}]P_1 + [(S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}]P_2 + [(S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}]P_3$
3. For $n \in N$, $S^n = S_0^n + [(S_0 + S_1)^n - S_0^n]P_1 + [(S_0 + S_1 + S_2)^n - (S_0 + S_1)^n]P_2 + [(S_0 + S_1 + S_2 + S_3)^n - (S_0 + S_1 + S_2)^n]P_3$
4. $e^S = e^{S_0} + [e^{S_0+S_1} - e^{S_0}]P_1 + [e^{S_0+S_1+S_2} - e^{S_0+S_1}]P_2 + [e^{S_0+S_1+S_2+S_3} - e^{S_0+S_1+S_2}]P_3$.

Proof of Theorem 1. (1,2). Assume that $S_0, S_0 + S_1, S_0 + S_1 + S_2, S_0 + S_1 + S_2 + S_3$ are invertible. We have the following:

$$K = S_0^{-1} + [(S_0 + S_1)^{-1} - S_0^{-1}]P_1 + [(S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}]P_2 + [(S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}]P_3$$

Let us compute the following:

$$\begin{aligned} SK &= S_0 S_0^{-1} + [S_0(S_0 + S_1)^{-1} - S_0 S_0^{-1} + S_1 S_0^{-1} + S_1(S_0 + S_1)^{-1} - S_1 S_0^{-1}]P_1 \\ &\quad + [S_0(S_0 + S_1 + S_2)^{-1} - S_0(S_0 + S_1)^{-1} + S_1(S_0 + S_1 + S_2)^{-1} \\ &\quad - S_1(S_0 + S_1)^{-1} + S_2(S_0 + S_1 + S_2)^{-1} - S_2(S_0 + S_1)^{-1} + S_2 S_0^{-1} \\ &\quad + S_2(S_0 + S_1)^{-1} - S_2 S_0^{-1}]P_2 + \\ [S_0(S_0 + S_1 + S_2 &+ S_3)^{-1} - S_0(S_0 + S_1 + S_2)^{-1} + S_1(S_0 + S_1 + S_2 + S_3)^{-1} \\ &- S_1(S_0 + S_1 + S_2)^{-1} + S_2(S_0 + S_1 + S_2 + S_3)^{-1} - S_2(S_0 + S_1 + S_2)^{-1} \\ &+ S_3(S_0 + S_1 + S_2 + S_3)^{-1} - S_3(S_0 + S_1 + S_2)^{-1} + S_3 S_0^{-1} + S_3(S_0 + S_1)^{-1} \\ &- S_3 S_0^{-1} + S_3(S_0 + S_1 + S_2 + S_3)^{-1} - S_3(S_0 + S_1)^{-1}]P_3 = \\ U_{n \times n} + [O_{n \times n}]P_1 &+ [O_{n \times n}]P_2 + [O_{n \times n}]P_3 = U_{n \times n} \end{aligned}$$

Thus, S is invertible and $S^{-1} = K$.

For the converse, if we suppose that S is invertible, then there exists $K = K_0 + K_1 P_1 + K_2 P_2 + K_3 P_3 \in 3 - SP_M$ such that $S \times K = U_{n \times n}$. This is equivalent to the following:

$$\left\{ \begin{array}{l} S_0 K_0 = U_{n \times n} \quad (1) \\ S_0 K_1 + S_1 K_0 + S_1 K_1 = O_{n \times n} \quad (2) \\ S_0 K_2 + S_1 K_2 + S_2 K_0 + S_2 K_1 + S_2 K_2 = O_{n \times n} \quad (3) \\ S_0 K_3 + S_1 K_3 + S_2 K_3 + S_3 K_0 + S_3 K_1 + S_3 K_2 = O_{n \times n} \quad (4) \end{array} \right.$$

Equation (1) implies that S_0 is invertible and $S_0^{-1} = K_0$.

By adding (1) to (2), we obtain $(S_0 + S_1)(K_0 + K_1) = U_{n \times n}$, so that $S_0 + S_1$ is invertible and $(S_0 + S_1)^{-1} = K_0 + K_1$, hence $K_1 = (S_0 + S_1)^{-1} - S_0^{-1}$.

By adding (1) to (2) to (3), we obtain $(S_0 + S_1 + S_2)(K_0 + K_1 + K_2) = U_{n \times n}$, so that $S_0 + S_1 + S_2$ is invertible and $(S_0 + S_1 + S_2)^{-1} = K_0 + K_1 + K_2$, hence $K_2 = (S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}$.

By adding (1) to (2) to (3) to (4), we obtain $(S_0 + S_1 + S_2 + S_3)(K_0 + K_1 + K_2 + K_3) = U_{n \times n}$, so that $S_0 + S_1 + S_2 + S_3$ is invertible and $(S_0 + S_1 + S_2 + S_3)^{-1} = K_0 + K_1 + K_2 + K_3$, hence $K_3 = (S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}$.

(3). For $n = 1$, it holds directly. We suppose that it is true. For $n = k$, we must prove it for $k + 1$.

$$\begin{aligned} S^{k+1} = SS^k &= S_0 S_0^k + \left[S_0(S_0 + S_1)^k - S_0 S_0^k + S_1 S_0^k + S_1(S_0 + S_1)^k - S_1 S_0^k \right] P_1 \\ &\quad + \left[S_0(S_0 + S_1 + S_2)^k - S_0(S_0 + S_1)^k + S_1(S_0 + S_1 + S_2)^k - S_1(S_0 + S_1)^k + S_2(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. - S_2(S_0 + S_1)^k + S_2 S_0^k + S_2(S_0 + S_1)^k - S_2 S_0^k \right] P_2 \\ &\quad + \left[S_0(S_0 + S_1 + S_2 + S_3)^k - S_0(S_0 + S_1 + S_2)^k + S_1(S_0 + S_1 + S_2 + S_3)^k - S_1(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. + S_2(S_0 + S_1 + S_2 + S_3)^k - S_2(S_0 + S_1 + S_2)^k + S_3(S_0 + S_1 + S_2 + S_3)^k - S_3(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. + S_3 S_0^k + S_3(S_0 + S_1)^k - S_3 S_0^k + S_3(S_0 + S_1 + S_2 + S_3)^k - S_3(S_0 + S_1)^k \right] P_3 \\ &= S_0^{k+1} + \left[(S_0 + S_1)(S_0 + S_1)^k - S_0 S_0^k \right] P_1 \\ &\quad + \left[(S_0 + S_1 + S_2)(S_0 + S_1 + S_2)^k - (S_0 + S_1)(S_0 + S_1)^k \right] P_2 \\ &\quad + \left[(S_0 + S_1 + S_2 + S_3)(S_0 + S_1 + S_2 + S_3)^k - (S_0 + S_1 + S_2)(S_0 + S_1 + S_2)^k \right] P_3 \\ &= S_0^{k+1} + \left[(S_0 + S_1)^{k+1} - S_0 S_0^{k+1} \right] P_1 + \left[(S_0 + S_1 + S_2)^{k+1} - (S_0 + S_1)^{k+1} \right] P_2 \\ &\quad + \left[(S_0 + S_1 + S_2 + S_3)^{k+1} - (S_0 + S_1 + S_2)^{k+1} \right] P_3 \end{aligned}$$

Therefore, the proof holds via induction.

$$(4). e^S = \sum_{n=0}^{\infty} \frac{S^n}{n!} = \sum_{n=0}^{\infty} \frac{S_0^n}{n!} + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1)^n}{n!} - \sum_{n=0}^{\infty} \frac{S_0^n}{n!} \right] P_1 + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2)^n}{n!} - \sum_{n=0}^{\infty} \frac{(S_0 + S_1)^n}{n!} \right] P_2 + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2 + S_3)^n}{n!} - \sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2)^n}{n!} \right] P_3 = e^{S_0} + [e^{S_0 + S_1} - e^{S_0}] P_1 + [e^{S_0 + S_1 + S_2} - e^{S_0 + S_1}] P_2 + [e^{S_0 + S_1 + S_2 + S_3} - e^{S_0 + S_1 + S_2}] P_3. \quad \square$$

Definition 4. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 \in 3 - SP_M$, then we define the following:

$$\det(S) = \det(S_0) + [\det(S_0 + S_1) - \det(S_0)]P_1 + [\det(S_0 + S_1 + S_2) - \det(S_0 + S_1)]P_2 + [\det(S_0 + S_1 + S_2 + S_3) - \det(S_0 + S_1 + S_2)]P_3$$

Theorem 2. S is invertible if, and only if, $\det(S)$ is invertible in $3 - SP_R$.

The proof holds via a similar discussion of the 2-plithogenic case.

Example 2. Consider the following matrix:

$$\begin{aligned} S &= \begin{pmatrix} 1 + P_1 + P_2 + P_3 & 5 + P_1 - 2P_2 \\ 1 + 4P_1 - P_2 + 3P_3 & P_1 + P_3 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 5 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix} P_1 + \begin{pmatrix} 1 & -2 \\ -1 & 0 \end{pmatrix} P_2 + \begin{pmatrix} 1 & 0 \\ 3 & 1 \end{pmatrix} P_3 \end{aligned}$$

We have $\det(S_0) = -5$, $\det(S_0 + S_1) = \det\begin{pmatrix} 2 & 6 \\ 5 & 1 \end{pmatrix} = -28$, $\det(S_0 + S_1 + S_2) = \det\begin{pmatrix} 3 & 4 \\ 4 & 1 \end{pmatrix} = -13$, $\det(S_0 + S_1 + S_2 + S_3) = \det\begin{pmatrix} 4 & 4 \\ 7 & 2 \end{pmatrix} = -20$.
Hence,

$$\det(S) = -5 + [-28 + 5]P_1 + [-13 + 28]P_2 + [-20 + 13]P_3 = -5 - 23P_1 + 15P_2 - 7P_3$$

Since $\det(S)$ is invertible, then S is invertible.

$$S_0^{-1} = \frac{-1}{5} \begin{pmatrix} 0 & -5 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ \frac{1}{5} & -\frac{1}{5} \end{pmatrix}$$

$$(S_0 + S_1)^{-1} = \frac{-1}{28} \begin{pmatrix} 1 & -6 \\ -5 & 2 \end{pmatrix} = \begin{pmatrix} -\frac{1}{28} & \frac{6}{28} \\ \frac{5}{28} & \frac{-2}{28} \end{pmatrix}$$

$$(S_0 + S_1 + S_2)^{-1} = \frac{-1}{13} \begin{pmatrix} 1 & -4 \\ -4 & 3 \end{pmatrix} = \begin{pmatrix} -\frac{1}{13} & \frac{4}{13} \\ \frac{4}{13} & \frac{-3}{13} \end{pmatrix}$$

$$(S_0 + S_1 + S_2 + S_3)^{-1} = \frac{-1}{20} \begin{pmatrix} 2 & -4 \\ -7 & 4 \end{pmatrix} = \begin{pmatrix} -\frac{2}{20} & \frac{4}{20} \\ \frac{7}{20} & \frac{-4}{20} \end{pmatrix}$$

$$S^{-1} = S_0^{-1} + [(S_0 + S_1)^{-1} - S_0^{-1}]P_1 + [(S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}]P_2 + [(S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}]P_3$$

$$S^{-1} = \begin{pmatrix} 0 & 1 \\ \frac{1}{5} & -\frac{1}{5} \end{pmatrix} + \left[\begin{pmatrix} -\frac{1}{28} & \frac{-22}{28} \\ \frac{-3}{28} & \frac{18}{28} \end{pmatrix} \right] P_1 + \left[\begin{pmatrix} -\frac{15}{364} & \frac{34}{364} \\ \frac{47}{364} & \frac{-58}{364} \end{pmatrix} \right] P_2 + \left[\begin{pmatrix} -\frac{6}{260} & \frac{-28}{260} \\ \frac{11}{260} & \frac{8}{260} \end{pmatrix} \right] P_3 =$$

$$\begin{pmatrix} -\frac{1}{28}P_1 - \frac{15}{364}P_2 - \frac{6}{260}P_3 & 1 - \frac{22}{28}P_1 + \frac{34}{364}P_2 - \frac{28}{260}P_3 \\ \frac{1}{5} - \frac{3}{28}P_1 + \frac{47}{364}P_2 + \frac{11}{260}P_3 & -\frac{1}{5} + \frac{18}{28}P_1 + \frac{58}{364}P_2 + \frac{8}{260}P_3 \end{pmatrix}$$

3.2. Symbolic 3-Plithogenic Eigen Values/Vectors

Theorem 3. Let $S = S_0 + S_1P_1 + S_2P_2 + S_3P_3 \in 3 - SP_M$, then $A = a_0 + a_1P_1 + a_2P_2 + a_3P_3$ is an eigen value of S if, and only if, a_0 eigen value of S_0 , $a_0 + a_1$ eigen value of $S_0 + S_1$, $a_0 + a_1 + a_2$ eigen value of $S_0 + S_1 + S_2$, $a_0 + a_1 + a_2 + a_3$ eigen value of $S_0 + S_1 + S_2 + S_3$.

In addition, $X = X_0 + X_1P_1 + X_2P_2 + X_3P_3$ is the corresponding eigen vector of A if, and only if, X_0 eigen vector of a_0 , $X_0 + X_1$ eigen vector of $a_0 + a_1$, $X_0 + X_1 + X_2$ eigen vector of $a_0 + a_1 + a_2$, $X_0 + X_1 + X_2 + X_3$ eigen vector of $a_0 + a_1 + a_2 + a_3$.

Proof of Theorem 3. From the equation $AS = AX$, we can obtain:

$$\left\{ \begin{array}{l} a_0S_0 = a_0X_0 \quad (5) \\ a_0S_1 + a_1S_0 + a_1S_1 = a_0X_1 + a_1X_0 + a_1X_1 \quad (6) \\ a_0S_2 + a_1S_2 + a_2S_1 + a_2S_0 = a_0X_2 + a_1X_2 + a_2X_2 + a_2X_1 + a_2X_0 \quad (7) \\ a_0S_3 + a_1S_3 + a_2S_3 + a_3S_3 + a_3S_0 + a_3S_1 + a_3S_2 = a_0X_3 + a_1X_3 + a_2X_3 + a_3X_3 + a_3X_0 + a_3X_1 + a_3X_2 \quad (8) \end{array} \right.$$

Equation (5) implies that a_0 is the eigen value of S_0 , with X_0 as the eigen vector.

By adding (5) to (6), we obtain $(a_0 + a_1)(S_0 + S_1) = (a_0 + a_1)(X_0 + X_1)$, which means that $a_0 + a_1$ is the eigen value of $S_0 + S_1$, with $X_0 + X_1$ as the eigen vector.

By adding (5) to (6) to (7), we obtain $(a_0 + a_1 + a_2)(S_0 + S_1 + S_2) = (a_0 + a_1 + a_2)(X_0 + X_1 + X_2)$, which means that $a_0 + a_1 + a_2$ is the eigen value of $S_0 + S_1 + S_2$, with $X_0 + X_1 + X_2$ as the eigen vector.

By adding (5) to (6) to (7) to (8), we obtain $(a_0 + a_1 + a_2 + a_3)(S_0 + S_1 + S_2 + S_3) = (a_0 + a_1 + a_2 + a_3)(X_0 + X_1 + X_2 + X_3)$, which means that $a_0 + a_1 + a_2 + a_3$ is the eigen value of $S_0 + S_1 + S_2 + S_3$, with $X_0 + X_1 + X_2 + X_3$ as the eigen vector.

Therefore, the proof is complete. \square

Example 3. Consider the following matrix:

$$S = \begin{pmatrix} 3 + 2P_1 - P_3 & 4 + P_2 - 3P_3 \\ 2P_1 + P_2 + P_3 & 5 + P_1 \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 0 & 5 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 2 & 1 \end{pmatrix}P_1 + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}P_2 + \begin{pmatrix} -1 & -3 \\ 1 & 0 \end{pmatrix}P_3 = A_0 + A_1P_1 + A_2P_2 + A_3P_3$$

The eigen values of $A_0 = \begin{pmatrix} 3 & 4 \\ 0 & 5 \end{pmatrix}$ are $\{3, 5\}$.

The eigen values of $A_0 + A_1 = \begin{pmatrix} 2 & 0 \\ 2 & 1 \end{pmatrix}$ are $\left\{ \frac{11+\sqrt{33}}{2}, \frac{11-\sqrt{33}}{2} \right\}$.

The eigen values of $A_0 + A_1 + A_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ are $\left\{ \frac{11+\sqrt{61}}{2}, \frac{11-\sqrt{61}}{2} \right\}$.

The eigen values of $A_0 + A_1 + A_2 + A_3 = \begin{pmatrix} -1 & -3 \\ 1 & 0 \end{pmatrix}$ are $\{8, 2\}$.

The eigen values of the symbolic 3-plithogenic matrix A are as follows:

$$U_1 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_2 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_3 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_4 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_5 = 3 + \left(\frac{11-\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11-\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_6 = 3 + \left(\frac{11-\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11-\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11-\sqrt{61}}{2} \right) P_3$$

$$U_7 = 3 + \left(\frac{11-\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11-\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11-\sqrt{61}}{2} \right) P_3$$

$$U_8 = 3 + \left(\frac{11-\sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11-\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11-\sqrt{61}}{2} \right) P_3$$

$$U_9 = 5 + \left(\frac{11+\sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_{10} = 5 + \left(\frac{11+\sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11+\sqrt{61}}{2} \right) P_3$$

$$U_{11} = 5 + \left(\frac{11+\sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11-\sqrt{61}}{2} \right) P_3$$

$$U_{12} = 5 + \left(\frac{11+\sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11-\sqrt{61}}{2} \right) P_3$$

$$U_{13} = 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 + \sqrt{61}}{2} \right) P_3$$

$$U_{14} = 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 + \sqrt{61}}{2} \right) P_3$$

$$U_{15} = 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 - \sqrt{61}}{2} \right) P_3$$

$$U_{16} = 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 - \sqrt{61}}{2} \right) P_3$$

Remark 3. If A has n eigen values for $0 \leq i \leq s$, then A has n^4 eigen values.

3.3. Symbolic 3-Plithogenic Diagonalization

Theorem 4. Let $T = T_0 + T_1P_1 + T_2P_2 + T_3P_3$ be an n -square symbolic 3-plithogenic matrix, then T is diagonalizable if, and only if, $T_0, T_0 + T_1, T_0 + T_1 + T_2, T_0 + T_1 + T_2 + T_3$ are diagonalizable.

Proof of Theorem 4. T are diagonalizable if, and only if, there exists $L = L_0 + L_1P_1 + L_2P_2 + L_3P_3$ and $D = D_0 + D_1P_1 + D_2P_2 + D_3P_3$ such that $T = L^{-1}DL$, where D is diagonal and L is invertible.

The equation $T = L^{-1}DL$ is equivalent to the following:

$$\begin{cases} T_0 = L_0^{-1}D_0L_0 \\ T_0 + T_1 = (L_0 + L_1)^{-1}(D_0 + D_1)(L_0 + L_1) \\ T_0 + T_1 + T_2 = (L_0 + L_1 + L_2)^{-1}(D_0 + D_1 + D_2)(L_0 + L_1 + L_2) \\ T_0 + T_1 + T_2 + T_3 = (L_0 + L_1 + L_2 + L_3)^{-1}(D_0 + D_1 + D_2 + D_3)(L_0 + L_1 + L_2 + L_3) \end{cases}$$

Which implies the proof. \square

Example 4. Consider the following matrix:

$$\begin{aligned} T &= \begin{pmatrix} 2 + P_1 + P_2 + 2P_3 & P_1 + P_2 \\ 3 + P_3 & -1 + 3P_2 + P_3 \end{pmatrix} \\ &= \begin{pmatrix} 2 & 0 \\ 3 & -1 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix} P_1 + \begin{pmatrix} 1 & 1 \\ 0 & 3 \end{pmatrix} P_2 + \begin{pmatrix} 2 & 0 \\ 1 & 1 \end{pmatrix} P_3 \\ &= T_0 + T_1P_1 + T_2P_2 + T_3P_3 \end{aligned}$$

$$T_0 = \begin{pmatrix} 2 & 0 \\ 3 & -1 \end{pmatrix} = L_0 D_0 L_0^{-1} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}^{-1}$$

$$\begin{aligned} T_0 + T_1 &= \begin{pmatrix} 3 & 1 \\ 3 & -1 \end{pmatrix} = M_1 K_1 M_1^{-1} \\ &= \begin{pmatrix} 1 & 1 \\ -2 + \sqrt{7} & -2 - \sqrt{7} \end{pmatrix} \begin{pmatrix} 1 + \sqrt{7} & 0 \\ 0 & 1 - \sqrt{7} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -2 + \sqrt{7} & -2 - \sqrt{7} \end{pmatrix}^{-1} \end{aligned}$$

$$\begin{aligned}
T_0 + T_1 + T_2 &= \begin{pmatrix} 4 & 2 \\ 3 & 2 \end{pmatrix} = M_2 K_2 M_2^{-1} \\
&= \begin{pmatrix} 1 & 1 \\ \frac{-1+\sqrt{7}}{2} & \frac{-1-\sqrt{7}}{2} \end{pmatrix} \begin{pmatrix} 3+\sqrt{7} & 0 \\ 0 & 3-\sqrt{7} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \frac{-1+\sqrt{7}}{2} & \frac{-1-\sqrt{7}}{2} \end{pmatrix}^{-1} \\
T_0 + T_1 + T_2 + T_3 &= \begin{pmatrix} 6 & 2 \\ 4 & 3 \end{pmatrix} = M_3 K_3 M_3^{-1} \\
&= \begin{pmatrix} 1 & 1 \\ \frac{-3+\sqrt{41}}{4} & \frac{-3-\sqrt{41}}{4} \end{pmatrix} \begin{pmatrix} \frac{9+\sqrt{41}}{4} & 0 \\ 0 & \frac{9-\sqrt{41}}{4} \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \frac{-3+\sqrt{41}}{4} & \frac{-3-\sqrt{41}}{4} \end{pmatrix}^{-1}
\end{aligned}$$

So that $L_1 = M_1 - L_0, L_2 = M_2 - M_1, L_3 = M_3 - M_2, D_1 = K_1 - D_0, D_2 = K_2 - K_1, D_3 = K_3 - K_2$.

3.4. Symbolic 4-Plithogenic Matrices

Definition 5. A square symbolic 4-plithogenic matrix is a matrix with symbolic 4-plithogenic entries.

Example 5.

$$A = \begin{pmatrix} 4 + P_1 + P_2 - 2P_3 + 4P_4 & 1 + P_3 + 2P_4 & 3 - P_1 + 5P_3 + 4P_4 \\ 5 - P_1 + P_2 + 3P_4 & 2 - P_1 + 11P_4 & 7P_1 + 4P_2 - P_3 + 6P_4 \\ 6P_2 + 4P_3 + 2P_4 & 5 + 11P_2 + P_3 + 11P_4 & P_1 + P_3 + 3P_4 \end{pmatrix}$$

is a symbolic 4-plithogenic 3×3 real matrix.

Remark 4. Any symbolic 4-plithogenic matrix can be written as follows: $A = A_0 + A_1 P_1 + A_2 P_2 + A_3 P_3 + A_4 P_4$.

For example, the matrix defined in the previous example can be written as follows:

$$A = \begin{pmatrix} 4 & 1 & 3 \\ 5 & 2 & 0 \\ 0 & 5 & 0 \end{pmatrix} + \begin{pmatrix} 1 & 0 & -1 \\ -1 & -1 & 7 \\ 0 & 0 & 1 \end{pmatrix} P_1 + \begin{pmatrix} 1 & 0 & 0 \\ 1 & 0 & 4 \\ 6 & 11 & 0 \end{pmatrix} P_2 + \begin{pmatrix} -2 & 1 & 5 \\ 0 & 0 & -1 \\ 4 & 1 & 1 \end{pmatrix} P_3 + \begin{pmatrix} 4 & 2 & 4 \\ 3 & 11 & 6 \\ 2 & 11 & 3 \end{pmatrix} P_4$$

Definition 6. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 + S_4 P_4, T = T_0 + T_1 P_1 + T_2 P_2 + T_3 P_3 + T_4 P_4$ be two n -square symbolic 4-plithogenic matrices, then the following is true:

$$S + T = (S_0 + T_0) + (S_1 + T_1)P_1 + (S_2 + T_2)P_2 + (S_3 + T_3)P_3 + (S_4 + T_4)P_4$$

$$S \times T = S_0 T_0 + (S_0 T_1 + S_1 T_0 + S_1 T_1)P_1 + (S_0 T_2 + S_2 T_1 + S_2 T_2 + S_2 T_0 + S_1 T_2)P_2 + (S_0 T_3 + S_1 T_3 + S_2 T_3 + S_3 T_3 + S_3 T_0 + S_3 T_1 + S_3 T_2)P_3 + (S_0 T_4 + S_1 T_4 + S_2 T_4 + S_3 T_4 + S_4 T_0 + S_4 T_1 + S_4 T_2 + S_4 T_3 + S_4 T_4)P_4$$

We denote the set of all symbolic 4-plithogenic n -square matrices using $(4 - SP_M)$.

It is clear that $(4 - SP_M, +, .)$. Is a ring.

Theorem 5. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 + S_4 P_4 \in 4 - SP_M$, then:

1. S is invertible if, and only if, $S_0, S_0 + S_1, S_0 + S_1 + S_2, S_0 + S_1 + S_2 + S_3, S_0 + S_1 + S_2 + S_3 + S_4$ are invertible.
2. $S^{-1} = S_0^{-1} + [(S_0 + S_1)^{-1} - S_0^{-1}]P_1 + [(S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}]P_2 + [(S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}]P_3 + [(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - (S_0 + S_1 + S_2 + S_3)^{-1}]P_4$
3. For $n \in \mathbb{N}$

4. $S^n = S_0^n + [(S_0 + S_1)^n - S_0^n]P_1 + [(S_0 + S_1 + S_2)^n - (S_0 + S_1)^n]P_2 + [(S_0 + S_1 + S_2 + S_3)^n - (S_0 + S_1 + S_2)^n]P_3 + [(S_0 + S_1 + S_2 + S_3 + S_4)^n - (S_0 + S_1 + S_2 + S_3)^n]P_4.$
5. $e^S = e^{S_0} + [e^{S_0+S_1} - e^{S_0}]P_1 + [e^{S_0+S_1+S_2} - e^{S_0+S_1}]P_2 + [e^{S_0+S_1+S_2+S_3} - e^{S_0+S_1+S_2}]P_3 + [e^{S_0+S_1+S_2+S_3+S_4} - e^{S_0+S_1+S_2+S_3}]P_4$

Proof of Theorem 5. (1,2). Assume that $S_0, S_0 + S_1, S_0 + S_1 + S_2, S_0 + S_1 + S_2 + S_3, S_0 + S_1 + S_2 + S_3 + S_4$ are invertible. We have the following:

$$K = S_0^{-1} + [(S_0 + S_1)^{-1} - S_0^{-1}]P_1 + [(S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}]P_2 + [(S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}]P_3 + [(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - (S_0 + S_1 + S_2 + S_3)^{-1}]P_4,$$

Let us compute the following:

$$\begin{aligned} SK &= S_0 S_0^{-1} + \left[S_0(S_0 + S_1)^{-1} - S_0 S_0^{-1} + S_1 S_0^{-1} + S_1(S_0 + S_1)^{-1} - S_1 S_0^{-1} \right] P_1 \\ &\quad + \left[S_0(S_0 + S_1 + S_2)^{-1} - S_0(S_0 + S_1)^{-1} + S_1(S_0 + S_1 + S_2)^{-1} - S_1(S_0 + S_1)^{-1} \right. \\ &\quad \left. + S_2(S_0 + S_1 + S_2)^{-1} - S_2(S_0 + S_1)^{-1} + S_2 S_0^{-1} + S_2(S_0 + S_1)^{-1} - S_2 S_0^{-1} \right] P_2 + \\ [S_0(S_0 + S_1 + S_2) &+ S_3]^{-1} - S_0(S_0 + S_1 + S_2)^{-1} + S_1(S_0 + S_1 + S_2 + S_3)^{-1} - S_1(S_0 + S_1 + S_2)^{-1} \\ &+ S_2(S_0 + S_1 + S_2 + S_3)^{-1} - S_2(S_0 + S_1 + S_2)^{-1} + S_3(S_0 + S_1 + S_2 + S_3)^{-1} \\ &- S_3(S_0 + S_1 + S_2)^{-1} + S_3 S_0^{-1} + S_3(S_0 + S_1)^{-1} - S_3 S_0^{-1} + S_3(S_0 + S_1 + S_2 + S_3)^{-1} \\ &- S_3(S_0 + S_1)^{-1}] P_3 + \\ [S_0(S_0 + S_1 + S_2) &+ S_3 + S_4]^{-1} - S_0(S_0 + S_1 + S_2 + S_3)^{-1} + S_1(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} \\ &- S_1(S_0 + S_1 + S_2 + S_3)^{-1} + S_2(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - S_2(S_0 + S_1 + S_2 + S_3)^{-1} \\ &+ S_3(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - S_3(S_0 + S_1 + S_2 + S_3)^{-1} + S_4 S_0^{-1} + S_4(S_0 + S_1)^{-1} \\ &- S_4 S_0^{-1} + S_4(S_0 + S_1 + S_2)^{-1} - S_4(S_0 + S_1)^{-1} + S_4(S_0 + S_1 + S_2 + S_3)^{-1} \\ &- S_4(S_0 + S_1 + S_2)^{-1} + S_4(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - S_4(S_0 + S_1 + S_2 + S_3)^{-1}] P_4 = \\ U_{n \times n} &+ [O_{n \times n}] P_1 + [O_{n \times n}] P_2 + [O_{n \times n}] P_3 + [O_{n \times n}] P_4 = U_{n \times n} \end{aligned}$$

Thus, S is invertible and $S^{-1} = K$.

For the converse, if we suppose that S is invertible, then there exists $K = K_0 + K_1 P_1 + K_2 P_2 + K_3 P_3 + K_4 P_4 \in 4 - SP_M$ such that $S \times K = U_{n \times n}$. This is equivalent to the following:

$$\left\{ \begin{array}{l} S_0 K_0 = U_{n \times n} \quad (9) \\ S_0 K_1 + S_1 K_0 + S_1 K_1 = O_{n \times n} \quad (10) \\ S_0 K_2 + S_1 K_2 + S_2 K_2 + S_2 K_1 + S_2 K_0 = O_{n \times n} \quad (11) \\ S_0 K_3 + S_1 K_3 + S_2 K_3 + S_3 K_3 + S_3 K_0 + S_3 K_1 + S_3 K_2 = O_{n \times n} \quad (12) \\ S_0 K_4 + S_1 K_4 + S_2 K_4 + S_3 K_4 + S_4 K_0 + S_4 K_1 + S_4 K_2 + S_4 K_3 + S_4 K_4 = O_{n \times n} \quad (13) \end{array} \right.$$

Equation (9) implies that S_0 is invertible and $S_0^{-1} = K_0$.

By adding (9) to (10), we obtain $(S_0 + S_1)(K_0 + K_1) = U_{n \times n}$, so that $S_0 + S_1$ is invertible and $(S_0 + S_1)^{-1} = K_0 + K_1$, hence $K_1 = (S_0 + S_1)^{-1} - S_0^{-1}$.

By adding (9) to (10) to (11), we obtain $(S_0 + S_1 + S_2)(K_0 + K_1 + K_2) = U_{n \times n}$, so that $S_0 + S_1 + S_2$ is invertible and $(S_0 + S_1 + S_2)^{-1} = K_0 + K_1 + K_2$, hence $K_2 = (S_0 + S_1 + S_2)^{-1} - (S_0 + S_1)^{-1}$.

By adding (9) to (10) to (11) to (12), we obtain $(S_0 + S_1 + S_2 + S_3)(K_0 + K_1 + K_2 + K_3) = U_{n \times n}$, so that $S_0 + S_1 + S_2 + S_3$ is invertible and $(S_0 + S_1 + S_2 + S_3)^{-1} = K_0 + K_1 + K_2 + K_3$, hence $K_3 = (S_0 + S_1 + S_2 + S_3)^{-1} - (S_0 + S_1 + S_2)^{-1}$.

By adding all equations, we obtain the following:

$$(S_0 + S_1 + S_2 + S_3 + S_4)(K_0 + K_1 + K_2 + K_3 + K_4) = U_{n \times n}$$

so that $S_0 + S_1 + S_2 + S_3 + S_4$ is invertible and $(S_0 + S_1 + S_2 + S_3 + S_4)^{-1} = K_0 + K_1 + K_2 + K_3 + K_4$; hence $K_4 = (S_0 + S_1 + S_2 + S_3 + S_4)^{-1} - (S_0 + S_1 + S_2 + S_3)^{-1}$.

(3). For $n = 1$, it holds directly. We suppose that it is true. For $n = k$, we must prove it for $k + 1$.

$$\begin{aligned} S^{k+1} = SS^k = S_0 S_0^k &+ \left[S_0(S_0 + S_1)^k - S_0 S_0^k + S_1 S_0^k + S_1(S_0 + S_1)^k - S_1 S_0^k \right] P_1 \\ &+ \left[S_0(S_0 + S_1 + S_2)^k - S_0(S_0 + S_1)^k + S_1(S_0 + S_1 + S_2)^k - S_1(S_0 + S_1)^k + S_2(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. - S_2(S_0 + S_1)^k + S_2 S_0^k + S_2(S_0 + S_1)^k - S_2 S_0^k \right] P_2 \\ &+ \left[S_0(S_0 + S_1 + S_2 + S_3)^k - S_0(S_0 + S_1 + S_2)^k + S_1(S_0 + S_1 + S_2 + S_3)^k - S_1(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. + S_2(S_0 + S_1 + S_2 + S_3)^k - S_2(S_0 + S_1 + S_2)^k + S_3(S_0 + S_1 + S_2 + S_3)^k - S_3(S_0 + S_1 + S_2)^k \right. \\ &\quad \left. + S_3 S_0^k + S_3(S_0 + S_1)^k - S_3 S_0^k + S_3(S_0 + S_1 + S_2 + S_3)^k - S_3(S_0 + S_1)^k \right] P_3 + \\ &\quad \left[S_0(S_0 + S_1 + S_2 + S_3 + S_4)^k - S_0(S_0 + S_1 + S_2 + S_3)^k + S_1(S_0 + S_1 + S_2 + S_3 + S_4)^k - \right. \\ &\quad \left. S_1(S_0 + S_1 + S_2 + S_3)^k + S_2(S_0 + S_1 + S_2 + S_3 + S_4)^k - S_2(S_0 + S_1 + S_2 + S_3)^k + S_3(S_0 + S_1 + \right. \\ &\quad \left. S_2 + S_3 + S_4)^k - S_3(S_0 + S_1 + S_2 + S_3)^k + S_4 S_0^k + S_4(S_0 + S_1)^k - S_4 S_0^k + S_4(S_0 + S_1 + S_2)^k - \right. \\ &\quad \left. S_4(S_0 + S_1)^k + S_4(S_0 + S_1 + S_2 + S_3)^k - S_4(S_0 + S_1 + S_2)^k + S_4(S_0 + S_1 + S_2 + S_3 + S_4)^k - \right. \\ &\quad \left. S_4(S_0 + S_1 + S_2 + S_3)^k \right] P_4 = S_0^{k+1} + \left[(S_0 + S_1)^{k+1} - S_0 S_0^{k+1} \right] P_1 + \left[(S_0 + S_1 + S_2)^{k+1} - (S_0 + S_1)^{k+1} \right] P_2 + \left[(S_0 + S_1 + S_2 + S_3)^{k+1} - (S_0 + S_1 + S_2)^{k+1} \right] P_3 + \left[(S_0 + S_1 + S_2 + S_3 + S_4)^n - (S_0 + S_1 + S_2 + S_3)^n \right] P_4. \end{aligned}$$

This is so that the proof holds by induction.

$$\begin{aligned} (4). e^S = \sum_{n=0}^{\infty} \frac{S^n}{n!} &= \sum_{n=0}^{\infty} \frac{S_0^n}{n!} + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1)^n}{n!} - \sum_{n=0}^{\infty} \frac{S_0^n}{n!} \right] P_1 + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2)^n}{n!} - \right. \\ &\quad \left. \sum_{n=0}^{\infty} \frac{(S_0 + S_1)^n}{n!} \right] P_2 + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2 + S_3)^n}{n!} - \sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2)^n}{n!} \right] P_3 + \left[\sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2 + S_3 + S_4)^n}{n!} - \right. \\ &\quad \left. \sum_{n=0}^{\infty} \frac{(S_0 + S_1 + S_2 + S_3)^n}{n!} \right] P_4 = e^{S_0} + [e^{S_0 + S_1} - e^{S_0}] P_1 + [e^{S_0 + S_1 + S_2} - e^{S_0 + S_1}] P_2 + [e^{S_0 + S_1 + S_2 + S_3} - \\ &\quad e^{S_0 + S_1 + S_2}] P_3 + [e^{S_0 + S_1 + S_2 + S_3 + S_4} - e^{S_0 + S_1 + S_2 + S_3}] P_4. \quad \square \end{aligned}$$

Definition 7. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 + S_4 P_4 \in 4 - SP_M$, then we define the following:

$$\det(S) = \det(S_0) + [\det(S_0 + S_1) - \det(S_0)] P_1 + [\det(S_0 + S_1 + S_2) - \det(S_0 + S_1)] P_2 + [\det(S_0 + S_1 + S_2 + S_3) - \det(S_0 + S_1 + S_2)] P_3 + [\det(S_0 + S_1 + S_2 + S_3 + S_4) - \det(S_0 + S_1 + S_2 + S_3)] P_4.$$

Theorem 6. S is invertible if, and only if, $\det(S)$ is invertible in $4 - SP_R$.

The proof holds via a similar discussion of the 3-plithogenic case.

3.5. Symbolic 4-Plithogenic Eigen Values/Vectors

Theorem 7. Let $S = S_0 + S_1 P_1 + S_2 P_2 + S_3 P_3 + S_4 P_4 \in 4 - SP_M$, then $A = a_0 + a_1 P_1 + a_2 P_2 + a_3 P_3 + a_4 P_4$ is an eigen value of S if, and only if, a_0 eigen value of S_0 , $a_0 + a_1$ eigen value of $S_0 + S_1$, $a_0 + a_1 + a_2$ eigen value of $S_0 + S_1 + S_2$, $a_0 + a_1 + a_2 + a_3$ eigen value of $S_0 + S_1 + S_2 + S_3$, $a_0 + a_1 + a_2 + a_3 + a_4$ eigen value of $S_0 + S_1 + S_2 + S_3 + S_4$.

In addition, $X = X_0 + X_1 P_1 + X_2 P_2 + X_3 P_3 + X_4 P_4$ is the corresponding eigen vector of A if, and only if, the X_0 eigen vector of a_0 , the $X_0 + X_1$ eigen vector of $a_0 + a_1$, the $X_0 + X_1 + X_2$ eigen vector of $a_0 + a_1 + a_2$, the $X_0 + X_1 + X_2 + X_3$ eigen vector of $a_0 + a_1 + a_2 + a_3$, the $X_0 + X_1 + X_2 + X_3 + X_4$ eigen vector of $a_0 + a_1 + a_2 + a_3 + a_4$.

Proof of Theorem 7. From the equation $AS = AX$, we obtain the following:

$$\left\{ \begin{array}{l} a_0 S_0 = a_0 X_0 \quad (14) \\ a_0 S_1 + a_1 S_0 + a_1 S_1 = a_0 X_1 + a_1 X_0 + a_1 X_1 \quad (15) \\ a_0 S_2 + a_1 S_2 + a_2 S_1 + a_2 S_0 = a_0 X_2 + a_1 X_2 + a_2 X_2 + a_2 X_1 + a_2 X_0 \quad (16) \\ a_0 S_3 + a_1 S_3 + a_2 S_3 + a_3 S_0 + a_3 S_1 + a_3 S_2 = a_0 X_3 + a_1 X_3 + a_2 X_3 + a_3 X_3 + a_3 X_0 + a_3 X_1 + a_3 X_2 \quad (17) \\ a_0 S_4 + a_1 S_4 + a_2 S_4 + a_3 S_4 + a_4 S_0 + a_4 S_1 + a_4 S_2 + a_4 S_3 + a_4 S_4 = a_0 X_4 + a_1 X_4 + a_2 X_4 + a_3 X_4 + a_4 X_0 + a_4 X_1 + a_4 X_2 + a_4 X_3 + a_4 X_4 \quad (18) \end{array} \right.$$

Equation (14) implies that a_0 is the eigen value of S_0 , with X_0 as the eigen vector.

By adding (14) to (15), we obtain $(a_0 + a_1)(S_0 + S_1) = (a_0 + a_1)(X_0 + X_1)$, which means that $a_0 + a_1$ is the eigen value of $S_0 + S_1$, with $X_0 + X_1$ as the eigen vector.

By adding (14) to (15) to (16), we obtain $(a_0 + a_1 + a_2)(S_0 + S_1 + S_2) = (a_0 + a_1 + a_2)(X_0 + X_1 + X_2)$, which means that $a_0 + a_1 + a_2$ is the eigen value of $S_0 + S_1 + S_2$, with $X_0 + X_1 + X_2$ as the eigen vector.

By adding (14) to (15) to (16) to (17), we obtain $(a_0 + a_1 + a_2 + a_3)(S_0 + S_1 + S_2 + S_3) = (a_0 + a_1 + a_2 + a_3)(X_0 + X_1 + X_2 + X_3)$, which means that $a_0 + a_1 + a_2 + a_3$ is the eigen value of $S_0 + S_1 + S_2 + S_3$, with $X_0 + X_1 + X_2 + X_3$ as the eigen vector.

By adding all equations, we obtain the following: $(a_0 + a_1 + a_2 + a_3 + a_4)(S_0 + S_1 + S_2 + S_3 + S_4) = (a_0 + a_1 + a_2 + a_3 + a_4)(X_0 + X_1 + X_2 + X_3 + X_4)$, which means that $a_0 + a_1 + a_2 + a_3 + a_4$ is the eigen value of $S_0 + S_1 + S_2 + S_3 + S_4$, with $X_0 + X_1 + X_2 + X_3 + X_4$ as the eigen vector.

This is so that the proof is complete. \square

Example 6. Consider the following matrix:

$$S = \begin{pmatrix} 3 + 2P_1 - P_3 - P_4 & 4 + P_2 - 3P_3 \\ 2P_1 + P_2 + P_3 - P_4 & 5 + P_1 + P_4 \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 0 & 5 \end{pmatrix} + \begin{pmatrix} 2 & 0 \\ 1 & 1 \end{pmatrix}P_1 + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}P_2 + \begin{pmatrix} -1 & -3 \\ 1 & 0 \end{pmatrix}P_3 + \begin{pmatrix} -1 & 0 \\ -1 & 1 \end{pmatrix}P_4 = A_0 + A_1P_1 + A_2P_2 + A_3P_3 + A_4P_4$$

The eigen values of $A_0 = \begin{pmatrix} 3 & 4 \\ 0 & 5 \end{pmatrix}$ are $\{3, 5\}$.

The eigen values of $A_0 + A_1 = \begin{pmatrix} 2 & 0 \\ 2 & 1 \end{pmatrix}$ are $\left\{ \frac{11+\sqrt{33}}{2}, \frac{11-\sqrt{33}}{2} \right\}$.

The eigen values of $A_0 + A_1 + A_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ are $\left\{ \frac{11+\sqrt{61}}{2}, \frac{11-\sqrt{61}}{2} \right\}$.

The eigen values of $A_0 + A_1 + A_2 + A_3 = \begin{pmatrix} -1 & -3 \\ 1 & 0 \end{pmatrix}$ are $\{8, 2\}$.

The eigen values of $A_0 + A_1 + A_2 + A_3 + A_4 = \begin{pmatrix} -2 & -3 \\ 0 & 1 \end{pmatrix}$ are $\{-2, 1\}$.

The eigen values of the symbolic 4-plithogenic matrix A are as follows:

$$U_1 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right)P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right)P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right)P_3 + (-10)P_4$$

$$U_2 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right)P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right)P_2 + \left(2 - \frac{11+\sqrt{61}}{2} \right)P_3 + (-4)P_4$$

$$U_3 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right)P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right)P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right)P_3 + (-7)P_4$$

$$U_4 = 3 + \left(\frac{11+\sqrt{33}}{2} - 3 \right)P_1 + \left(\frac{11-\sqrt{61}}{2} - \frac{11+\sqrt{33}}{2} \right)P_2 + \left(2 - \frac{11+\sqrt{61}}{2} \right)P_3 + (-1)P_4$$

$$U_5 = 3 + \left(\frac{11-\sqrt{33}}{2} - 3 \right)P_1 + \left(\frac{11+\sqrt{61}}{2} - \frac{11-\sqrt{33}}{2} \right)P_2 + \left(8 - \frac{11+\sqrt{61}}{2} \right)P_3 + (-10)P_4$$

$$\begin{aligned}
U_6 &= 3 + \left(\frac{11 - \sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-4) P_4 \\
U_7 &= 3 + \left(\frac{11 - \sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-7) P_4 \\
U_8 &= 3 + \left(\frac{11 - \sqrt{33}}{2} - 3 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-1) P_4 \\
U_9 &= 5 + \left(\frac{11 + \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 + \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 + \sqrt{61}}{2} \right) P_3 + (-10) P_4 \\
U_{10} &= 5 + \left(\frac{11 + \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 + \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 + \sqrt{61}}{2} \right) P_3 + (-4) P_4 \\
U_{11} &= 5 + \left(\frac{11 + \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 + \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-7) P_4 \\
U_{12} &= 5 + \left(\frac{11 + \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 + \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-1) P_4 \\
U_{13} &= 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 + \sqrt{61}}{2} \right) P_3 + (-10) P_4 \\
U_{14} &= 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 + \sqrt{61}}{2} \right) P_3 + (-4) P_4 \\
U_{15} &= 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 - \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(8 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-7) P_4 \\
U_{16} &= 5 + \left(\frac{11 - \sqrt{33}}{2} - 5 \right) P_1 + \left(\frac{11 + \sqrt{61}}{2} - \frac{11 - \sqrt{33}}{2} \right) P_2 + \left(2 - \frac{11 - \sqrt{61}}{2} \right) P_3 + (-1) P_4.
\end{aligned}$$

We continue using the same process to obtain all other eigen values.

Remark 5. If A has n eigen values for $0 \leq i \leq s$, then A has n^5 eigen values.

3.6. Symbolic 4-Plithogenic Diagonalization

Theorem 8. Let $T = T_0 + T_1 P_1 + T_2 P_2 + T_3 P_3 + T_4 P_4$ be an n -square symbolic 4-plithogenic matrix, then T is diagonalizable if, and only if, $T_0, T_0 + T_1, T_0 + T_1 + T_2, T_0 + T_1 + T_2 + T_3, T_0 + T_1 + T_2 + T_3 + T_4$ are diagonalizable.

Proof of Theorem 8. T are diagonalizable if, and only if, there exists $L = L_0 + L_1 P_1 + L_2 P_2 + L_3 P_3 + L_4 P_4$ and $D = D_0 + D_1 P_1 + D_2 P_2 + D_3 P_3 + D_4 P_4$, such that $T = L^{-1}DL$, where D is diagonal and L is invertible.

The equation $T = L^{-1}DL$ is equivalent to the following:

$$\left\{
\begin{aligned}
T_0 &= L_0^{-1} D_0 L_0 \\
T_0 + T_1 &= (L_0 + L_1)^{-1} (D_0 + D_1) (L_0 + L_1) \\
T_0 + T_1 + T_2 &= (L_0 + L_1 + L_2)^{-1} (D_0 + D_1 + D_2) (L_0 + L_1 + L_2) \\
T_0 + T_1 + T_2 + T_3 &= (L_0 + L_1 + L_2 + L_3)^{-1} (D_0 + D_1 + D_2 + D_3) (L_0 + L_1 + L_2 + L_3) \\
T_0 + T_1 + T_2 + T_3 + T_4 &= (L_0 + L_1 + L_2 + L_3 + L_4)^{-1} (D_0 + D_1 + D_2 + D_3 + D_4) (L_0 + L_1 + L_2 + L_3 + L_4)
\end{aligned}
\right.$$

This implies the proof. \square

4. Conclusions

In this scientific work, we have studied the symbolic 3-plithogenic square real matrices and the symbolic 4-plithogenic square real matrices from many different perspectives, and we have presented many algorithms and theorems to describe the invertibility, the eigen values, vectors, and diagonalization of these matrices by transforming them into classical matrices. In the future, we aim to study the symbolic 5-plithogenic and 6-plithogenic real square matrices in a similar approach to that used for the symbolic 3-plithogenic matrices and symbolic 4-plithogenic matrices.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The author declares no conflict of interest.

References

1. Smarandache, F. Introduction to the Symbolic Plithogenic Algebraic Structures (revisited). *Neutrosophic Sets Syst.* **2023**, *53*, 39.
2. Smarandache, F. *Plithogenic Algebraic Structures*. Chapter in “*Nidus idearum Scilogs, V: Joining the Dots*”, 3rd ed.; Pons Publishing: Brussels, Belgium, 2019; pp. 123–125.
3. Smarandache, F. Plithogenic Set, an Extension of Crisp, Fuzzy, Intuitionistic Fuzzy, and Neutrosophic Sets—Revisited. *Neutrosophic Sets Syst.* **2018**, *21*, 153–166.
4. Taffach, N.M.; Hatip, A. A Review on Symbolic 2-Plithogenic Algebraic Structures. *Galoitica J. Math. Struct. Appl.* **2023**, *5*, 8–16. [[CrossRef](#)]
5. Taffach, N.M.; Hatip, A. A Brief Review on The Symbolic 2-Plithogenic Number Theory and Algebraic Equations. *Galoitica J. Math. Struct. Appl.* **2023**, *5*, 36–44. [[CrossRef](#)]
6. Merkepcı, H.; Abobala, M. On the Symbolic 2-Plithogenic Rings. *Int. J. Neutrosophic Sci.* **2023**, *20*, 115–122. [[CrossRef](#)]
7. Taffach, N. An Introduction to Symbolic 2-Plithogenic Vector Spaces Generated from the Fusion of Symbolic Plithogenic Sets and Vector Spaces. *Neutrosophic Sets Syst.* **2023**, *54*, 45–57.
8. Taffach, N.; Othman, K.B. An Introduction to Symbolic 2-Plithogenic Modules Over Symbolic 2-Plithogenic Rings. *Neutrosophic Sets Syst.* **2023**, *54*, 33–44.
9. Ali, R.; Hasan, Z. An Introduction to the Symbolic 3-Plithogenic Modules. *Galoitica J. Math. Struct. Appl.* **2023**, *6*, 13–17. [[CrossRef](#)]
10. Ali, R.; Hasan, Z. An Introduction to the Symbolic 3-Plithogenic Vector Spaces. *Galoitica J. Math. Struct. Appl.* **2023**, *6*, 8–12. [[CrossRef](#)]
11. Rawashdeh, A. An Introduction to the Symbolic 3-plithogenic Number Theory. *Neoma J. Math. Comput. Sci.* **2023**. [[CrossRef](#)]
12. Othman, B.K. On Some Algorithms for Solving Symbolic 3-Plithogenic Equations. *Neoma J. Math. Comput. Sci.* **2023**. [[CrossRef](#)]
13. Albasheer, O.; Hajjari, A.; Dalla, R. On The Symbolic 3-Plithogenic Rings and Their Algebraic Properties. *Neutrosophic Sets Syst.* **2023**, *54*, 57–67.
14. Alfahal, A.; Alhasan, Y.; Abdulfatah, R.; Mehmood, A.; Kadhim, M. On Symbolic 2-Plithogenic Real Matrices and Their Algebraic Properties. *Int. J. Neutrosophic Sci.* **2023**, *21*. [[CrossRef](#)]
15. Abobala, M.; Hatip, A.; Olgun, N.; Broumi, S.; Salama, A.A.; Khaled, E.H. The algebraic creativity in the Neutrosophic Square Matrices. *Neutrosophic Sets Syst.* **2021**, *40*, 1–11.
16. Abobala, M. On Refined Neutrosophic Matrices and Their Applications in Refined Neutrosophic Algebraic Equations. *J. Math.* **2021**, *2021*, 5531093. [[CrossRef](#)]
17. Olgun, N.; Hatip, A.; Bal, M.; Abobala, M. A Novel Approach To Necessary and Sufficient Conditions For The Diagonalization of Refined Neutrosophic Matrices. *Int. J. Neutrosophic Sci.* **2021**, *16*, 72–79. [[CrossRef](#)]
18. Merkepcı, H.; Abobala, M. The Application of AH-isometry in the Study of Neutrosophic Conic Sections. *Galoitica J. Math. Struct. Appl.* **2022**, *2*, 18–22.
19. Abobala, M.; Zeina, M.B. A Study of Neutrosophic Real Analysis by Using One Dimensional Geometric AH-Isometry. *Galoitica J. Math. Struct. Appl.* **2023**, *3*, 18–24. [[CrossRef](#)]
20. Von Shtawzen, O. Conjectures for Invertible Diophantine Equations of 3-Cyclic and 4-Cyclic Refined Integers. *J. Neutrosophic Fuzzy Syst.* **2022**, *3*, 32–36. [[CrossRef](#)]
21. Abobala, M.; Hatip, A.; Bal, M. A Review on Recent Advantages in Algebraic Theory of Neutrosophic Matrices. *Int. J. Neutrosophic Sci.* **2021**, *17*, 68–86. [[CrossRef](#)]
22. Abobala, M.; Ziena, B.M.; Doeves, R.; Hussein, Z. The Representation of Refined Neutrosophic Matrices by Refined Neutrosophic Linear Functions. *Int. J. Neutrosophic Sci.* **2022**, *19*, 342–349. [[CrossRef](#)]

23. Ahmad, K.; Thjeel, N.; AbdulZahra, M.; Jaleel, R. On the Classification of n-Refined Neutrosophic Rings and Its Applications in Matrix Computing Algorithms and Linear Systems. *Int. J. Neutrosophic Sci.* **2022**, *18*, 8–15.
24. Abobala, M. On the Representation of Neutrosophic Matrices by Neutrosophic Linear Transformations. *J. Math.* **2021**, *2021*, 5591576. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.