

Article

Some Korovkin-Type Approximation Theorems Associated with a Certain Deferred Weighted Statistical Riemann-Integrable Sequence of Functions

Hari Mohan Srivastava ^{1,2,3,4,*} , Bidu Bhusan Jena ⁵  and Susanta Kumar Paikray ⁵ ¹ Department of Mathematics and Statistics, University of Victoria, Victoria, BC V8W 3R4, Canada² Department of Medical Research, China Medical University Hospital, China Medical University, Taichung 40402, Taiwan³ Department of Mathematics and Informatics, Azerbaijan University, 71 Jeyhun Hajibeyli Street, Baku AZ1007, Azerbaijan⁴ Section of Mathematics, International Telematic University Uninettuno, I-00186 Rome, Italy⁵ Department of Mathematics, Veer Surendra Sai University of Technology, Burla 768018, India; bidumath.05@gmail.com (B.B.J.); skpaikray_math@vssut.ac.in (S.K.P.)

* Correspondence: harimsri@math.uvic.ca

Abstract: Here, in this article, we introduce and systematically investigate the ideas of deferred weighted statistical Riemann integrability and statistical deferred weighted Riemann summability for sequences of functions. We begin by proving an inclusion theorem that establishes a relation between these two potentially useful concepts. We also state and prove two Korovkin-type approximation theorems involving algebraic test functions by using our proposed concepts and methodologies. Furthermore, in order to demonstrate the usefulness of our findings, we consider an illustrative example involving a sequence of positive linear operators in conjunction with the familiar Bernstein polynomials. Finally, in the concluding section, we propose some directions for future research on this topic, which are based upon the core concept of statistical Lebesgue-measurable sequences of functions.

Keywords: Riemann and Lebesgue integrals; statistical Riemann and Lebesgue integral; deferred weighted Riemann summability; Banach space; Bernstein polynomials; positive linear operators; Korovkin-type approximation theorems; Lebesgue-measurable sequences of functions

MSC: 40A05; 40G15; 33C45; 41A36

Citation: Srivastava, H.M.; Jena, B.B.; Paikray, S.K. Some Korovkin-Type Approximation Theorems Associated with a Certain Deferred Weighted Statistical Riemann-Integrable Sequence of Functions. *Axioms* **2022**, *11*, 128. <https://doi.org/10.3390/axioms11030128>

Academic Editor: Yurii Kharkevych

Received: 7 February 2022

Accepted: 11 March 2022

Published: 12 March 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction and Motivation

The relatively more familiar theory of ordinary convergence is one of the most important topics of study of sequence spaces. It has indeed gradually progressed to a very high level of development. Two prominent researchers, Fast [1] and Steinhaus [2], independently created a new idea in the theory of sequence spaces, which is known as *statistical convergence*. This fruitful concept is extremely valuable for studies in various areas of pure and applied mathematical sciences. It is remarkably more powerful than the traditional convergence and has provided a vital area of research in recent years. Furthermore, such a concept is closely related to the study of Real Analysis, Analytic Probability theory and Number theory, and so on. For some recent related developments on this subject, the reader can see, for example, the works in [3–18].

Suppose that $\mathfrak{E} \subseteq \mathbb{N}$. Moreover, let

$$\mathfrak{E}_k = \{\eta : \eta \leq k \text{ and } \eta \in \mathfrak{E}\}.$$

Then, the natural (or asymptotic) density $d(\mathfrak{E})$ of \mathfrak{E} is

$$d(\mathfrak{E}) = \lim_{k \rightarrow \infty} \frac{|\mathfrak{E}_k|}{k} = \tau,$$

where τ is a real and finite number, and $|\mathfrak{E}_k|$ is the cardinality of \mathfrak{E}_k .

A sequence (u_n) is said to be statistically convergent to α if, for each $\epsilon > 0$,

$$\mathfrak{E}_\epsilon = \{\eta : \eta \in \mathbb{N} \text{ and } |u_\eta - \alpha| \geq \epsilon\}$$

has zero natural density (see [1,2]). Thus, for every $\epsilon > 0$,

$$d(\mathfrak{E}_\epsilon) = \lim_{k \rightarrow \infty} \frac{|\mathfrak{E}_\epsilon|}{k} = 0.$$

We write

$$\text{stat} \lim_{k \rightarrow \infty} u_k = \alpha.$$

For a closed and bounded interval $\mathcal{I} := [a, b] \subset \mathbb{R}$, we define the partition of $[a, b]$ as an ordered set that is finite and we denote it as follows:

$$\mathfrak{P} := \{(r_0, r_1, \dots, r_k) : a = r_0 < r_1 < \dots < r_k = b\}.$$

We now divide the interval $[a, b]$ into the following non-overlapping subintervals:

$$\mathcal{I}_1 := [r_0, r_1], \quad \mathcal{I}_2 := [r_1, r_2], \quad \dots, \quad \mathcal{I}_k := [r_{k-1}, r_k].$$

The resulting partition \mathfrak{P} is then given by

$$\mathfrak{P} := \{[r_{i-1}, r_i] : i = 1, 2, 3, \dots, k\}.$$

Next, in order to find the norm of the partition \mathfrak{P} , we have

$$\|\mathfrak{P}\| := \max\{r_1 - r_0, r_2 - r_1, r_3 - r_2, \dots, r_k - r_{k-1}\}.$$

Let γ_i ($i = 1, 2, 3, \dots, k$) be a point that is chosen arbitrarily from each of the subintervals $(\mathcal{I})_{i=1}^k$. We refer to these points as the tags of the subintervals. We also call the subintervals associated with the tags the tagged partitions of \mathcal{I} . We denote it as follows:

$$\mathcal{P} := \{([r_{i-1}, r_i]; \gamma_i) : i = 1, 2, 3, \dots, k\}.$$

Let $[a, b] \subset \mathbb{R}$. Suppose that, for each $i \in \mathbb{N}$, there is a function $h_i : [a, b] \rightarrow \mathbb{R}$. We thus construct the sequence $(h_i)_{i \in \mathbb{N}}$ of functions over the closed interval $[a, b]$.

We now define a subsequence $(h_i)_i^k$ of functions with respect to the Riemann sum associated with a tagged partition \mathcal{P} as follows:

$$\delta(h_i; \mathcal{P}) := \sum_{i=1}^k h(\gamma_i)(r_i - r_{i-1}).$$

We next recall the definition of the Riemann integrability.

A sequence $(h_k)_{k \in \mathbb{N}}$ of functions is Riemann-integrable to h on $[a, b]$ if, for each $\epsilon > 0$, there exists $\sigma_\epsilon > 0$ such that, for any tagged partition \mathcal{P} of $[a, b]$ with $\|\mathcal{P}\| < \sigma_\epsilon$, we have

$$|\delta(h_k; \mathcal{P}) - h| < \epsilon.$$

The definition of statistically Riemann-integrable functions is given as follows.

Definition 1. A sequence $(h_k)_{k \in \mathbb{N}}$ of functions is statistically Riemann-integrable to h on $[a, b]$ if, for every $\epsilon > 0$ and for each $x \in [a, b]$, there exists $\sigma_\epsilon > 0$, and for any tagged partition \mathcal{P} of $[a, b]$ with $\|\mathcal{P}\| < \sigma_\epsilon$, the set

$$\mathfrak{E}_\epsilon = \{\eta : \eta \in \mathbb{N} \text{ and } |\delta(h_\eta; \mathcal{P}) - h| \geq \epsilon\}$$

has zero natural density. That is, for every $\epsilon > 0$,

$$d(\mathfrak{E}_\epsilon) = \lim_{k \rightarrow \infty} \frac{|\mathfrak{E}_\epsilon|}{k} = 0.$$

We write

$$\text{statRie} \lim_{k \rightarrow \infty} \text{ffi}(h_k; \mathcal{P}) = h.$$

By making use of Definition 1, we first establish an inclusion theorem as Theorem 1 below.

Theorem 1. If a sequence of functions (h_k) is Riemann-integrable to h over $[a, b]$, then (h_k) is statistically Riemann-integrable to the same function h over $[a, b]$.

Proof. Given $\epsilon > 0$, there exists $\sigma_\epsilon > 0$. Suppose that \mathcal{P} is any tagged partition of $[a, b]$ such that $\|\mathcal{P}\| < \sigma_\epsilon$. Then

$$|\delta(h_k; \mathcal{P}) - h| < \epsilon.$$

Since, for each $\epsilon > 0$, \mathcal{P} is any tagged partition of $[a, b]$ such that $\|\mathcal{P}\| < \sigma_\epsilon$, so we have

$$\lim_{k \rightarrow \infty} \frac{1}{k} |\{\eta : \eta \in \mathbb{N} \text{ and } |\delta(h_k; \mathcal{P}) - h| \geq \epsilon\}| \leq \lim_{k \rightarrow \infty} |\delta(h_k; \mathcal{P}) - h| < \epsilon.$$

Consequently, by Definition 1, we get

$$\text{statRie} \lim_{k \rightarrow \infty} \delta(h_k; \mathcal{P}) = h,$$

which completes the proof of Theorem 1. \square

Remark 1. In order to demonstrate that the converse of Theorem 1 is not true, we consider Example 1 below.

Example 1. Let $h_k : [0, 1] \rightarrow \mathbb{R}$ be a sequence of functions defined by

$$h_k(x) = \begin{cases} \frac{1}{2} & (x \in \mathbb{Q} \cap [0, 1]; k = j^2, j \in \mathbb{N}) \\ \frac{1}{n} & (\text{otherwise}). \end{cases} \tag{1}$$

It is easily seen that the sequence (h_k) of functions is statistically Riemann-integrable to 0 over the closed interval $[0, 1]$, but it is not Riemann-integrable (in the usual sense) over $[0, 1]$.

Motivated mainly by the above-mentioned investigations and developments, we introduce and study the ideas of deferred weighted statistical Riemann integrability and statistical deferred weighted Riemann summability of sequences of real-valued functions. We first prove an inclusion theorem connecting these two potentially useful concepts. We then state and prove two Korovkin-type approximation theorems with algebraic test functions based on the methodologies and techniques that we have adopted here. Furthermore, we consider an illustrative example involving a positive linear operator in conjunction with the familiar Bernstein polynomials, which shows the effectiveness of our findings. Finally, based upon the core concept of statistical Lebesgue-measurable sequences of functions, we

suggest some possible directions for future research on this topic in the concluding section of our study.

2. Deferred Weighted Statistical Riemann Integrability

Let (ϕ_k) and (φ_k) be sequences of non-negative integers with the regularity conditions given

$$\phi_k < \varphi_k \quad \text{and} \quad \lim_{k \rightarrow \infty} \varphi_k = +\infty.$$

Moreover, let (p_i) be a sequence of non-negative real numbers with

$$P_k = \sum_{i=\phi_k+1}^{\varphi_k} p_i.$$

We then define the deferred weighted summability mean for $\sigma(h_k; \mathcal{P})$ associated with tagged partition \mathcal{P} as follows:

$$\mathcal{W}(\delta(h_k; \mathcal{P})) = \frac{1}{P_k} \sum_{q=\phi_k+1}^{\varphi_k} p_q \delta(h_q; \mathcal{P}). \tag{2}$$

We now present the following definitions for our proposed study.

Definition 2. A sequence $(h_k)_{k \in \mathbb{N}}$ of functions is said to be deferred weighted statistically Riemann-integrable to h on $[a, b]$ if, for all $\epsilon > 0$, there exists $\sigma_\epsilon > 0$, and for any tagged partition \mathcal{P} of $[a, b]$ with $\|\mathcal{P}\| < \sigma_\epsilon$, the following set

$$\{\eta : \eta \leq P_k \quad \text{and} \quad p_\eta |\delta(h_\eta; \mathcal{P}) - h| \geq \epsilon\}$$

has zero natural density. Thus, for every $\epsilon > 0$, we have

$$\lim_{k \rightarrow \infty} \frac{|\{\eta : \eta \leq P_k \quad \text{and} \quad p_\eta |\delta(h_\eta; \mathcal{P}) - h| \geq \epsilon\}|}{P_k} = 0.$$

We write

$$DWR_{\text{stat}} \lim_{k \rightarrow \infty} \delta(h_k; \mathcal{P}) = h.$$

Definition 3. A sequence $(h_k)_{k \in \mathbb{N}}$ of functions is said to statistically deferred weighted Riemann summable to h on $[a, b]$ if, for all $\epsilon > 0 \exists \sigma_\epsilon > 0$ and for any tagged partition \mathcal{P} of $[a, b]$ with $\|\mathcal{P}\| < \sigma_\epsilon$, the set

$$\{\eta : \eta \leq k \quad \text{and} \quad |\mathcal{W}(\delta(h_\eta; \mathcal{P})) - h| \geq \epsilon\}$$

has zero natural density. Thus, for all $\epsilon > 0$, we have

$$\lim_{k \rightarrow \infty} \frac{|\{\eta : \eta \leq k \quad \text{and} \quad |\mathcal{W}(\delta(h_\eta; \mathcal{P})) - h| \geq \epsilon\}|}{k} = 0.$$

We write

$$\text{stat}_{DWR} \lim_{k \rightarrow \infty} \text{ffi}(h_k; \mathcal{P}) = h.$$

An inclusion theorem between the two new potentially useful notions in Definitions 2 and 3 is now given by Theorem 2 below.

Theorem 2. If the sequence $(h_k)_{k \in \mathbb{N}}$ of functions is deferred weighted statistically Riemann-integrable to a function h over $[a, b]$, then it is statistically deferred weighted Riemann summable to the same function h over $[a, b]$, but not conversely.

Proof. Suppose that the sequence $(h_k)_{k \in \mathbb{N}}$ is deferred weighted statistically Riemann-integrable to a function h on $[a, b]$. Then, by Definition 2, we have

$$\lim_{k \rightarrow \infty} \frac{|\{\eta : \eta \leq P_k \text{ and } p_\eta |\delta(h_\eta; \mathcal{P}) - h| \geq \epsilon\}|}{P_k} = 0.$$

Now, if we choose the two sets as follows,

$$\mathcal{O}_\epsilon = \{\eta : \eta \leq P_k \text{ and } p_\eta |\delta(h_\eta; \mathcal{P}) - h| \geq \epsilon\}$$

and

$$\mathcal{O}_\epsilon^c = \{\eta : \eta \leq P_k \text{ and } p_\eta |\delta(h_\eta; \mathcal{P}) - h| < \epsilon\},$$

then we have

$$\begin{aligned} |\mathcal{W}(\delta(h_k; \mathcal{P})) - h| &= \left| \frac{1}{P_k} \sum_{q=\phi_k+1}^{\varphi_k} p_q \delta(h_q; \mathcal{P}) - h \right| \\ &\leq \left| \frac{1}{P_k} \sum_{q=\phi_k+1}^{\varphi_k} p_q [\delta(h_q; \mathcal{P}) - h] \right| + \left| \frac{1}{P_k} \sum_{q=\phi_k+1}^{\varphi_k} p_q h - h \right| \\ &\leq \frac{1}{P_k} \sum_{\substack{q=\phi_k+1 \\ (\eta \in \mathcal{O}_\epsilon)}}^{\varphi_k} p_q |\delta(h_q; \mathcal{P}) - h| + \frac{1}{P_k} \sum_{\substack{q=\phi_k+1 \\ (\eta \in \mathcal{O}_\epsilon^c)}}^{\varphi_k} p_q |\delta(h_q; \mathcal{P}) - h| \\ &\quad + |h| \left| \frac{1}{P_k} \sum_{q=\phi_k+1}^{\varphi_k} p_q - 1 \right| \\ &\leq \frac{1}{P_k} |\mathcal{O}_\epsilon| + \frac{1}{P_k} |\mathcal{O}_\epsilon^c|. \end{aligned}$$

We thus obtain

$$|\mathcal{W}(\delta(h_k; \mathcal{P})) - h| < \epsilon.$$

Hence, clearly, the sequence of functions (h_k) is statistically deferred weighted Riemann-summable to h over $[a, b]$. \square

The following example shows that the converse statement of Theorem 2 is not true.

Example 2. Let $h_k : [0, 1] \rightarrow \mathbb{R}$ be a sequence of functions of the form given by

$$h_k(x) = \begin{cases} 0 & (x \in \mathbb{Q} \cap [0, 1]; k \text{ is even}) \\ 1 & (x \in \mathbb{R} - \mathbb{Q} \cap [0, 1]; k \text{ is odd}), \end{cases} \tag{3}$$

where

$$\phi_k = 2k \quad \varphi_k = 4k \quad \text{and} \quad p_k = 1.$$

The above-specified sequence (h_k) of functions trivially indicates that it is neither Riemann-integrable nor deferred weighted statistically Riemann-integrable. However, as per our proposed mean (2), it is easy to see that

$$\begin{aligned} \mathcal{W}(\delta(h_k; \mathcal{P})) &= \frac{1}{\varphi_k - \phi_k} \sum_{q=\phi_k+1}^{\varphi_k} \delta(h_q; \mathcal{P}) \\ &= \frac{1}{2k} \sum_{m=2k+1}^{4k} \delta(h_q; \mathcal{P}) = \frac{1}{2}. \end{aligned}$$

Thus, clearly, the sequence (h_k) of functions has deferred weighted Riemann sum $\frac{1}{2}$ under the tagged partition \mathcal{P} . Therefore, the sequence (h_k) of functions is statistically deferred weighted Riemann-summable to $\frac{1}{2}$ over $[0, 1]$, but it is not deferred weighted statistically Riemann-integrable over $[0, 1]$.

3. Korovkin-Type Approximation Theorems via the $\mathcal{W}(\delta(h_k; \mathcal{P}))$ -Mean

Many researchers have worked toward extending (or generalizing) the approximation-theoretic aspects of the Korovkin-type approximation theorems in several different areas of mathematics, such as (for example) probability space, measurable space, sequence spaces, and so on. In Real Analysis, Harmonic Analysis and other related fields, this notion is immensely useful. In this regard, we have chosen to refer the interested reader to the recent works (see, for example, [19–28]).

Let $C[0, 1]$ be the space of all continuous real-valued functions defined on $[0, 1]$. Suppose also that it is a Banach space with the norm $\|\cdot\|_\infty$. Then, for $h \in C[0, 1]$, the norm of h is given by

$$\|h\|_\infty = \sup\{|h(\rho)| : 0 \leq \rho \leq 1\}.$$

We say that $\mathfrak{G}_j : C[0, 1] \rightarrow C[0, 1]$ is a sequence of positive linear operators, if

$$\mathfrak{G}_j(h; \rho) \geq 0 \quad \text{as} \quad h \geq 0.$$

Now, in view of our above-proposed definitions, we state and prove the following Korovkin-type approximation theorems.

Theorem 3. Let $\mathfrak{G}_j : C[0, 1] \rightarrow C[0, 1]$ be a sequence of positive linear operators. Then, for $h \in C[0, 1]$,

$$\text{DWR}_{\text{stat}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(h; \rho) - h(\rho)\|_\infty = 0 \tag{4}$$

if and only if

$$\text{DWR}_{\text{stat}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(1; \rho) - 1\|_\infty = 0, \tag{5}$$

$$\text{DWR}_{\text{stat}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(\rho; \rho) - \rho\|_\infty = 0 \tag{6}$$

and

$$\text{DWR}_{\text{stat}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(\rho^2; \rho) - \rho^2\|_\infty = 0. \tag{7}$$

Proof. Since each of the following functions

$$h_0(\rho) = 1, \quad h_1(\rho) = 2\rho \quad \text{and} \quad h_2(\rho) = 3\rho^2$$

belongs to $C[0, 1]$ and is continuous on $[0, 1]$, the implication given by (4) obviously implies (5) to (7).

In order to complete the proof of Theorem 3, we first assume that the conditions (5) to (7) hold true. If $h \in C[0, 1]$, then there exists a constant $\mathcal{L} > 0$ such that

$$|h(\rho)| \leq \mathcal{L} \quad (\forall \rho \in [0, 1]).$$

We thus find that

$$|h(r) - h(\rho)| \leq 2\mathcal{L} \quad (r, \rho \in [0, 1]). \tag{8}$$

Clearly, for given $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(r) - f(\rho)| < \epsilon \tag{9}$$

whenever

$$|r - \rho| < \delta \quad \text{for all } r, \rho \in [0, 1].$$

If we now choose

$$\mu_1 = \mu_1(r, \rho) = (2r - 2\rho)^2.$$

If

$$|r - \rho| \geq \delta,$$

then we obtain

$$|h(r) - h(\rho)| < \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho). \tag{10}$$

Thus, from Equations (9) and (10), we get

$$|h(r) - h(\rho)| < \epsilon + \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho),$$

which implies that

$$-\epsilon - \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho) \leq h(r) - h(\rho) \leq \epsilon + \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho). \tag{11}$$

Now, since $\mathfrak{G}_m(1; \rho)$ is monotone and linear, by applying the operator $\mathfrak{G}_m(1; \rho)$ to the inequality (11), we get

$$\begin{aligned} \mathfrak{G}_m(1; \rho) \left(-\epsilon - \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho) \right) &\leq \mathfrak{G}_m(1; \rho) (h(r) - h(\rho)) \\ &\leq \mathfrak{G}_m(1; \rho) \left(\epsilon + \frac{2\mathcal{L}}{\theta^2} \mu_1(r, \rho) \right). \end{aligned}$$

We note that ρ is fixed, and so $h(\rho)$ is a constant number. Therefore, we have

$$\begin{aligned} -\epsilon \mathfrak{G}_m(1; \rho) - \frac{2\mathcal{L}}{\theta^2} \mathfrak{G}_m(\mu_1; \rho) &\leq \mathfrak{G}_m(h; \rho) - h(\rho) \mathfrak{G}_m(1; \rho) \\ &\leq \epsilon \mathfrak{G}_m(1; \rho) + \frac{2\mathcal{L}}{\theta^2} \mathfrak{G}_m(\mu_1; \rho). \end{aligned} \tag{12}$$

We also know that

$$\mathfrak{G}_m(h; \rho) - h(\rho) = [\mathfrak{G}_m(h; \rho) - h(\rho) \mathfrak{G}_m(1; \rho)] + h(\rho) [\mathfrak{G}_m(1; \rho) - 1]. \tag{13}$$

Thus, by using (12) and (13), we obtain

$$\mathfrak{G}_m(h; \rho) - h(\rho) < \epsilon \mathfrak{G}_m(1; \rho) + \frac{2\mathcal{L}}{\theta^2} \mathfrak{G}_m(\mu_1; \rho) + h(\rho) [\mathfrak{G}_m(1; \rho) - 1]. \tag{14}$$

We now estimate $\mathfrak{G}_m(\mu_1; \rho)$ as follows:

$$\begin{aligned} \mathfrak{G}_m(\mu_1; \rho) &= \mathfrak{G}_m((2r - 2\rho)^2; \rho) = \mathfrak{G}_m(2r^2 - 8\rho r + 4\rho^2; \rho) \\ &= \mathfrak{G}_m(4r^2; \rho) - 8t \mathfrak{G}_m(r; \rho) + 4\rho^2 \mathfrak{G}_m(1; \rho) \\ &= 4[\mathfrak{G}_m(r^2; \rho) - \rho^2] - 8t[\mathfrak{G}_m(r; \rho) - \rho] \\ &\quad + 4\rho^2[\mathfrak{G}_m(1; \rho) - 1], \end{aligned}$$

so that, in view of (14), we obtain

$$\begin{aligned} \mathfrak{G}_m(h; \rho) - h(\rho) &< \epsilon \mathfrak{G}_m(1; \rho) + \frac{2\mathcal{L}}{\theta^2} \{4[\mathfrak{G}_m(r^2; \rho) - \rho^2] \\ &\quad - 8\rho[\mathfrak{G}_m(r; \rho) - \rho] + 4\rho^2[\mathfrak{G}_m(1; \rho) - 1]\} \\ &\quad + h(\rho)[\mathfrak{G}_m(1; \rho) - 1]. \\ &= \epsilon[\mathfrak{G}_m(1; \rho) - 1] + \epsilon + \frac{2\mathcal{L}}{\theta^2} \{4[\mathfrak{G}_m(r^2; \rho) - \rho^2] \\ &\quad - 8\rho[\mathfrak{G}_m(r; \rho) - \rho] + 4\rho^2[\mathfrak{G}_m(1; \rho) - 1]\} \\ &\quad + h(\rho)[\mathfrak{G}_m(1; \rho) - 1]. \end{aligned}$$

Furthermore, since $\epsilon > 0$ is arbitrary, we can write

$$\begin{aligned} |\mathfrak{G}_m(h; \rho) - h(\rho)| &\leq \epsilon + \left(\epsilon + \frac{8\mathcal{L}}{\theta^2} + \mathcal{L} \right) |\mathfrak{G}_m(1; \rho) - 1| \\ &\quad + \frac{16\mathcal{L}}{\theta^2} |\mathfrak{G}_m(r; \rho) - \rho| + \frac{8\mathcal{L}}{\theta^2} |\mathfrak{G}_m(r^2; \rho) - \rho^2| \\ &\leq \mathcal{A} (|\mathfrak{G}_m(1; \rho) - 1| + |\mathfrak{G}_m(r; \rho) - \rho| \\ &\quad + |\mathfrak{G}_m(r^2; \rho) - \rho^2|), \end{aligned} \tag{15}$$

where

$$\mathcal{A} = \max \left(\epsilon + \frac{8\mathcal{L}}{\theta^2} + \mathcal{L}, \frac{16\mathcal{L}}{\theta^2}, \frac{8\mathcal{L}}{\theta^2} \right).$$

Now, for a given $\omega > 0$, there exists $\epsilon > 0$ ($\epsilon < \omega$) such that

$$\mathfrak{T}_m(\rho; \omega) = \{m : m \leq P_k \text{ and } p_m |\mathfrak{G}_m(h; \rho) - h(\rho)| \geq \omega\}.$$

Furthermore, for $\nu = 0, 1, 2$, we have

$$\mathfrak{T}_{\nu, m}(\rho; \omega) = \left\{ m : m \leq P_k \text{ and } p_m |\mathfrak{G}_m(h; \rho) - h_\nu(\rho)| \geq \frac{\omega - \epsilon}{3\mathcal{A}} \right\},$$

so that

$$\mathfrak{T}_m(\rho; \omega) \leq \sum_{\nu=0}^2 \mathfrak{T}_{\nu, m}(\rho; \omega).$$

Clearly, we obtain

$$\frac{\|\mathfrak{T}_m(\rho; \omega)\|_{\mathcal{C}[0,1]}}{P_k} \leq \sum_{\nu=0}^2 \frac{\|\mathfrak{T}_{\nu, m}(\rho; \omega)\|_{\mathcal{C}[0,1]}}{P_k}. \tag{16}$$

Now, using the above assumption about the implications in (5) to (7) and by Definition 2, the right-hand side of (16) tends to zero as $n \rightarrow \infty$. Consequently, we get

$$\lim_{k \rightarrow \infty} \frac{\|\mathfrak{T}_m(\rho; \omega)\|_{\mathcal{C}[0,1]}}{P_k} = 0 \quad (\delta, \omega > 0).$$

Therefore, the implication (4) holds true. \square

Theorem 4. Let $\mathfrak{G}_j : C[0,1] \rightarrow C[0,1]$ be a sequence of positive linear operators. Then, for $h \in C[0,1]$,

$$\text{stat}_{\text{DWR}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(h; \rho) - h(\rho)\|_\infty = 0 \tag{17}$$

if and only if

$$\text{stat}_{\text{DWR}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(1; \rho) - 1\|_\infty = 0, \tag{18}$$

$$\text{stat}_{\text{DWR}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(\rho; \rho) - \rho\|_\infty = 0 \tag{19}$$

and

$$\text{stat}_{\text{DWR}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_j(\rho^2; \rho) - \rho^2\|_\infty = 0. \tag{20}$$

Proof. The proof of Theorem 4 is similar to the proof of Theorem 3. Therefore, we choose to skip the details involved. \square

In view of Theorem 4, here, we consider an illustrative example. In this connection, we now recall the following operator:

$$\rho(1 + \rho D) \quad \left(D = \frac{d}{d\rho} \right), \tag{21}$$

which was used by Al-Salam [29] and, more recently, by Viskov and Srivastava [30].

Example 3. Consider the Bernstein polynomials $\mathfrak{B}_n(h; \beta)$ on $C[0, 1]$ given by

$$\mathfrak{B}_k(h; \beta) = \sum_{\varrho=0}^k f\left(\frac{\varrho}{k}\right) \binom{k}{\varrho} \beta^\varrho (1 - \beta)^{k-\varrho} \quad (\beta \in [0, 1]; k = 0, 1, \dots). \tag{22}$$

Here, in this example, we introduce the positive linear operators on $C[0, 1]$ under the composition of the Bernstein polynomials and the operators given by (21) as follows:

$$\mathfrak{G}_\varrho(h; \beta) = [1 + h_\varrho] \beta (1 + \beta D) \mathfrak{B}_\varrho(h; \beta) \quad (\forall h \in C[0, 1]), \tag{23}$$

where (h_ϱ) is the same as mentioned in Example 2.

We now estimate the values of each of the testing functions 1, β and β^2 by using our proposed operators (23) as follows:

$$\mathfrak{G}_\varrho(1; \beta) = [1 + h_\varrho] \beta (1 + \beta D) 1 = [1 + h_\varrho] \beta,$$

$$\mathfrak{G}_\varrho(t; \beta) = [1 + h_\varrho] \beta (1 + \beta D) \beta = [1 + h_\varrho] \beta (1 + \beta)$$

and

$$\begin{aligned} \mathfrak{G}_\varrho(t^2; \beta) &= [1 + h_\varrho] \beta (1 + \beta D) \left\{ \beta^2 + \frac{\beta(1 - \beta)}{\varrho} \right\} \\ &= [1 + h_\varrho] \left\{ \beta^2 \left(2 - \frac{3\beta}{\varrho} \right) \right\}. \end{aligned}$$

Consequently, we have

$$\text{stat}_{\text{DWR}} \lim_{\varrho \rightarrow \infty} \|\mathfrak{G}_\varrho(1; \beta) - 1\|_\infty = 0, \tag{24}$$

$$\text{stat}_{\text{DWR}} \lim_{\varrho \rightarrow \infty} \|\mathfrak{G}_\varrho(\beta; \beta) - \beta\|_\infty = 0 \tag{25}$$

and

$$\text{stat}_{\text{DWR}} \lim_{\varrho \rightarrow \infty} \|\mathfrak{G}_\varrho(\beta^2; \beta) - \beta^2\|_\infty = 0, \tag{26}$$

that is, the sequence $\mathfrak{G}_\rho(h; \beta)$ satisfies the conditions (18) to (20). Therefore, by Theorem 4, we have

$$\text{stat}_{\text{DWR}} \lim_{\rho \rightarrow \infty} \|\mathfrak{G}_\rho(h; \beta) - h\|_\infty = 0.$$

Hence, the given sequence (h_k) of functions mentioned in Example 2 is statistically deferred weighted Riemann-summable, but not deferred weighted statistically Riemann-integrable. Therefore, our above-proposed operators defined by (23) satisfy Theorem 4. However, they do not satisfy for statistical versions of deferred weighted Riemann-integrable sequence of functions (see Theorem 3).

4. Concluding Remarks and Directions for Further Research

In this concluding section of our present investigation, we further observe the potential usefulness of our Theorem 4 over Theorem 3 as well as over the classical versions of the Korovkin-type approximation theorems.

Remark 2. Let us consider the sequence $(h_\rho)_{\rho \in \mathbb{N}}$ of functions in Example 2. Suppose also that (h_ρ) is statistically deferred weighted Riemann-summable, so that

$$\text{stat}_{\text{DWR}} \lim_{\rho \rightarrow \infty} \text{ffi}(h_\rho; \mathcal{P}) = \frac{1}{2} \text{ on } [0, 1].$$

We then find that

$$\text{stat}_{\text{DWR}} \lim_{k \rightarrow \infty} \|\mathfrak{G}_k(h_\rho; \mathfrak{a}) - f_\rho(\mathfrak{a})\|_\infty = 0 \quad (\rho = 0, 1, 2). \quad (27)$$

Thus, by Theorem 4, we immediately get

$$\text{stat}_{\text{DWR}} \lim_{j \rightarrow \infty} \|\mathfrak{G}_k(h; \mathfrak{a}) - h(\mathfrak{a})\|_\infty = 0, \quad (28)$$

where

$$h_0(\rho) = 1, \quad h_1(\rho) = \rho \quad \text{and} \quad h_2(\rho) = \rho^2.$$

Now, the given sequence (h_k) of functions is statistically deferred weighted Riemann-summable, but neither deferred weighted statistically Riemann-integrable nor classically Riemann-integrable. Therefore, our Korovkin-type approximation Theorem 4 properly works under the operators defined in the Equation (23), but the classical as well as statistical versions of the deferred weighted Riemann-integrable sequence of functions do not work for the same operators. Clearly, this observation leads us to the fact that our Theorem 4 is a non-trivial extension of Theorem 3 as well as the classical Korovkin-type approximation theorem [31].

Remark 3. Motivated by some recently published results by Jena et al. [32] and Srivastava et al. [33], we choose to draw the attention of the interested readers toward the potential for further research associated with the analogous notion of statistical Lebesgue-measurable sequences of functions.

Author Contributions: Formal analysis, H.M.S. and S.K.P.; Investigation, B.B.J.; Methodology, S.K.P.; Supervision, H.M.S. and S.K.P.; Writing—original draft, B.B.J.; Writing—review & editing, H.M.S. and S.K.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare that they have no conflicts of interest.

References

1. Fast, H. Sur la convergence statistique. *Colloq. Math.* **1951**, *2*, 241–244. [\[CrossRef\]](#)
2. Steinhaus, H. Sur la convergence ordinaire et la convergence asymptotique. *Colloq. Math.* **1951**, *2*, 73–74.

3. Akdag, S. Weighted equi-statistical convergence of the Korovkin-type approximation theorems. *Results Math.* **2017**, *72*, 1073–1085. [[CrossRef](#)]
4. Balcerzak, M.; Dems, K.; Komisarski, A. Statistical convergence and ideal convergence for sequences of functions. *J. Math. Anal. Appl.* **2007**, *328*, 715–729. [[CrossRef](#)]
5. Braha, N.L.; Loku, V.; Srivastava, H.M. Λ^2 -Weighted statistical convergence and Korovkin and Voronovskaya type theorems. *Appl. Math. Comput.* **2015**, *266*, 675–686. [[CrossRef](#)]
6. Et, M.; Baliarsingh, P.; Küçxukaslan, H.S.K.M. On μ -deferred statistical convergence and strongly deferred summable functions. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2021**, *115*, 1–14. [[CrossRef](#)]
7. Ghosal, S.; Mandal, S. Rough weighted I - $\alpha\beta$ -statistical convergence in locally solid Riesz spaces. *J. Math. Anal. Appl.* **2021**, *506*, 125681. [[CrossRef](#)]
8. Ghosal, S.; Banerjee, M. Rough weighted statistical convergence on locally solid Riesz spaces. *Positivity* **2021**, *25*, 1789–1804. [[CrossRef](#)]
9. Guessab, A.; Schmeisser, G. Convexity results and sharp error estimates in approximate multivariate integration. *Math. Comput.* **2003**, *73*, 1365–1384. [[CrossRef](#)]
10. Alotaibi, A. Generalized weighted statistical convergence for double sequences of fuzzy numbers and associated Korovkin-type approximation theorem. *J. Funct. Spaces* **2020**, *2020*, 9298650. [[CrossRef](#)]
11. Özger, F. Applications of generalized weighted statistical convergence to approximation theorems for functions of one and two variables. *Numer. Funct. Anal. Optim.* **2020**, *41*, 1990–2006. [[CrossRef](#)]
12. Demirci, K.; Dirik, F.; Yıldız, S. Deferred Nörlund statistical relative uniform convergence and Korovkin-type approximation theorem. *Commun. Fac. Sci. Univ. Ank. Ser. A Math. Statist.* **2021**, *70*, 279–289. [[CrossRef](#)]
13. Mohiuddine, S.A.; Hazarika, B. On strongly almost generalized difference lacunary ideal convergent sequences of fuzzy numbers. *J. Comput. Anal. Appl.* **2017**, *23*, 925–936.
14. Móricz, F. Tauberian conditions under which statistical convergence follows from statistical summability $(C, 1)$. *J. Math. Anal. Appl.* **2002**, *275*, 277–287. [[CrossRef](#)]
15. Agrawal, P.N.; Acu, A.-M.; Chauhan, R.; Garg, T. Approximation of Bögel continuous functions and deferred weighted A -statistical convergence by Bernstein-Kantorovich type operators on a triangle. *J. Math. Inequal.* **2021**, *15*, 1695–1711. [[CrossRef](#)]
16. Saini, K.; Raj, K.; Mursaleen, M. Deferred Cesàro and deferred Euler equi-statistical convergence and its applications to Korovkin-type approximation theorem. *Internat. J. Gen. Syst.* **2021**, *50*, 567–579. [[CrossRef](#)]
17. Söylemez, D. A Korovkin type approximation theorem for Balázs type Bleimann, Butzer and Hahn operators via power series statistical convergence. *Math. Slovaca* **2022**, *72*, 153–164. [[CrossRef](#)]
18. Turan, C.; Duman, O. Fundamental properties of statistical convergence and lacunary statistical convergence on time scales. *Filomat* **2017**, *31*, 4455–4467. [[CrossRef](#)]
19. Altomare, F. Korovkin-type theorems and approximation by positive linear operators. *Surv. Approx. Theory* **2010**, *5*, 92–164.
20. Braha, N.L. Some weighted equi-statistical convergence and Korovkin-type theorem. *Results Math.* **2016**, *70*, 433–446. [[CrossRef](#)]
21. Braha, N.L.; Srivastava, H.M.; Et, M. Some weighted statistical convergence and associated Korovkin and Voronovskaya type theorems. *J. Appl. Math. Comput.* **2021**, *65*, 429–450. [[CrossRef](#)]
22. Guessab, A.; Schmeisser, G. Two Korovkin-type theorems in multivariate approximation. *Banach J. Math. Anal.* **2008**, *2*, 121–128. [[CrossRef](#)]
23. Jena, B.B.; Paikray, S.K.; Dutta, H. On various new concepts of statistical convergence for sequences of random variables via deferred Cesàro mean. *J. Math. Anal. Appl.* **2020**, *487*, 123950. [[CrossRef](#)]
24. Karakuş, S.; Demirci, K.; Duman, O. Equi-statistical convergence of positive linear operators. *J. Math. Anal. Appl.* **2008**, *339*, 1065–1072. [[CrossRef](#)]
25. Çınar, S.; Yıldız, S.; Demirci, K. Korovkin type approximation via triangular A -statistical convergence on an infinite interval. *Turk. J. Math.* **2021**, *45*, 929–942. [[CrossRef](#)]
26. Parida, P.; Paikray, S.K.; Jena, B.B. Generalized deferred Cesàro equi-statistical convergence and analogous approximation theorems. *Proyecciones J. Math.* **2020**, *39*, 307–331. [[CrossRef](#)]
27. Srivastava, H.M.; Jena, B.B.; Paikray, S.K. Statistical probability convergence via the deferred Nörlund mean and its applications to approximation theorems. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2020**, *114*, 1–14. [[CrossRef](#)]
28. Srivastava, H.M.; Jena, B.B.; Paikray, S.K. Statistical deferred Nörlund summability and Korovkin-type approximation theorem. *Mathematics* **2020**, *8*, 636. [[CrossRef](#)]
29. Al-Salam, W.A. Operational representations for the Laguerre and other polynomials. *Duke Math. J.* **1964**, *31*, 127–142. [[CrossRef](#)]
30. Viskov, O.V.; Srivastava, H.M. New approaches to certain identities involving differential operators. *J. Math. Anal. Appl.* **1994**, *186*, 1–10. [[CrossRef](#)]
31. Korovkin, P.P. Convergence of linear positive operators in the spaces of continuous functions. *Dokl. Akad. Nauk. SSSR (New Ser.)* **1953**, *90*, 961–964. (In Russian)
32. Jena, B.B.; Paikray, S.K.; Dutta, H. A new approach to Korovkin-type approximation via deferred Cesàro statistical measurable convergence. *Chaos Solitons Fractals* **2021**, *148*, 111016. [[CrossRef](#)]
33. Srivastava, H.M.; Jena, B.B.; Paikray, S.K. Statistical Riemann and Lebesgue integrable sequence of functions with Korovkin-type approximation theorems. *Axioms* **2021**, *10*, 229. [[CrossRef](#)]