

Table S1a: logistic regression analysis considering only sociodemographic characteristics associated with FSUs

| Model 1 | B | S.E. | Wald | p | Exp(B) | 95% CI for EXP |
|----------------|--------|-------|-------|-------|--------|----------------|
| Current Age | -0.025 | 0.010 | 6.689 | 0.010 | 0.975 | 0.956-0.994 |
| Single status | 0.169 | 0.370 | .207 | 0.649 | 1.184 | .573-2.445 |
| Male gender | -0.269 | .277 | .943 | 0.332 | 0.764 | 0.444-1.316 |
| Constant | -0.899 | 0.595 | 2.283 | 0.131 | 0.407 | |

R² Nagelkerke=0.041

Table S1b: logistic regression analysis considering sociodemographic and clinical characteristics associated with FSUs

| Model 2 | B | S.E. | Wald | p | Exp(B) | 95% CI for EXP |
|----------------------------|--------|-------|-------|-------|--------|----------------|
| Current Age | 0.012 | 0.013 | 0.909 | 0.340 | 1.012 | 0.987-1.037 |
| Single status | -0.014 | 0.407 | 0.001 | 0.974 | 0.987 | .444-2.191 |
| Male gender | -0.408 | 0.327 | 1.555 | 0.212 | 0.665 | 0.350-1.263 |
| Personality disorder | 0.766 | 0.390 | 3.865 | 0.049 | 2.152 | 1.002-4.620 |
| Substance use disorder | 0.349 | 0.519 | 0.454 | 0.500 | 1.418 | 0.513-3.921 |
| Age at onset | -0.050 | 0.018 | 7.631 | 0.006 | 0.951 | 0.918-0.986 |
| Length of hospitalisation | 0.030 | 0.012 | 5.970 | 0.015 | 1.031 | 1.006-1.056 |
| Non suicidal self-injuries | 0.567 | 0.465 | 1.486 | 0.223 | 1.763 | 0.709-4.385 |
| Psychiatric comorbidity | 0.086 | 0.360 | 0.058 | 0.810 | 1.090 | 0.538-2.207 |
| Presence of illicit drugs | 0.067 | 0.487 | 0.019 | 0.890 | 1.069 | 0.412-2.778 |
| Alcohol | -0.182 | 0.400 | 0.208 | 0.648 | 0.833 | 0.381-1.824 |
| Cannabinoid | 0.994 | 0.437 | 5.187 | 0.023 | 2.703 | 1.149-6.362 |
| Cocaine | 0.387 | 0.402 | 0.923 | 0.337 | 1.472 | 0.669-3.239 |
| Benzodiazepine therapy | 0.548 | 0.362 | 2.286 | 0.131 | 1.730 | 0.850-3.520 |
| Other therapy | 0.524 | 0.345 | 2.299 | 0.129 | 1.688 | 0.858-3.322 |
| Constant | -2.557 | 0.858 | 8.890 | 0.003 | 0.078 | |

R² Nagelkerke=0.224