

**Table S6 – Summary of histological forms of head and neck tumors afflicted by PNS**

First author	Nader et al.	Baulch et al.	Gandhi et al.	Nemzek et al.	Tomura et al.	Schmalfuss et al.	Hanna et al.	Warren et al.	Chang et al.	Shimamoto et al.	Majoie et al.	Summary
<b>Patients' number with PNS who had reference standard test</b>	<b>21 (out of 58)</b>	<b>33</b>	<b>25</b>	<b>19</b>	<b>12</b>	<b>7 (out of 15)</b>	<b>26</b>	<b>48</b>	<b>8</b>	<b>11 (out of 13)</b>	<b>2 (out of 15)</b>	<b>171</b>
<b>Tumor histology</b>												
SCC	9 (42.9%)	27(81.8%)	22(88%)	10(52.6%)	11(91.7%)	2(28.6%)		48(100%)*				<b>129</b>
ACC	3(14.3%)	2(6.1%)		4(21%)		2(28.6%)	26(100%)			11(100%)		<b>48</b>
Melanoma		3(9.1%)	1(4%)#						8(100%)		1(50%)	<b>13</b>
Mucoepidermoid carcinoma	1 (4.8%)			1(5.3%)		1(14.3%)						<b>3</b>
Salivary ductal carcinoma	3(14.3%)			1(5.3%)								<b>4</b>
Salivary gland tumor		1(3%)										<b>1</b>
nasopharyngeal angiofibroma					1(8.3%)							<b>1</b>
BCC			1(4%)									<b>1</b>
BCC with squamous differentiation	1 (4.8%)											<b>1</b>
small round cell neoplasm												
Lymphoma											1(50%)	<b>1</b>



[illegible]