

Table S1: Regression results for percentile plots of sonographic parameters. Model fit was performed for 10th, 50th and 90th percentile for each sonographic ratio fitted by: ratio ~ **a+b*GA+c*GA²+d*GA³**. AC = abdominal circumference, BPD = biparietal diameter, CI = confidence interval, FL = femur length, HC = head circumference, pct = percentile, TCD = transcerebellar diameter.

Ratio	Estimate	95% CI	<i>p</i> value
HC (10th pct)			
a	-5.5328	-21.4299 – 10.3644	0.4951
b	2.1830	0.0809 – 4.2852	0.0418
c	0.4936	0.4037 – 0.5836	<.0001
d	-0.0086	-0.0098 – -0.0073	<.0001
HC (50th pct)			
a	-3.9716	-17.7150 – 9.7719	0.5711
b	2.1841	0.4173 – 3.9508	0.0154
c	0.5179	0.4441 – 0.5917	<.0001
d	-0.0090	-0.0100 – -0.0080	<.0001
HC (90th pct)			
a	13.2957	-5.5018 – 32.0932	0.1656
b	0.3035	-2.1751 – 2.7821	0.8103
c	0.6126	0.5066 – 0.7186	<.0001
d	-0.0102	-0.0117 – -0.0088	<.0001
BPD (10th pct)			
a	8.3921	2.5741 – 14.2102	0.0047
b	-0.4590	-1.2274 – 0.3093	0.2416
c	0.1721	0.1394 – 0.2048	<.0001
d	-0.0027	-0.0032 – -0.0023	<.0001
BPD (50th pct)			
a	2.2356	-2.8324 – 7.3037	0.3872
b	0.3997	-0.2703 – 1.0696	0.2423
c	0.1463	0.1175 – 0.1751	<.0001
d	-0.0024	-0.0028 – -0.0021	<.0001
BPD (90th pct)			
a	6.7349	0.7516 – 12.7182	0.0274
b	-0.0910	-0.8707 – 0.6888	0.8191
c	0.1743	0.1413 – 0.2073	<.0001
d	-0.0028	-0.0033 – -0.0024	<.0001
AC (10th pct)			
a	-69.0745	-87.7419 – -50.4071	<.0001
b	9.5145	7.0579 – 11.9711	<.0001
c	0.0936	-0.0110 – 0.1983	0.0794
d	-0.0020	-0.0034 – -0.0006	0.0061
AC (50th pct)			
a	-70.9656	-84.3577 – -57.5735	<.0001
b	10.2846	8.5417 – 12.0275	<.0001
c	0.0727	-0.0011 – 0.1465	0.0535
d	-0.0016	-0.0026 – -0.0006	0.0022
AC (90th pct)			
a	-46.3402	-68.7357 – -23.9446	<.0001
b	7.4990	4.5507 – 10.4472	<.0001
c	0.2056	0.0804 – 0.3309	0.0013
d	-0.0033	-0.0050 – -0.0016	0.0002
FL (10th pct)			

a	-39.8647	-44.1893 – -35.5401	<.0001
b	4.2383	3.6761 – 4.8005	<.0001
c	-0.0348	-0.0583 – -0.0112	0.0038
d	0.0000	-0.0004 – 0.0003	0.8293
FL (50th pct)			
a	-40.2490	-43.2771 – -37,2210	<.0001
b	4.4077	4.0241 – 4.7912	<.0001
c	-0.0385	-0.0544 – -0.0227	<.0001
d	0.0000	-0.0002 – 0.0002	0.8453
FL (90th pct)			
a	-29.7584	-33.4226 – -26.0942	<.0001
b	3.2440	2.7802 – 3.7078	<.0001
c	0.0114	-0.0077 – 0.0305	0.2405
d	-0.0006	-0.0009 – -0.0004	<.0001
TCD (10th pct)			
a	16.3571	11.5841 – 21.1301	<.0001
b	-1.5250	-2.1720 – -0.8780	<.0001
c	0.1092	0.0809 – 0.1376	<.0001
d	-0.0013	-0.0017 – -0,0009	<.0001
TCD (50th pct)			
a	17.1858	13.3614 – 21.0101	<.0001
b	-1.4903	-2.0216 – -0.9590	<.0001
c	0.1048	0.0809 – 0.1287	<.0001
d	-0.0011	-0.0014 – -0,0008	<.0001
TCD (90th pct)			
a	18.3239	12.7680 – 23.8798	<.0001
b	-1.4625	-2.2075 – -0.7176	<.0001
c	0.1004	0.0680 – 0.1327	<.0001
d	-0.0009	-0.0014 – -0,0005	<.0001
FL/AC (10th pct)			
a	-0.1660	-0.1934 – -0.1386	<.0001
b	0.0385	0.0352 – 0.0418	<.0001
c	-0.0013	-0.0014 – -0,0012	<.0001
d	0.0000	0.0000 – 0.0000	<.0001
FL/AC (50th pct)			
a	-0.1325	-0.1544 – -0,1106	<.0001
b	0.0364	0.0337 – 0.0391	<.0001
c	-0.0012	-0.0013 – -0,0011	<.0001
d	0.0000	0.0000 – 0.0000	<.0001
FL/AC (90th pct)			
a	-0.0662	-0.1013 – -0,0311	0.0002
b	0.0308	0.0266 – 0.0351	<.0001
c	-0.0010	-0.0012 – -0,0008	<.0001
d	0.0000	0.0000 – 0.0000	<.0001
TCD/AC (10th pct)			
a	34.1176	31.3786 – 36.8566	<.0001
b	-2.3949	-2.7367 – -2.0530	<.0001
c	0.0853	0.0713 – 0.0993	<.0001
d	-0.0010	-0.0012 – -0.0008	<.0001
TCD/AC (50th pct)			
a	39.1917	37.5438 – 40.8396	<.0001
b	-2.8566	-3.0659 – -2.6473	<.0001

c	0.1018	0.0931 – 0.1104	<.0001
d	-0.0011	-0.0013 – -0.0010	<.0001
TCD/AC (90th pct)			
a	45.2301	42.3682 – 48.0919	<.0001
b	-3.3964	-3.7590 – -3.0337	<.0001
c	0.1200	0.1050 – 0.1350	<.0001
d	-0.0013	-0.0015 – -0.0011	<.0001
HC/AC (10th pct)			
a	15.3218	14.1559 – 16.4878	<.0001
b	-0.4693	-0.6111 – -0.3276	<.0001
c	0.0172	0.0116 – 0.0228	<.0001
d	-0.0002	-0.0003 – -0.0002	<.0001
HC/AC (50th pct)			
a	18.2726	17.1183 – 19.4270	<.0001
b	-0.7345	-0.8759 – -0.5932	<.0001
c	0.0272	0.0216 – 0.0328	<.0001
d	-0.0004	-0.0004 – -0.0003	<.0001
HC/AC (90th pct)			
a	18.5081	16.8963 – 20.1200	<.0001
b	-0.6491	-0.8422 – -0.4559	<.0001
c	0.0231	0.0156 – 0.0305	<.0001
d	-0.0003	-0.0004 – -0.0002	<.0001

Table S2: Regression results for percentile plots of sonographic parameters. Model fit was performed for 3th and 97th percentile for each sonographic ratio fitted by: $\text{ratio} \sim a + b \cdot \text{GA} + c \cdot \text{GA}^2 + d \cdot \text{GA}^3$. AC = abdominal circumference, BPD = biparietal diameter, CI = confidence interval, FL = femur length, HC = head circumference, pct=percentile, TCD = transcerebellar diameter.

ratio	Estimate	95% CI	<i>p</i> value
HC (3th pct)			
a	-10.2006	-33.4466 – 13.0455	0.3897
b	2.5757	-0.4888 – 5.6403	0.0995
c	0.4719	0.3415 – 0.6024	<.0001
d	-0.0083	-0.0101 – -0.0065	<.0001
HC (97th pct)			
a	16.2585	-10.0805 – 42.5975	0.2263
b	0.1701	-3.3262 – 3.6665	0.9240
c	0.6214	0.4706 – 0.7722	<.0001
d	-0.0103	-0.0124 – -0.0082	<.0001
BPD (3th pct)			
a	13.4294	4.4180 – 22.4407	0.0035
b	-1.1306	-2.2904 – 0.0292	0.0561
c	0.1943	0.1461 – 0.2425	<.0001
d	-0.0030	-0.0036 – -0.0023	<.0001
BPD (97th pct)			
a	12.7560	3.4725 – 22.0395	0.0071
b	-0.7646	-1.9337 – 0.4046	0.1999
c	0.2035	0.1560 – 0.2511	<.0001
d	-0.0032	-0.0038 – -0.0026	<.0001
AC (3th pct)			
a	-63.2831	-88.4957 – -38.0704	<.0001
b	8.6451	5.3611 – 11.9292	<.0001
c	0.1172	-0.0197 – 0.2542	0.0933
d	-0.0023	-0.0041 – -0.0004	0.0147
AC (97th pct)			
a	-33.1537	-63.8501 – -2.4573	0.0343
b	5.9026	2.0297 – 9.7755	0.0028
c	0.2862	0.1275 – 0.4448	0.0004
d	-0.0044	-0.0065 – -0.0023	<.0001
FL (3th pct)			
a	-37.3809	-42.98931 – -31.7686	<.0001
b	3.9113	3.2008 – 4.6218	<.0001
c	-0.0243	-0.0536 – 0.0050	0.1035
d	-0.0002	-0.0005 – 0.0002	0.4323
FL (97th pct)			
a	-25.7335	-32.4664 – -19.0006	<.0001
b	2.7815	1.9403 – 3.6227	<.0001
c	0.0329	-0.0013 – 0.0670	0.0593
d	-0.0009	-0.0014 – -0.0005	<.0001
TCD (3th pct)			
a	7.4550	0.2833 – 14.6268	0.0416
b	-0.4535	-1.4037 – 0.4968	0.3496
c	0.0666	0.0259 – 0.1073	0.0013
d	-0.0008	-0.0014 – -0.0002	0.0049
TCD (97th pct)			

a	20.8028	6.7502 – 34.8553	0.0037
b	-1.7493	-3.7387 – 0.2400	0.0848
c	0.1122	0.0209 – 0.2035	0.0160
d	-0.0010	-0.0024 – 0.0003	0.1384
FL/AC (3th pct)			
a	-0.1938	-0.2343 – -0.1532	<.0001
b	0.0412	0.0362 – 0.0463	<.0001
c	-0.0014	-0.0016 – -0.0012	<.0001
d	0.0000	0.0000 – 0.0000	<.0001
FL/AC (97th pct)			
a	-0.0506	-0.1246 – 0.0234	0.1804
b	0.0299	0.0208 – 0.0389	<.0001
c	-0.0010	-0.0013 – -0.0006	<.0001
d	0.0000	0.0000 – 0.0000	<.0001
TCD/AC (3th pct)			
a	31.9001	28.1809 – 35.6192	<.0001
b	-2.2270	-2.6940 – -1.7600	<.0001
c	0.0810	0.0618 – 0.1002	<.0001
d	-0.0010	-0.0012 – -0.0007	<.0001
TCD/AC (97th pct)			
a	48.6127	43.1549 – 54.0704	<.0001
b	-3.7614	-4.4600 – -3.0629	<.0001
c	0.1347	0.1055 – 0.1638	<.0001
d	-0.0015	-0.0019 – -0.0011	<.0001
HC/AC (3th pct)			
a	12.9014	11.3303 – 14.4725	<.0001
b	-0.2144	-0.4006 – -0.0281	0.0241
c	0.0071	-0.0001 – 0.0143	0.0522
d	-0.0001	-0.0002 – 0.0000	0.0185
HC/AC (97th pct)			
a	20.7330	18.3564 – 23.1095	<.0001
b	-0.8811	-1.1565 – -0.6056	<.0001
c	0.0325	0.0222 – 0.0429	<.0001
d	-0.0004	-0.0006 – -0.0003	<.0001

Table S3: Results of logistic regression analysis for the gestational-age depended association between sonographic parameters and small gestational age. Model fit was performed for each parameter separately, including gestational age, sonographic parameter and an interaction term between gestational age and parameter. AC = abdominal circumference, BPD = biparietal diameter, CI = confidence interval, FL = femur length, HC = head circumference, TCD = transcerebellar diameter.

Sonographic parameter	Gestational age in weeks	Odds ratio	95% CI	<i>p</i> value
HC	14+0 – 18+6	0.977	0.963 – 0.991	0.0013
	19+0 – 23+6	0.965	0.957 – 0.974	<.0001
	24+0 – 28+6	0.952	0.944 – 0.961	<.0001
	29+0 – 33+6	0.931	0.920 – 0.941	<.0001
	34+0 – 38+6	0.888	0.868 – 0.908	<.0001
	≥39+0	0.897	0.802 – 1.004	0.0595
BPD	14+0 – 18+6	0.933	0.888 – 0.979	0.0052
	19+0 – 23+6	0.893	0.869 – 0.917	<.0001
	24+0 – 28+6	0.858	0.833 – 0.883	<.0001
	29+0 – 33+6	0.801	0.772 – 0.831	<.0001
	34+0 – 38+6	0.707	0.662 – 0.754	<.0001
	≥39+0	0.709	0.492 – 1.022	0.0649
AC	14+0 – 18+6	0.968	0.953 – 0.983	<.0001
	19+0 – 23+6	0.935	0.926 – 0.943	<.0001
	24+0 – 28+6	0.927	0.916 – 0.938	<.0001
	29+0 – 33+6	0.920	0.909 – 0.931	<.0001
	34+0 – 38+6	0.885	0.866 – 0.904	<.0001
	≥39+0	0.849	0.692 – 1.042	0.1167
FL	14+0 – 18+6	0.884	0.835 – 0.936	<.0001
	19+0 – 23+6	0.805	0.776 – 0.834	<.0001
	24+0 – 28+6	0.773	0.740 – 0.807	<.0001
	29+0 – 33+6	0.735	0.701 – 0.771	<.0001
	34+0 – 38+6	0.700	0.651 – 0.753	<.0001
	≥39+0	0.314	0.091 – 1.087	0.0675
TCD	14+0 – 18+6	0.873	0.772 – 0.986	0.0284
	19+0 – 23+6	0.861	0.811 – 0.913	<.0001
	24+0 – 28+6	0.854	0.811 – 0.898	<.0001
	29+0 – 33+6	0.920	0.886 – 0.955	<.0001
	34+0 – 38+6	0.891	0.850 – 0.933	<.0001
	≥39+0	0.806	0.606 – 1.073	0.1401
FL/AC	14+0 – 18+6	0.839	0.739 – 0.953	0.0068
	19+0 – 23+6	1.053	0.979 – 1.133	0.1639
	24+0 – 28+6	1.247	1.122 – 1.385	<.0001
	29+0 – 33+6	1.396	1.255 – 1.553	<.0001
	34+0 – 38+6	1.977	1.675 – 2.333	<.0001
	≥39+0	2.973	0.886 – 9.973	0.0776
TCD/AC	14+0 – 18+6	1.668	1.72 – 2.028	<.0001

	19+0 – 23+6	2.249	1.999 – 2.531	<.0001
	24+0 – 28+6	2.421	2.065 – 2.838	<.0001
	29+0 – 33+6	1.962	1.732 – 2.224	<.0001
	34+0 – 38+6	1.600	1.394 – 1.836	<.0001
	>=39+0	1.252	0.713 – 2.199	0.4342
HC/AC	14+0 – 18+6	1.069	1.033 – 1.106	0.0001
	19+0 – 23+6	1.125	1.105 – 1.144	<.0001
	24+0 – 28+6	1.171	1.139 – 1.203	<.0001
	29+0 – 33+6	1.155	1.125 – 1.185	<.0001
	34+0 – 38+6	1.172	1.129 – 1.215	<.0001
	>=39+0	1.184	0.972 – 1.443	0.0941

Table S4: Results of logistic regression analysis for the gestational-age depended association between sonographic parameters and fetal growth restriction. Model fit was performed for each parameter separately, including gestational age, sonographic parameter and an interaction term between gestational age and parameter. AC = abdominal circumference, BPD = biparietal diameter, CI = confidence interval, FL = femur length, HC = head circumference, TCD = transcerebellar diameter.

Sonographic parameter	Gestational age in weeks	Odds ratio	95% CI	<i>p</i> value
HC	14+0 – 18+6	0.960	0.915 – 1.007	0.0912
	19+0 – 23+6	0.956	0.935 – 0.977	<.0001
	24+0 – 28+6	0.921	0.904 – 0.938	<.0001
	29+0 – 33+6	0.908	0.890 – 0.926	<.0001
	34+0 – 38+6	0.852	0.818 – 0.887	<.0001
	≥39+0	0.679	0.091 – 5.061	0.7057
BPD	14+0 – 18+6	0.906	0.774 – 1.060	0.2172
	19+0 – 23+6	0.837	0.778 – 0.901	<.0001
	24+0 – 28+6	0.775	0.732 – 0.821	<.0001
	29+0 – 33+6	0.757	0.713 – 0.804	<.0001
	34+0 – 38+6	0.626	0.558 – 0.703	<.0001
	≥39+0	0.038	NA	0.8967
AC	14+0 – 18+6	0.951	0.902 – 1.002	0.0614
	19+0 – 23+6	0.893	0.869 – 0.918	<.0001
	24+0 – 28+6	0.886	0.864 – 0.909	<.0001
	29+0 – 33+6	0.890	0.869 – 0.911	<.0001
	34+0 – 38+6	0.837	0.798 – 0.878	<.0001
	≥39+0	0.684	NA	0.9619
FL	14+0 – 18+6	0.822	0.676 – 0.999	0.0488
	19+0 – 23+6	0.694	0.625 – 0.770	<.0001
	24+0 – 28+6	0.663	0.608 – 0.724	<.0001
	29+0 – 33+6	0.654	0.601 – 0.713	<.0001
	34+0 – 38+6	0.579	0.503 – 0.666	<.0001
	≥39+0	0.005	NA	0.8686
TCD	14+0 – 18+6	0.865	0.595 – 1.259	0.4494
	19+0 – 23+6	0.906	0.784 – 1.047	0.1816
	24+0 – 28+6	0.756	0.690 – 0.828	<.0001
	29+0 – 33+6	0.908	0.854 – 0.966	0.0023
	34+0 – 38+6	0.874	0.812 – 0.940	0.0003
	≥39+0	1.133	0.632 – 2.030	0.6755
FL/AC	14+0 – 18+6	0.753	0.506 – 1.121	0.1625
	19+0 – 23+6	1.052	0.876 – 1.265	0.5853
	24+0 – 28+6	1.231	1.036 – 1.463	0.0183
	29+0 – 33+6	1.361	1.149 – 1.612	0.0004
	34+0 – 38+6	2.209	1.736 – 2.811	<.0001
	≥39+0	NA	NA	0.9151
TCD/AC	14+0 – 18+6	2.165	1.271 – 3.686	0.0045

	19+0 – 23+6	3.327	2.581 – 4.288	<.0001
	24+0 – 28+6	3.173	2.474 – 4.070	<.0001
	29+0 – 33+6	2.511	2.053 – 3.072	<.0001
	34+0 – 38+6	2.000	1.612 – 2.482	<.0001
	>=39+0	NA	NA	0.9533
HC/AC	14+0 – 18+6	1.068	0.965 – 1.183	0.2048
	19+0 – 23+6	1.193	1.147 – 1.241	<.0001
	24+0 – 28+6	1.236	1.182 – 1.293	<.0001
	29+0 – 33+6	1.192	1.145 – 1.241	<.0001
	34+0 – 38+6	1.223	1.155 – 1.295	<.0001
	>=39+0	5.035	NA	0.9379

Table S5: Evaluation of gestational age-dependent cut-off values, 10th percentile (SGA). AC = abdominal circumference, BPD = biparietal diameter, FL = femur length, GA = Gestational age, HC = head circumference, TCD = transcerebellar diameter.

Parameter	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Youden Index (J)
HC					
Total	0.55	0.91	0.43	0.94	0.46
GA: 14+0– 18+6	0.35	0.90	0.18	0.95	0.25
19+0– 23+6	0.44	0.91	0.30	0.95	0.35
24+0– 28+6	0.65	0.90	0.56	0.92	0.55
29+0– 33+6	0.65	0.93	0.69	0.92	0.58
34+0– 38+6	0.64	0.90	0.65	0.90	0.54
BPD					
Total	0.51	0.88	0.35	0.93	0.39
GA: 14+0– 18+6	0.28	0.88	0.14	0.95	0.16
19+0– 23+6	0.36	0.88	0.21	0.94	0.24
24+0– 28+6	0.64	0.87	0.51	0.92	0.51
29+0– 33+6	0.65	0.87	0.55	0.91	0.52
34+0– 38+6	0.67	0.86	0.57	0.90	0.53
AC					
Total	0.74	0.90	0.48	0.96	0.64
GA: 14+0– 18+6	0.48	0.89	0.22	0.96	0.37
19+0– 23+6	0.65	0.90	0.37	0.97	0.55
24+0– 28+6	0.82	0.90	0.63	0.96	0.72
29+0– 33+6	0.83	0.89	0.65	0.96	0.72
34+0– 38+6	0.90	0.91	0.74	0.97	0.81
FL					
Total	0.65	0.90	0.44	0.95	0.55
GA: 14+0– 18+6	0.56	0.89	0.24	0.97	0.45
19+0– 23+6	0.59	0.90	0.34	0.96	0.49
24+0– 28+6	0.72	0.88	0.56	0.94	0.60
29+0– 33+6	0.70	0.90	0.62	0.93	0.60
34+0– 38+6	0.67	0.89	0.62	0.91	0.56
TCD					
Total	0.24	0.91	0.25	0.90	0.15
GA: 14+0– 18+6	0.19	0.92	0.13	0.94	0.11
19+0– 23+6	0.24	0.90	0.18	0.93	0.14
24+0– 28+6	0.30	0.91	0.39	0.86	0.21
29+0– 33+6	0.20	0.92	0.39	0.83	0.12
34+0– 38+6	0.25	0.91	0.44	0.81	0.16
FL/AC					
Total	0.40	0.78	0.19	0.91	0.18
GA: 14+0– 18+6	0.05	0.86	0.02	0.93	0.09
19+0– 23+6	0.21	0.84	0.10	0.92	0.05
24+0– 28+6	0.32	0.79	0.24	0.85	0.11
29+0– 33+6	0.60	0.58	0.26	0.86	0.18
34+0– 38+6	0.88	0.40	0.29	0.92	0.28
TCD/AC					
Total	0.23	0.94	0.33	0.90	0.17

GA: 14+0– 18+6	0.19	0.92	0.13	0.94	0.11
19+0– 23+6	0.22	0.94	0.24	0.93	0.16
24+0– 28+6	0.31	0.95	0.54	0.87	0.26
29+0– 33+6	0.24	0.97	0.64	0.84	0.21
34+0– 38+6	0.13	0.96	0.50	0.80	0.09
HC/AC					
Total	0.31	0.91	0.31	0.91	0.22
GA: 14+0– 18+6	0.19	0.90	0.11	0.94	0.09
19+0– 23+6	0.30	0.91	0.23	0.94	0.21
24+0– 28+6	0.36	0.91	0.44	0.87	0.27
29+0– 33+6	0.34	0.91	0.48	0.85	0.25
34+0– 38+6	0.27	0.94	0.55	0.82	0.21

Table S6: Evaluation of gestational age-dependent cut-off values, 3rd percentile (FGR). AC = abdominal circumference, BPD = biparietal diameter, FL = femur length, GA = Gestational age, HC = head circumference, TCD = transcerebellar diameter.

Parameter	Sensitivity	Specificity	Positive predictive value	Negative predictive value	Youden Index (J)
HC					
Total	0.64	0.97	0.39	0.99	0.61
GA: 14+0–18+6	0.63	0.96	0.10	1.00	0.59
19+0–23+6	0.53	0.97	0.19	0.99	0.50
24+0–28+6	0.77	0.96	0.56	0.98	0.73
29+0–33+6	0.60	0.97	0.63	0.97	0.57
34+0–38+6	0.67	0.97	0.68	0.97	0.64
BPD					
Total	0.45	0.98	0.40	0.98	0.43
GA: 14+0–18+6	0.38	0.97	0.08	1.00	0.35
19+0–23+6	0.37	0.98	0.19	0.99	0.35
24+0–28+6	0.59	0.97	0.58	0.97	0.56
29+0–33+6	0.41	0.99	0.78	0.96	0.40
34+0–38+6	0.42	0.98	0.71	0.95	0.40
AC					
Total	0.87	0.97	0.51	1.00	0.84
GA: 14+0–18+6	0.63	0.97	0.12	1.00	0.60
19+0–23+6	0.80	0.97	0.28	1.00	0.77
24+0–28+6	0.91	0.98	0.74	0.99	0.89
29+0–33+6	0.87	0.97	0.72	0.99	0.84
34+0–38+6	0.98	0.98	0.82	1.00	0.96
FL					
Total	0.62	0.99	0.58	0.99	0.61
GA: 14+0–18+6	0.88	0.97	0.18	1.00	0.85
19+0–23+6	0.71	0.97	0.41	1.00	0.68
24+0–28+6	0.72	0.99	0.78	0.98	0.71
29+0–33+6	0.59	0.99	0.86	0.97	0.58
34+0–38+6	0.40	1.00	1.00	0.94	0.40
TCD					
Total	0.20	0.97	0.17	0.98	0.17
GA: 14+0–18+6	0.25	0.97	0.06	0.99	0.22
19+0–23+6	0.25	0.97	0.09	0.99	0.22
24+0–28+6	0.30	0.96	0.34	0.96	0.26
29+0–33+6	0.13	0.98	0.34	0.94	0.11
34+0–38+6	0.10	0.97	0.23	0.92	0.07
FL/AC					
Total	1.00	0.01	0.03	0.99	0.01
GA: 14+0–18+6	0.88	0.09	0.01	0.99	-0.03
19+0–23+6	1.00	0.00	0.01	–	0.00
24+0–28+6	1.00	0.00	0.06	–	0.00
29+0–33+6	1.00	0.00	0.07	–	0.00
34+0–38+6	1.00	0.00	0.09	–	0.00

TCD/AC						
Total		0.30	0.97	0.24	0.98	0.27
GA: 14+0-18+6		0.13	0.97	0.03	0.99	0.10
19+0-23+6		0.32	0.97	0.13	0.99	0.29
24+0-28+6		0.34	0.97	0.40	0.96	0.31
29+0-33+6		0.28	0.97	0.46	0.95	0.25
34+0-38+6		0.25	0.98	0.43	0.93	0.23
HC/AC						
Total		0.13	0.99	0.25	0.97	0.12
GA: 14+0-18+6		0.00	0.98	0.00	0.99	-0.02
19+0-23+6		0.22	0.99	0.18	0.99	0.21
24+0-28+6		0.17	1.00	0.73	0.95	0.17
29+0-33+6		0.07	1.00	0.83	0.93	0.07
34+0-38+6		0.06	1.00	1.00	0.91	0.06