

Insecticidal Activity of Lemongrass Essential Oil as an Eco-friendly Agent Against the Black Cutworm *Agrotis ipsilon* (Lepidoptera: Noctuidae)

Moataz A. M. Moustafa ¹, Mona Awad ¹, Alia Amer ², Nancy N. Hassan ¹, El-Desoky S. Ibrahim ¹, Hayssam M. Ali ^{3,4}, Mohammad Akrami ⁵ and Mohamed Z. M. Salem ^{6,*}

¹ Department of Economic Entomology and Pesticides, Faculty of Agriculture, Cairo University, 12613 Giza, Egypt; moataz.moustafa79@gmail.com (M.M.); mona.awad2003@gmail.com (M.A.); whitehorse3050@gmail.com (N.N.H.); moat_mon@yahoo.com (E.D.S.I.)

² Medicinal and Aromatic plants Department, Horticulture Research Institute, Agricultural Research Center, Giza, Egypt (A.A.: dr_aliaamer@yahoo.com)

³ Botany and Microbiology Department, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia; hayhassan@ksu.edu.sa

⁴ Timber Trees Research Department, Sabahia Horticulture Research Station, Horticulture Research Institute, Agriculture Research Center, Alexandria 21526, Egypt

⁵ Department of Engineering, University of Exeter, Exeter EX4 4QF, UK; m.akrami@exeter.ac.uk (M.A.)

⁶ Forestry and Wood Technology Department, Faculty of Agriculture (El-Shatby), Alexandria University, Alexandria 21545, Egypt; zidan_forest@yahoo.com (M.Z.M.S.)

* Correspondence: zidan_forest@yahoo.com; mohamed-salem@alexu.edu.eg




Figure S1. GC-MS chromatogram of the identified chemical compounds in the essential oil from *Cymbopogon citratus* leaves.