

Supplementary Materials
Article

Effect of Different Light Spectrum in *Helicoverpa armigera* Larvae during HearNPV Induced Tree-Top Disease


Mandira Katuwal Bhattarai ^{1,†}, Upendra Raj Bhattarai ^{1,†}, Ji-nian Feng ^{2,*} and Dun Wang ^{1,*}
¹ State Key Laboratory of Crop Stress Biology for Arid Areas, Northwest A&F University, Yangling 712100, China; mandirakat123@gmail.com (M.K.B.); upendrarajbhattarai@gmail.com (U.R.B.)

² Department of Entomology, Northwest A&F University, Yangling 712100, China

* Correspondence: jnianf@nwsuaf.edu.cn (J.F.); wanghande@nwsuaf.edu.cn (D.W.); Tel.: +86-29-8709-1511 (D.W.)

† These authors contributed equally.

Received: 9 October 2018; Accepted: 20 November 2018; Published: date






Figure S1–S9. Climbing behavior of uninfected larvae in different light treatments. Y-axis represents the vertical position of the larvae in centimeter from the base of the cylinder; X-axis represents different hour post light treatments until pupation; Error bar represents standard error mean (SEM); S1–S4 illustrates larval climbing behavior in different light treatments from below (L: D, 12h: 12h); S5 during complete dark (L: D, 0h: 24h); and S6–S9 during different light treatments from above (L: D, 12h: 12h) as mentioned in the title of each line graph. .



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).