

## Supplementary tables

**Supplementary Table S1.** Characteristics of the patients who suffered an IE complicated by vascular embolism.

Characteristics	N = 6, n (%), or Mean ± SD
<b>Valve</b>	
Native	6 (100)
Aortic IE	3 (50)
Mitral IE	3 (50)
Ring abscess and/or severe valve leakage	3 (50)
<b>Microbiology</b>	
<i>Enterococci</i>	2 (33.3)
<i>Staphylococcus aureus</i>	2 (33.3)
<i>Streptococcus gallolyticus</i>	1 (16.7)
<i>Escherichia coli</i>	1 (16.7)
<b>Treatment</b>	
Antibiotic treatment duration (weeks)	6.0 ± 3.5
Aminoglycoside use	6 (100)
Indication for surgery	5 (83.3)
Surgery	1 (16.7)
One year mortality	3 (50)

IE: infective endocarditis.

**Supplementary Table S2.** Characteristics of the cases of IE complicated by death or graft loss (without death over the study period).

Characteristics	IE Complicated by Death	IE Complicated by Graft Loss and Not Death
	N = 8 n (%) or Mean ± SD	N = 5 n (%) or Mean ± SD
<b>Valve</b>		
Native	8 (87.5)	3 (60)
Prosthetic	1 (12.5)	2 (40)
Aortic IE	4 (50)	2 (40)
Mitral IE	4 (50)	1 (20)
Mitral and aortic IE	0 (0)	2 (40)
Ring abscess and/or severe valve leakage	4 (50)	1 (20)
<b>Microbiology</b>		
<i>Enterococci</i>	4 (50)	2 (40)
<i>Staphylococcus aureus</i>	3 (37.5)	0 (0)
<i>Escherichia coli</i>	1 (10)	(0)
<i>Staphylococcus epidermidis</i>	0 (0)	2 (40)
No documentation	0 (0)	1 (20)
<b>Treatment</b>		
Antibiotic treatment duration (weeks)	NA *	6 ± 0
Aminoglycoside use	7 (87.5)	2 (40)
Indication for surgery	6 (75)	1 (20)
Surgery	1 (12.5)	0 (0)

\* Not Applicable due to early death.