

Table S2: Morphometric data of *Ascarops strongylina* by different authors, presented as mean \pm standard deviation, in millimeters

	This study (n=10)	Dakova and Panayotova-Pencheva [19]
Host	Wild boar	Pig
Male		
Length	13.4 \pm 0.73	12
	0.336 \pm	
Width	0.005	0.33
Pharynx	0.10 \pm 0.01	0.086
Esophagus	3.65 \pm 0.12	3.2
	0.295 \pm	
Excretory pore	0.023	-
	0.256 \pm	
Nerve ring	0.012	-
	0.515 \pm	
Smaller spicule	0.117	0.575
	2.940 \pm	
Bigger spicule	0.151	2.1
Cloaca to the posterior end	0.25 \pm 0.037	0.253
Gubernacle length	0.071 \pm 0.08	0.055
Gubernacle width	0.054 \pm 0.01	0.038
Caudal ala width	0.33 \pm 0.093	0.29
Female		
	17.945 \pm	
Length	1.09	16.52 \pm 2.76
	0.375 \pm	
Width	0.033	0.357.8 \pm 0.037
Pharynx	0.11 \pm 0.008	0.083 \pm 0.008
Esophagus	3.460 \pm 0.48	3.36 \pm 0.4
	0.417 \pm	
Excretory pore	0.011	-
Nerve ring	0.288 \pm 0.03	-
Vulva to the anterior end	8.525 \pm 0.6	8.120 \pm 1.34
	0.305 \pm	
Anus to the posterior end	0.017	0.232 \pm 0.059