

Table S1. Susceptibility of the *S. aureus* isolates to antimicrobial agents

| Strain | Antimicrobial agent | | | | | | | | | | | | | | | |
|---------------|---------------------|-----|-----|----|-----|---|----|-----|----|----|---|----|-----|----|----|---|
| | FOX | CPT | CIP | CN | TEC | E | TE | TGC | DA | QD | C | FD | LZD | RD | VA | |
| B863 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B864 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B865 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |
| B870 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B904 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B928 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B929 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B937 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |
| B938 | S | S | S | S | S | R | S | S | D+ | S | S | S | S | S | S | S |
| B939 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B1014 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B1198 | S | S | S | S | S | R | S | S | D+ | S | S | S | S | S | S | S |
| B1207 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |
| B1209 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B1252 | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S | S |
| B1258 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |
| B1265 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |
| B1270 | S | S | S | S | S | R | S | S | D | S | S | S | S | S | S | S |

Note: fusidic acid (FD), cefoxitin (FOX), ceftaroline (CPT), ciprofloxacin (CIP), clindamycin (DA), chloramphenicol (C), erythromycin (E), gentamicin (CN), linezolid (LZD), quinupristin-dalfopristin (QD), rifampicin (RD), teicoplanin (TEC), tetracycline (TE), tigecycline (TGC), Vancomycin (VA). S – susceptible; R - Resistant All strains were susceptible to clindamycin but the table indicates the phenotype of inducible resistance on erythromycin resistant strains.