

Supplementary Materials:

Table S1. Microorganisms previously documented in the literature as associated with secondary infections resulting from snake bites [8,23,44,46,73,80-83].

| Aerobic Bacteria | | Anaerobic Bacteria |
|------------------------------------|---------------------------------|-----------------------------------|
| Gram-positive | Gram-negative | |
| <i>Bacillus subtilis</i> | <i>Aeromonas hydrophila</i> | <i>Bacteroides fragilis</i> |
| <i>Bacillus megaterium</i> | <i>Bordetella</i> species | <i>Clostridium perfringens</i> |
| <i>Bacillus thuringiensis</i> | <i>Citrobacter freundii</i> | <i>Clostridium sporogenes</i> |
| <i>Bacillus cereus</i> | <i>Citrobacter amalonaticus</i> | <i>Cetobacterium</i> species |
| <i>Corynebacterium diphtheriae</i> | <i>Chryseobacterium</i> species | <i>Clostridium</i> species |
| <i>Enterococcus</i> species | <i>Escherichia coli</i> | <i>Peptostreptococcus</i> species |
| <i>Lactococcus lactis</i> | <i>Enterobacter aerogenes</i> | |
| <i>Micrococcus luteus</i> | <i>Klebsiella pneumoniae</i> | |
| <i>Micrococcus glutamicus</i> | <i>Morganella morganii</i> | |
| <i>Rhodococcus</i> species | <i>Proteus vulgaris</i> | |
| <i>Staphylococcus aureus</i> | <i>Proteus mirabilis</i> | |
| <i>Strptococcus</i> species | <i>Providencia rettgeri</i> | |
| <i>Vagococcus</i> species | <i>Pseudomonas aeruginosa</i> | |
| | <i>Salmonella typhi</i> | |
| | <i>Serratia marcescens</i> | |
| | <i>Serratia liquefaciens</i> | |
| | <i>Salmonella</i> species | |
| | <i>Stenotrophomonas</i> species | |
| | <i>Shewanella</i> species | |
| | <i>Vibrio cholera</i> | |
| | <i>Yersinia enterocolitica</i> | |