

Table S1. Primers used for the determination of multilocus sequence type (MLST) in this study

Gene name	Primer name	Sequences (5' to 3' end)	Amplicon size	References	
<i>gapA</i>	gapA-F	TGAAATATGACTCCACTCACGG	662	[1]	
	gapA-R	CTTCAGAAGCGGCTTTGATGGCTT			
<i>infB</i>	infB-F	CTCGCTGCTGGACTATATTCG	462		
	infB-R	CGCTTTCAGCTCAAGAACTTC			
<i>mdh</i>	mdh-F	CCCAACTCGCTTCAGGTTTCAG	756		
	mdh-R	CCGTTTTTCCCCAGCAGCAG			
<i>pgi</i>	pgi-F	GAGAAAAACCTGCCTGTACTGCTGGC	718		
	pgi-R	CGCGCCACGCTTATAGCGGTTAAT			
<i>phoE</i>	phoE-F	ACCTACCGCAACACCGACTTCTTCGG	602		
	phoE-R	TGATCAGAACTGGTAGGTGAT			
<i>rpoB</i>	rpoB-F	GGCGAAATGGCWGAGAACCA	1075		
	rpoB-R	GAGTCTTCGAAGTTGTAACC			
<i>wzi</i>	wzi_for2	GTGCCGCGAGCGCTTTCTATCTTGGTATTCC	580		[2]
	wzi_rev	GAGAGCCACTGGTTCAGAAAYTTSACCGC			

Table S2. Primers used for the detection of carbapenemases and extended-spectrum β -lactamases in this study

Gene name	Primer name	Sequences (5' to 3' end)	Amplicon size (bp)	References	
Primers for carbapenemases					
<i>bla</i> _{IMP}	IMP-F	GGAATAGAGTGGCTTAAAYTCTC	233	[3]	
	IMP-R	GGTTTAAAYAAAACAACCACC			
<i>bla</i> _{VIM}	VIM-F	CCGATGGTGTGGTTCGC	397		
	VIM-R	CGGTCCAATGCGCAGCAC			
<i>bla</i> _{OXA-48}	OXA-F	CCCAATAGCTTGATCGCCC	466		
	OXA-R	CATCAAGTTC AACCCAACCG			
<i>bla</i> _{KPC}	KPC-Fm	CGTCTAGTTCTGCTGTCTTGTC	798		
	KPC-Rm	CTTGTCATCCTTGTTAGGCGC			
<i>bla</i> _{NDM}	NDM1-F	ATGGAATTGCCCAATATTATGC	813		[4]
	NDM1-R	TCAGCGCAGCTTGTCGGC			
Primers for TEM, SHV, CTX-M-group 1, 2, 9, 8/25, CMY, DHA					
<i>bla</i> _{TEM}	MultiTSO-T_for	CATTTCCGTGTCGCCCTTATTC	800	[5]	
	MultiTSO-T_rev	CGTTCATCCATAGTTGCCTGAC			
<i>bla</i> _{SHV}	MultiTSO-S_for	AGCCGCTTGAGCAAATTA AAC	713		
	MultiTSO-S_rev	ATCCCGCAGATAAATCACCAC			
<i>bla</i> _{CTX-M-gp1}	MultiCTXMGp1_for	TTAGGAARTGTGCCGCTGYA	688		
	MultiCTXMGp1-2_rev	CGATATCGTTGGTGGTRCCAT			
<i>bla</i> _{CTX-M-gp2}	MultiCTXMGp2_for	CGTTAACGGCACCATGAC	404		
	MultiCTXMGp1-2_rev	CGATATCGTTGGTGGTRCCAT			
<i>bla</i> _{CTX-M-gp9}	MultiCTXMGp9_for	TCAAGCCTGCCGATCTGGT	561		
	MultiCTXMGp9_rev	TGATTCTCGCCGCTGAAG			
<i>bla</i> _{CTX-M-gp8/25}	CTX-Mg8/25_for	AACRCRCAGACGCTCTAC	326		
	CTX-Mg8/25_rev	TCGAGCCGGAASGTGYAT			
<i>bla</i> _{DHA}	MultiCaseDHA_for	TGATGGCACAGCAGGATATTC	997		
	MultiCaseDHA_rev	GCTTTGACTCTTTCGGTATTCG			
Gene name	Primer name	Sequences (5' to 3' end)	Amplicon size (bp)	References	
<i>bla</i> _{CMY}	MultiCaseCIT_for	CGAAGAGGCAATGACCAGAC	538	[5]	
	MultiCaseCIT_rev	ACGGACAGGGTTAGGATAGY			

Table S3. Primers used for genes of outer membrane proteins and for the detection of gene expression levels in this study

Gene name	Primer name	Sequences (5' to 3' end)	Amplicon size (bp)	References
Outer membrane proteins				
<i>ompK35</i>	ompK35for	AACTTATTGACGGCAGTGGC	1,132	[6]
	ompK35rev	TTGGTAAACGATACCCACGG		
<i>ompK36</i>	ompK36for	GCAGTGGCATAATAAAAGGCA	1,144	
	ompK36rev	ACTGGTAAACCAGGCCCA		
Gene expression levels				
<i>acrB</i>	acrBF	CGATAACCTGATGTACATGTCC	206	[7]
	acrBR	CCGACAACCATCAGGAAGCT		
<i>oqxB</i>	oqxB1806F	GAGCGAGATCGGGATGAATAC	84	[8]
	oqxB1890R	CGGCGTGTGGTGAACGTG		
<i>ramA</i>	ramAF	ATCGTCGAGTGGATTGATGA	85	[9]
	ramAR	AGATGCCATTTCGAATACCC		
<i>rpoB</i>	rpoBF	AAGGCGAATCCAGCTTGTTTCAGC	147	[7]
	rpoBR	TGACGTTGCATGTTTCGCACCCATCA		

References

1. Diancourt, L.; Passet, V.; Verhoef, J.; Grimont, P.A.; Brisse, S. Multilocus sequence typing of *Klebsiella pneumoniae* nosocomial isolates. *J Clin Microbiol* **2005**, *43*, 4178-4182, doi:10.1128/JCM.43.8.4178-4182.2005.
2. Brisse, S.; Passet, V.; Haugaard, A.B.; Babosan, A.; Kassis-Chikhani, N.; Struve, C.; Decre, D. wzi Gene sequencing, a rapid method for determination of capsular type for *Klebsiella* strains. *J Clin Microbiol* **2013**, *51*, 4073-4078, doi:10.1128/JCM.01924-13.
3. Poirel, L.; Walsh, T.R.; Cuvillier, V.; Nordmann, P. Multiplex PCR for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis* **2011**, *70*, 119-123, doi:10.1016/j.diagmicrobio.2010.12.002.
4. Huang, T.W.; Chen, T.L.; Chen, Y.T.; Lauderdale, T.L.; Liao, T.L.; Lee, Y.T.; Chen, C.P.; Liu, Y.M.; Lin, A.C.; Chang, Y.H., et al. Copy Number Change of the NDM-1 sequence in a multidrug-resistant *Klebsiella pneumoniae* clinical isolate. *PloS one* **2013**, *8*, e62774, doi:10.1371/journal.pone.0062774.
5. Dallenne, C.; Da Costa, A.; Decre, D.; Favier, C.; Arlet, G. Development of a set of multiplex PCR assays for the detection of genes encoding important beta-lactamases in Enterobacteriaceae. *The Journal of antimicrobial chemotherapy* **2010**, *65*, 490-495, doi:10.1093/jac/dkp498.
6. Landman, D.; Bratu, S.; Quale, J. Contribution of OmpK36 to carbapenem susceptibility in KPC-producing *Klebsiella pneumoniae*. *Journal of medical microbiology* **2009**, *58*, 1303-1308, doi:10.1099/jmm.0.012575-0.
7. Doumith, M.; Ellington, M.J.; Livermore, D.M.; Woodford, N. Molecular mechanisms disrupting porin expression in ertapenem-resistant *Klebsiella* and *Enterobacter* spp. clinical isolates from the UK. *The Journal of antimicrobial chemotherapy* **2009**, *63*, 659-667, doi:10.1093/jac/dkp029.
8. Bialek-Davenet, S.; Lavigne, J.P.; Guyot, K.; Mayer, N.; Tournebize, R.; Brisse, S.; Leflon-Guibout, V.; Nicolas-Chanoine, M.H. Differential contribution of AcrAB and OqxAB efflux pumps to multidrug resistance and virulence in *Klebsiella pneumoniae*. *The Journal of antimicrobial chemotherapy* **2015**, *70*, 81-88, doi:10.1093/jac/dku340.
9. Veleba, M.; Higgins, P.G.; Gonzalez, G.; Seifert, H.; Schneiders, T. Characterization of RarA, a novel AraC family multidrug resistance regulator in *Klebsiella pneumoniae*. *Antimicrobial agents and chemotherapy* **2012**, *56*, 4450-4458, doi:10.1128/AAC.00456-12.