

Table S2: Data set for Injuries of Leopards -

Obliterate changes (Completely dissapeared markings)					
Method - Count					
ID Code	Injury No	Injury Area	Reference photo	Number of spots at Capture	Number of spots at recapture
YM 16	D 1	Near Muzzle (R)	5 B	10	0
YM 16	D 2	Upper side of face (L)	5 C	4	0
YM 01	U 2	Under the eye ®	6 B	7	0
YM 01	U 4	Forehead (F1)	6 D	15	7
YF 03	U 1	Mystacial spots (L)	6 E	2	0
YM 07	U 1	Forehead (F1)	6 G	10	4
YM 32	U 1	Forehead (F1) (F2)	6 H	15	10
YM 3	U1	Forehed F1		11	7
YF 5	U1	Forehead F1		15	12
YM 2	U1	Forehead (F1)		18	9
YM 11	U1	Forehead (F1)		14	10
YM 53	U1	Forehead (F1)		12	10
Rejig changes (Markings where the formation has changed)					
Method - Presence absence survey					
ID Code	Injury No	Injury Area	Reference photo	Rank of change in spot or rosette pattern formation at recaptur	
YM 07	D 1	Below Mystacial area M2 (R)	5A	2	
YM 27	D 1	Flank (L)	5D	2	
YM 59	D 1	Forehead (F2)	5E	2	
YM 59	D 2	Forehead (F3)	5F	2	
YM 01	U 1	Side of face (Right of Guide arch(R))	6A	1	
YM 01	U 3	Mystacial area (R)	6C	1	
YF 01	U 1	Forehead (F1)	6F	1	
YM 3	U2	Forehead (F2 & F3)		2	
YF 3	U2	Forehead (F1)		1	
YF 6	U1	Forehead (F1)		1	
YF 20	U1	Forehead (F1 & F3)		1	
YF 24	U1	Forehead (F1)		1	
YM 53	U2	Forehead (F2)		1	
Ranking guide for Rejig changes					
Rank 1 Prominance reduced					
Rank 2 Prominence has reduced					