


## Supplementary Material

# Hematite nanoparticles addition to serpentine/pyroxenes by-products of magnesite mining enrichment process for the production of refractories

Kyriaki Kalaitzidou<sup>1,\*</sup>, Evangelia Pagona<sup>1</sup>, Paraskevas Stratigopoulos<sup>1</sup>, Xanthi Ntampou<sup>1</sup>, Vasileios Zaspalis<sup>1</sup>, Anastasios Zouboulis<sup>2</sup> and Manassis Mitrakas<sup>1</sup>

- <sup>1</sup> Department of Chemical Engineering, School of Engineering, Aristotle University of Thessaloniki, Thessaloniki, Greece; evangelpg@cheng.auth.gr (E.P.), stratigop@cheng.auth.gr (P.S.), xntampou@cheng.auth.gr (X.N.), zaspalis@auth.gr (V.Z.)
  - <sup>2</sup> Department of Chemistry, School of Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece; zoubouli@chem.auth.gr (A.Z.)
- \* Correspondence: [kalaitzidou@cheng.auth.gr](mailto:kalaitzidou@cheng.auth.gr) (K.K.)



**Figure S1.** Pseudobinary phase diagram of the system  $\text{MgFe}_2\text{O}_4$ – $\text{Mg}_2\text{SiO}_4$  caption from Abruz et al., 1981 [1].

## References

1. Ambrúz, V. J.; Havlica, V.; Kanclir, E. Phase equilibria in the system  $\text{MgFe}_2\text{O}_4\text{—Mg}_2\text{SiO}_4$ . *Chem. Zvesti.* **1981**, *35*, 51–55.