

blinded to the participants' exposures/interventions ?																	
9. Was the loss to follow-up after baseline 20% or less? Were those lost to follow-up accounted for in the analysis?	No	No	No	Yes	Yes	Yes	Yes	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes	Yes	Yes
10. Did the statistical methods examine changes in outcome measures from before to after the intervention? Were statistical tests carried out that provided p values for the pre-to-post changes?	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
11. Were outcome measures of interest taken multiple times before the intervention and multiple times after the intervention (i.e., did they use an interrupted time-series design)?	Yes	Yes	Yes	No	Yes	Yes	No	No	No	No	No	Yes	No	No	No	No	Yes
12. If the intervention was conducted at a group level (e.g., a whole hospital, a community, etc.) did the statistical analysis take into account the use of individual-level data to determine effects at the group level?	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
Rating	Good	Fair	Fair	Fair	Good	Good	Fair	Fair	Good	Fair	Poor	Fair	Fair	Fair	Fair	Fair	Good