

Characterization of antioxidant and α -glucosidase inhibitory compounds of *Cratoxylum formosum* ssp. *pruniflorum* and optimization of extraction condition

Hee Won Ahn ¹, Le Nguyen Thanh ², Le Quoc Khanh ³, Se Hwan Ryu ¹, Solip Lee ¹, Sang Won Yeon ¹, Hak Hyun Lee ¹, Ayman Turk ¹, Ki Yong Lee ⁴, Bang Yeon Hwang ¹ and Mi Kyeong Lee ^{1,*}

¹College of Pharmacy, Chungbuk National University, Cheongju 28160, Republic of Korea

²Department of Medicinal Chemistry Technology, Institute of Marine Biochemistry, Vietnam
Academy of Science & Technology, Hanoi, 10000, Vietnam

³Hatinh Pharmaceutical Company (HADIPHAR), Hatinh 45000, Vietnam

⁴College of Pharmacy, Korea University, Sejong 47236, Republic of Korea




Figure S1. ¹H NMR spectrum of compound **16** (methanol-*d*₄, 400 MHz)




Figure S2. ¹³C NMR spectrum of compound **16** (methanol-*d*₄, 100 MHz)




Figure S3. HSQC spectrum of compound 16.






Figure S4. HMBC spectrum of compound 16.



Chemical Formula: $C_{28}H_{32}O_6$
 Exact Mass: 464.2199
 Molecular Weight: 464.5580
463.2124 (Calcd. for $C_{28}H_{31}O_6^-$ 463.2126)

Figure S5. ESI-TOF-MS spectrum of compound 16.




Figure S6. ¹H NMR spectrum of compound **19** (methanol-*d*₄, 400 MHz)




Figure S7. ¹³C NMR spectrum of compound **19** (methanol-*d*₄, 100 MHz)




Figure S8. HSQC spectrum of compound 19.






Figure S9. HMBC spectrum of compound 19.



Chemical Formula: $C_{19}H_{20}O_9$

Exact Mass: 392.1107

Molecular Weight: 392.3600

391.1034(calcd. for $C_{19}H_{19}O_9^-$) 391.1035)

Figure S10. ESI-TOF-MS spectrum of compound 19.