

Supplementary Table S1. Correlation of the degree of tubulointerstitial infiltrate with the urinary concentration of the cytokines and the analytical variables.

	Spearman Rho	p-Value
ITAC/CXCL11	0.574	0.001*
MIG/CXCL9	0.376	0.049*
ILB1	0.009	0.965
CXCL10	0.614	0.001*
IL6	0.342	0.075
IL17	0.398	0.041*
TNFalfa	0.318	0.099
IFN alfa	0.403	0.034*
MCP1	0.472	0.013*
EGF	0.41	0.03*
Serum creatinine	0.045	0.82
Serum CRP	0.435	0.021*
Urinary leukocyte count	0.317	0.101
Urinary erythrocyte count	0.011	0.957
Blood eosinophil count	0.25	0.199

Supplementary Table S2. Value of the cytokines at diagnosis and follow-up.

	Diagnosis sample	Follow-up sample	p-Value
I-TAC/CXCL11	44.34 ± 20.74	4.36 ± 2.05	0.002*
MIG/CXCL9	1044 ± 457.3	22.86 ± 8.67	0.007*
IL1B	5.8 ± 2.99	17.73 ± 15.89	0.82
CXCL10	441.3 ± 248.7	53.02 ± 23.83	0.02*
IL6	456 ± 408.2	18.98 ± 5.72	0.1
IL17	11.69 ± 9.86	1.48 ± 0.36	0.15
TNF alfa	8.65 ± 3.81	3.55 ± 0.16	0.08
IFN alfa	30.63 ± 26.48	2.58 ± 1.51	0.3
MCP1	14106 ± 6727	6735 ± 3877	0.27
EGF	4461 ± 1212	11727 ± 2940	0.016*
Creatinine	291.7 ± 47.67	123.8 ± 13.09	0.002*
CRP (mg/mL)	113.7 ± 42.2	5.52 ± 3.21	0.008*
Urinary leukocyte count (cells/uL)	756.3 ± 670.1	20 ± 12.7	0.4
Eosinophil blood count (/10e6)	305.5 ± 78.74	237.3 ± 58.42	0.78