

Left ventricular diastolic function following anthracycline-based chemotherapy in patients with breast cancer without previous cardiac disease – a meta-analysis

Raluca I Mincu, PhD¹; Lena Lampe¹; Amir A. Mahabadi, MD¹, Rainer Kimmig, MD², Tienush Rassaf, MD¹; Matthias Totzeck, MD^{1*}

¹Department of Cardiology and Vascular Medicine, West German Heart and Vascular Center, University Hospital Essen, Hufelandstraße 55, 45147 Essen, Germany

²Clinic for Obstetrics and Gynecology, Essen University Hospital, Essen, Germany

Supplementary material

#Corresponding author:

Dr. Matthias Totzeck

University Hospital Essen

West German Heart and Vascular Center

Department for Cardiology and Vascular Medicine

Hufelandstr. 55, 45147 Essen, Germany

Tel.: +49 201 723 84805

Fax: +49 201 723 5401

E-mail: Matthias.Totzeck@uk-essen.de

Figure S1.

PRIMA flowchart for study selection




Figure S2. Funnel plot of standard error by standard difference in means to assess the risk of bias for the analysis of E/A changes before and after anthracycline-based chemotherapy.




Figure S3. Funnel plot of standard error by standard difference in means to assess the risk of bias for the analysis of E/e' changes before and after anthracycline-based chemotherapy.




Figure S4. Funnel plot of standard error by standard difference in means to assess the risk of bias for the analysis of LVEF changes before and after anthracycline-based chemotherapy.

