

# Effectiveness of Species- and Trichothecene-Specific Primers in Monitoring *Fusarium graminearum* Species Complex in Small Grain–Pea Intercropping Systems

Vesna Župunski <sup>1\*</sup>, Radivoje Jevtić <sup>1</sup>, Milosav Grčak <sup>2</sup>, Mirjana Lalošević <sup>1</sup>, Branka Orbović <sup>1</sup>, Dalibor Živanov <sup>1</sup> and Desimir Knežević <sup>2</sup>

<sup>1</sup> Department of Small Grains, Institute of Field and Vegetable Crops, 21000 Novi Sad, Serbia; vesna.zupunski@ifvcns.ns.ac.rs (V.Ž.); radivoje.jevtic@ifvcns.ns.ac.rs (R.J.); mirjana.lalosevic@ifvcns.ns.ac.rs (M.L.); dalibor.zivanov@ifvcns.ns.ac.rs (D.Ž.); branka.orbovic@ifvcns.ns.ac.rs (B.O.)

<sup>2</sup> Department of Plant Protection, Faculty of Agriculture, University of Priština–Kosovska Mitrovica, 38219 Lešak, Serbia; milosav.grcak@pr.ac.rs (M.G.); deskooa@ptt.rs (D.K.)

\* Correspondence: vesna.zupunski@ifvcns.ns.ac.rs Tel.: +381214898214

**Table S2.** Relative frequencies of *F. graminearum* s. stricto, *F. graminearum* s. lato and unidentified (FGSC) per crop.

Crop	sensu stricto	sensu lato	unidentified	Total number of isolates per crop	Relative frequency (sensu stricto)	Relative frequency (sensu lato)	Relative frequency (unidentified)
Wheat	3	0	0	22	77.27	13.63	9.09
Wheat	7	2	1				
Wheat	4	0	0				
Wheat	3	1	1				
Rye	2	1	0	11	72.72	27.27	0
Rye	4	2	0				
Rye	1	0	0				
Rye	1	0	0				
Triticale	3	3	1	19	36.84	47.36	15.78
Triticale	2	1	1				
Triticale	2	4	0				
Triticale	0	1	1				
Oat	0	0	0	1	0	100	0
Oat	0	0	0				
Oat	0	0	0				
Oat	0	1	0				
<b>Total</b>	<b>32</b>	<b>16</b>	<b>5</b>	<b>53</b>			

The counts of *F. graminearum* s. stricto, *F. graminearum* s. lato and unidentified *Fusarium* species per each crop were divided by the total number of isolates per crop to obtain relative frequency.