

Supplementary Materials

Phytotoxicity and Accumulation of Copper-Based Nanoparticles in *Brassica* under Cadmium Stress

Shiqi Wang ^{1,2,3}, Yutong Fu ^{1,2,3}, Shunan Zheng ⁴, Yingming Xu ^{2,3} and Yuebing Sun ^{2,3,*}

¹ College of Resources and Environment, Northeast Agricultural University, Harbin 150030, China; wwp875822270@163.com (S.W.); 18845141255@163.com (Y.F.)

² Key Laboratory of Original Agro-Environmental Pollution Prevention and Control, Ministry of Agriculture and Rural Affairs (MARA), Agro-Environmental Protection Institute, Tianjin 300191, China; ymxu1999@126.com

³ Tianjin Key Laboratory of Agro-Environment and Agro-Product Safety, Agro-Environmental Protection Institute, MARA, Tianjin 300191, China;

⁴ Rural Energy & Environment Agency, MARA, Beijing 100125, China; zhengshunan1234@163.com

* Correspondence: sunyuebing2008@126.com

Table S1. The composition and concentration of Hoagland nutrient solution

TK	Composition	Concentration(mg/L)
TK1	K ₂ SO ₄	130.7
	KCl	7.5
TK2	KH ₂ PO ₄	34.0
	MgSO ₄ ·H ₂ O	160.2
TK3	Ca(NO ₃) ₂ ·4H ₂ O	472.3
	H ₃ BO ₃	0.6
TK4	CuSO ₄ ·5H ₂ O	2.5 × 10 ⁻²
	ZnSO ₄ ·7H ₂ O	0.3
	MnSO ₄ ·H ₂ O	0.2
TK5	(NH ₄) ₆ Mo ₇ O ₂₄ ·4H ₂ O	6.2 × 10 ⁻³
	FeSO ₄ ·7H ₂ O	27.8
	Na ₂ EDTA·2H ₂ O	37.2