

Bioactive Glasses Containing Strontium or Magnesium Ions to Enhance the Biological Response in Bone Regeneration

Sílvia Rodrigues Gavinho ¹, Ana Sofia Pádua ², Laura Isabel Vilas Holz ³, Isabel Sá-Nogueira ^{4,5}, Jorge Carvalho Silva ², João Paulo Borges ⁶, Manuel Almeida Valente ^{1,*} and Manuel Pedro Fernandes Graça ¹

- ¹ I3N and Physics Department, Aveiro University, 3810-193 Aveiro, Portugal; silviagavinho@ua.pt (S.R.G.); mpfg@ua.pt (M.P.F.G.)
 - ² I3N-CENIMAT and Physics Department, NOVA School of Science and Technology, Campus de Caparica, 2829-516 Caparica, Portugal; as.padua@campus.fct.unl.pt (A.S.P.); jcs@fct.unl.pt (J.C.S.)
 - ³ TEMA and Mechanical Engineering Department, Aveiro University, 3810-193 Aveiro, Portugal; lauraholz@ua.pt
 - ⁴ Associate Laboratory i4HB—Institute for Health and Bioeconomy, NOVA School of Science and Technology, NOVA University Lisbon, 2819-516 Caparica, Portugal; isn@fct.unl.pt
 - ⁵ UCIBIO—Applied Molecular Biosciences Unit, Department of Life Sciences, NOVA School of Science and Technology, NOVA University Lisbon, 2819-516 Caparica, Portugal
 - ⁶ I3N-CENIMAT and Materials Science Department, NOVA School of Science and Technology, Campus de Caparica, 2829-516 Caparica, Portugal; jpb@fct.unl.pt
- * Correspondence: mav@ua.pt




Figure S1. Disc diffusion method using (a) *Escherichia coli* K12 DSM498; (b) *Staphylococcus aureus* COL MRSA and (c) *Streptococcus mutans* DSM20523. The *S. mutans* strain used in this work displays

resistance to 10 µg of gentamicin. Discs impregnated with bi-distilled water (-) and gentamicin (10 µg) (G).