

Supplementary Materials

Fluorescence tumor-imaging using a thermo-responsive molecule with an emissive aminoquinoline derivative

Takeru Araki ¹, Yasufumi Fuchi ², Shuhei Murayama ^{3,4}, Ryoma Shiraishi ¹, Tokimi Oyama ², Mariko Aso ¹, Ichio Aoki ⁴, Shigeki Kobayashi ², Ken-ichi Yamada ¹ and Satoru Karasawa ^{2,5,*}

¹ Graduate School of Pharmaceutical Sciences, Kyushu University, 3-1-1 Maidashi, Higashi-Ku, Fukuoka 812-8582, Japan; araking417@gmail.com (T.A.); oirawawhitydaze0709@gmail.com (R.S.); aso@phar.kyushu-u.ac.jp (M.A.); kenyamada@phar.kyushu-u.ac.jp (K.Y.)

² Faculty of Pharmaceutical Sciences, Showa Pharmaceutical University, Machida, Tokyo 194-8543, Japan; fuchi@ac.shoyaku.ac.jp (Y.F.); b14030@ug.shoyaku.ac.jp (T.O.); kobayasi@ac.shoyaku.ac.jp (S.K.)

³ Department of Bioanalytical Chemistry, School of Pharmacy, Showa University, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8555, Japan; s.murayama@pharm.showa-u.ac.jp


⁴ Department of Molecular Imaging and Theranostics, National Institute of Radiological Sciences (NIRS), QST, Anagawa 4-9-1, Inage, Chiba 263-8555, Japan; iaoki.jp@gmail.com

⁵ PRESTO, Japan Science and Technology Agency, Kawaguchi 332-0012, Japan

* Correspondence: karasawa@ac.shoyaku.ac.jp; Tel.: +81-427-211-553

Table of Contents

1. Synthetic route of TFMAQ-diEg4 .	S3
2. Copies of ¹ H NMR spectrum of TFMAQ-diEg4 .	S4
3. Copies of ¹ H NMR spectrum of TFMAQ-Eg4 .	S5
4. Temperature dependence of the fluorescence spectra of TFMAQ-diEg4 .	S6
5. Illustrations showing the process of controlling the body temperature, in mice.	S7



Scheme S1. Synthetic route of the **TFMAQ-diEg4**.





Figure S1. ^1H NMR spectrum of TFMAQ-diEg4 in CDCl_3 . Arrow indicates ^1H of





Figure S2. ^1H NMR spectrum of TFMAQ-Eg4 in CDCl_3 . Arrow indicates ^1H of water.




Figure S3. Temperature dependence of the fluorescence spectra of TFMAQ-diEg4.




Figure S4. Illustrations showing the process of controlling the body temperature, in mice (a) and (b) indicate the conditions of without and with local heating around tumor tissues, respectively.