

Supplementary Files

Magnetic Halloysite Nanotubes based SERS Biosensor Enhanced with Au@Ag Core-Shell Nanotags for Bisphenol A Determination

Sen Li,^a Defu He^{a,b}, Shuning Li^{a,c}, Ruipeng Chen^{a,c}, Yuan Peng^a, Shuang Li^a, Dianpeng Han^a, Yu Wang^a, Kang Qin^a, Shuyue Ren^a, Ping Chen^{b*}, Zhixian Gao^{a*}

^a Tianjin Key Laboratory of Risk Assessment and Control Technology for Environment and Food Safety, Tianjin Institute of Environmental and Operational Medicine, Tianjin 300050, China

^b School of Food Science and Engineering, Jilin Agricultural University, Changchun 130118, China

^c State Key Laboratory of Food Nutrition and Safety, Tianjin University of Science & Technology, Tianjin 300457, China

* Corresponding Author: gaozhx@163.com; ccchenping@sina.com




Figure S1. Hysteresis curves for MNTs@AuNPs.




Figure S2. FTIR spectra of HNTs, MNTs and MNTs-NH₂.




Figure S3. Zeta potential of HNTs, MNTs and MNTs-NH₂ and MNTs@AuNPs.










Figure S4. X-ray diffraction patterns of MNTs and MNTs@AuNPs.










Figure S5. (a) X-ray photoelectron spectra of MNTs and MNTs@AuNPs. (b-g) XPS scan curves for the Au, C, N, O, Si, and Al elements in MNTs@AuNPs.




Figure S6. Comparison of Raman peak intensities at 1077 cm^{-1} for estradiol (E2), 4-tert-butylpyridine (4-tBP), 4,4'-dihydroxybiphenyl (DOD), hexenestrol (DES), and bisphenol A (BPA) by the reporter probes. The SERS intensities of the experimental group (S) and the control group (S0) were detected. Error bars indicate the standard deviation of the three replicate experiments.