

Supplementary Material

Antimicrobial resistance challenged with platinum(II) and palladium(II) complexes containing 1,10-phenanthroline and 5-amino-1,3,4- thiadiazole-2(3H)-thione in *Campylobacter jejuni*

Meiry Leandra de Lacerda¹, Daise Aparecida Rossi³, Eduarda Cristina Alves Lourenzatto³, Micaela Guidotti Takeuchi³, Wesley Almeida Souza^{1,2}, Raphael Tristão Cruvinel Silva¹, Luma Gonçalves Julio³, Wendell Guerra^{1*} and Roberta Torres de Melo^{3*}

¹Institute of Chemistry, Federal University of Uberlândia, Uberlândia, MG, Brasil

²Institute of Exact and Earth Sciences, Federal University of Mato Grosso, Pontal do Araguaia, MT, Brazil

³Laboratory of Molecular Epidemiology, Federal University of Uberlândia, Uberlândia, MG, Brasil

*Correspondence: wendell.guerra@ufu.br and roberta-melo@hotmail.com

Figure S1. ^1H NMR spectrum (400 MHz, DMSO-*d*6) of complex **II**.




Figure S2. ^{13}C NMR spectrum of **II** (100 MHz, $\text{DMSO}-d_6$) highlighting the region between 125.0 and 170.0 ppm.




Figure S3. ^{195}Pt NMR spectrum of **I**.

