



## Review

# Factors Influencing the Implementation of Antimicrobial Stewardship in Primary Care: A Narrative Review

Véronique Suttels <sup>1,\*</sup>, Mathias Van Singer <sup>1,†</sup>, Lauren Catherine Clack <sup>2</sup>, Catherine Plüss-Suard <sup>3</sup>, Anne Niquille <sup>4,5</sup>, Yolanda Mueller <sup>6,†</sup> and Noémie Boillat Blanco <sup>1,†</sup>

<sup>1</sup> Infectious Diseases Service, Lausanne University Hospital and University of Lausanne, 1011 Lausanne, Switzerland

<sup>2</sup> Institute for Implementation Science in Health Care, Medical Faculty, University of Zurich, 8006 Zürich, Switzerland

<sup>3</sup> Swiss Centre for Antibiotic Resistance, Institute for Infectious Diseases, University of Bern, 3001 Bern, Switzerland

<sup>4</sup> Center for Primary Care and Public Health (Unisanté), Pharmacy University of Lausanne, 1011 Lausanne, Switzerland

<sup>5</sup> Institute of Pharmaceutical Sciences of Western Switzerland, University of Lausanne, 1011 Lausanne, Switzerland

<sup>6</sup> Center for Primary Care and Public Health (Unisanté), Department of Family Medicine, 1011 Lausanne, Switzerland

\* Correspondence: veronique.suttels@chuv.ch

† These authors contribute equally to this work.

**Abstract:** Antimicrobial resistance (AMR) is directly driven by inappropriate use of antibiotics. Although the majority of antibiotics (an estimated 80%) are consumed in primary care settings, antimicrobial stewardship (AMS) activities in primary care remain underdeveloped and factors influencing their implementation are poorly understood. This can result in promising stewardship activities having little-to-no real-world impact. With this narrative review, we aim to identify and summarize peer-reviewed literature reporting on (1) the nature and impact of AMS interventions in primary care and (2) the individual and contextual factors influencing their implementation. Reported activities included AMS at different contextual levels (individual, collective and policy). AMS activities being often combined, it is difficult to evaluate them as stand-alone interventions. While some important individual and contextual factors were reported (difficulty to reach physicians leading to a low uptake of interventions, tight workflow of physicians requiring implementation of flexible and brief interventions and AMS as a unique opportunity to strengthen physician-patients relationship), this review identified a paucity of information in the literature about the factors that support or hinder implementation of AMS in primary care settings. In conclusion, identifying multilevel barriers and facilitators for AMS uptake is an essential step to explore before implementing primary care AMS interventions.

**Keywords:** antimicrobial stewardship; antimicrobial resistance; primary care; qualitative



**Citation:** Suttels, V.; Van Singer, M.; Clack, L.C.; Plüss-Suard, C.; Niquille, A.; Mueller, Y.; Boillat Blanco, N. Factors Influencing the Implementation of Antimicrobial Stewardship in Primary Care: A Narrative Review. *Antibiotics* **2023**, *12*, 30. <https://doi.org/10.3390/antibiotics12010030>

Academic Editor: Dino Sgarabotto

Received: 18 November 2022

Revised: 20 December 2022

Accepted: 22 December 2022

Published: 24 December 2022



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Antimicrobial resistance (AMR) is directly driven by inappropriate use of antibiotics [1,2]. Antibiotic consumption in the outpatient setting represents 80% or more of total antibiotic consumption in Europe and the United States [3–7]. Inappropriate prescribing accounts for up to fifty per cent of all antibiotic consumption in outpatient care (including unnecessary prescription or inappropriate selection, dosing and duration of treatment) [8]. Inappropriate use of antibiotics is common for the most frequent infections usually managed by primary care physicians such as upper and lower respiratory tract infections (RTI) and urinary tract infections (UTI) [9–11].

Antimicrobial stewardship (AMS) programs, defined as “coordinated interventions designed to improve and measure the appropriate use of antibiotics”, are developed to tackle the challenge of over-consumption of antibiotics [12]. AMS remains the cornerstone of the global effort to slow the spread of antimicrobial resistance and maximize the benefits of antibiotic treatment [13]. Such programs are common practice in acute care settings, such as hospitals, while AMS remains virtually absent in outpatient practice [14,15]. This setting presents a challenge to AMS efforts, in part, because of its partner heterogeneity (private practices with varying numbers of general practitioners (GPs) and specialists; public and private outpatient clinics) and important geographic spread [13]. While some AMS interventions show promising and effective results in clinical trials, there is little evidence that such interventions make their way into standard practice, leading to frustratingly low real-world impact. To bridge the gap between effectiveness and successful implementation, understanding of the individual and contextual factors affecting antibiotic prescribing and uptake of AMS activities in primary care is primordial and listed as a key priority in AMS research [16]. Understanding such factors is critical to designing implementation processes tailored to the primary care setting. Considering that the common indications for which antibiotics are prescribed are usually managed by primary care physicians, this is the focus of the current review.

With this narrative review, we aimed to summarize peer-reviewed literature reporting on (1) the nature and impact of AMS activities targeting physicians in primary care and (2) the specific behavioral and contextual factors that influence the uptake of AMS activities in primary care.

## 2. Methods

We searched for peer-reviewed literature in PubMed and Google scholar databases published prior to October 2022. We considered reviews and primary studies published since 2000 in English. Search terms included “Antimicrobial stewardship”, “AMS”, “primary care”, “qualitative”. Regarding selection criteria, we included clinical trials, reviews and meta-analyses that reported AMS interventions in primary care and nursing homes and/or observational quantitative and qualitative studies, clinical trials, reviews and meta-analyses that evaluated factors influencing the implementation of AMS in the community (primary care, nursing homes and pharmacies). We excluded studies which targeted a different setting, such as emergency department or hospital. References of included studies were also screened to identify further relevant articles meeting the selection criteria (having inclusion criteria and not having exclusion criteria) of this narrative review. For included articles, reported AMS activities were identified and categorized according to different contextual levels: (1) Individual level behavior change; (2) Collective (team, organization) level change, and (3) Structural/policy/legal level change as suggested by the Consolidated Framework for Implementation Research (CFIR) [17].

## 3. Results

In this narrative review, we discuss 30 studies reporting on factors influencing the implementation of antimicrobial stewardship in primary care. The characteristics of the studies are shown in Table 1 and a summary of the findings in Table 2.

### 3.1. Reviews on AMS Activities in Primary Care

A multitude of AMS activities targeting physicians, patients in primary care and/or the public have been reported [18]. A 2014 meta-analysis including 50 studies on AMS interventions in primary care identified that programs including communication skills training and laboratory testing were associated with a significant reduction in antimicrobial consumption. Evidence behind the impact of other stewardships activities was lower [19].

To guide implementation of the broad range of AMS intervention possibilities, the CDC proposed a framework of four core components targeting different contextual levels in 2016: commitment, action for policy, tracking & reporting and education & expertise [13].

In the following paragraphs, we will briefly review the impact of different AMS interventions and detail the factors influencing their implementation by based on the different contextual levels previously defined.

**Table 1.** Characteristics of studies on factors influencing the implementation of antimicrobial stewardship activities and categorized according to different contextual levels.

Contextual Level	Type of Intervention	First Author Publication Year	Study Design	Location	Sample Size		
Individual level change	Education of patients	O'Connor 2018 McKay R 2016	Narrative review Systematic Review	NA NA	NA NA		
	Education of GPs	Jeffs 2020 D'Hulster 2022	Qualitative study Clinical trial	Canada Belgium	23 HWs 10375 GPs		
	Communication skills	Jeffs 2020	Qualitative study	Canada	23 HWs		
		Fletcher-Lartey 2016	Bi-annual survey and qualitative study	Australia	32 GPs		
		Kumar 2003 D'Hulster 2022	Qualitative study Clinical trial	United Kingdom Belgium	40 GPs 10375 GPs		
		Lecky 2020	Qualitative study	United Kingdom	20 GPs and 29 patients		
		Delayed prescribing	Spurling 2017 Dallas 2020 Høye 2010 Ryves 2016	Systematic Review Qualitative study Qualitative study Qualitative study	NA Australia Norway England	22 HWs 33 GPs 32 GPs 32 GPs	
	Electronic clinical decision support tools	Forest 2014 Kortteisto 2012	Narrative review Qualitative study	NA Finland	NA 48 HWs		
	Biomarkers at the point-of-care: C reactive protein and procalcitonin	Jeffs 2020	Qualitative study	Canada	23 HWs		
		Martínez-González 2022	Web-based survey	Switzerland	188 GPs		
		Lecky 2020 Lopez-Vazquez 2011	Web-based survey Systematic Review	United Kingdom NA	428 GPs NA		
		Little 2019	Discussion of a clinical trial	NA	NA		
		Borek 2021	Qualitative study	England	50 HWs		
		Cals 2010	Qualitative study	Netherlands	20 GPs		
		Geis 2022	Qualitative study	Switzerland	12 GPs		
		Knusli 2022	Secondary analysis of a clinical trial	Switzerland	60 GPs		
		Collective level change	Guideline dissemination	Md Rezal 2015 Martinez-Gonzales 2020 Plate 2020 Hoorn 2019	Systematic Review Cross-sectional study Cross-sectional study Review	NA Switzerland Switzerland NA	NA 155'292 patients 163 GPs practices and 1352 patients NA
			Multifaceted intervention deployed by a large health care organization	Madaras-Kelly 2021	Post-implementation survey	USA	Unknown
	Provider feedback		Szymczak 2014	Qualitative study	USA	24 pediatricians 26 GPs and 26 pediatricians	
			Zetts 2020	Qualitative study	USA		
			Roche 2022 Laur 2021	Qualitative study Qualitative study	Ireland Canada	12 GPs 18 GPs	
	Quality circles		Elango 2018	Survey and Qualitative study	USA	31 HCWs	
Policy level change	Education of the public		None				
	Governmental strategies		Mauffrey 2016	Qualitative study	France	30 HCWs	

Abbreviations: NA: not applicable; HW: healthcare worker; GP: general practitioner.

### 3.2. Individual Level Behavior Change

#### 3.2.1. Education of Patients

Education for patients can be provided passively by the clinician with the help of leaflets and/or more actively in conjunction with education of prescribers on communication skills, which can enable shared decision-making and explore the patients' values and preferences [20]. Visual decision-making aids are also an attractive candidate for patients' education on AMS in primary care [21]. These easy-to-understand infographics on the benefits, and harms to expect of a certain treatment or screening test have already been

evaluated for cancer screening improving patients' informed decision making and reducing the number of screening tests [22].

**Table 2.** Summary of barriers and facilitators influencing the implementation of AMS activities.

Contextual Level	AMS Activity	Barriers	Facilitators
<b>Individual level behavior change</b>	<i>Education of patients</i>	Perceived patient desire to receive antibiotics	Part of GP duty Providing reassurance and a clear plan
	<i>Education of general practitioners</i>	Low participation Time pressure (linked to patient volume)	Flexible and relevant learning strategies Easy to access information, resources and reminders Creating a heightened awareness about AMR
	<i>Communication skills training</i>	Low participation Time pressure (short consultation time) Misunderstanding of depth of knowledge of the patient Phone consultations	
	<i>Delayed prescribing</i>	Loss of control over management decisions Less suitable for patients not fully understanding AB indications Heterogeneity between and within practices	High patient satisfaction Perceived as a safety net Educational and empowering to patients Improving patient-physician relationship Avoiding after-hours consultation
	<i>Electronic clinical decision support tools</i>	Interruption of workflow and additional time pressure Inflexibility of the application	High quality application Perception of content as relevant
	<i>Biomarkers at the point-of-care: C reactive protein and procalcitonin</i>	Reduced use over time Perceived as of limited clinical value	Addresses diagnostic uncertainty Helpful for unexperienced GPs “Social tool” to negotiate treatment and educate the patient
<b>Collective (team, organization) level change</b>	<i>Guideline dissemination</i>	Older GPs Concern of adverse patient outcome without ABs	Easy access
	<i>Multifaceted intervention deployed by a large health care organization</i>	Time pressure	Site champions being comfortable delivering the bundle intervention
	<i>Provider feedback</i>	Skepticism about the usefulness of audit data Complex reports to read No perceived impact on prescribing	Easy understandable visual data representation Feedback complemented by nurses education and communication skills training
	<i>Quality circles</i>		High readiness Positive group dynamic
<b>Structural/policy/legal level change</b>	<i>Education of the public</i>		Mass public campaigns coupled to GP education
	<i>Government strategies</i>	Directly targeting prescribers	Bottom-up approach and indirect interventions (e.g., local guidelines, reimbursement restrictions, restricted reporting of susceptibility tests, mass public campaigns)

Educational interventions targeting patients are a key component in AMS but there is a lack of evidence demonstrating their impact as a stand-alone intervention [23].

According to a qualitative study conducted in France, primary care prescribers believe that patients' education is part of their duty [24]. Depending on the setting, between 10 to 75% of patients reported the desire to receive antibiotic treatment [25]. This desire to receive antibiotics may affect their satisfaction, especially in the absence of information, reassurance and a clear plan from the physician [25,26]. However, the perception of patients' desire by the GP, rather than patients' actual desire, is strongly associated with inappropriate antibiotic consumption highlighting the need to also provide education to GPs [25,26]. Pharmacists also provide education for patients by counseling them about the appropriate use of antibiotics, improving adherence to antibiotic prescriptions (including delayed ones) or decreasing patient perception of the need for an antibiotic in viral illnesses when they are not indicated, such as in acute RTIs [27].

### 3.2.2. Education of General Practitioners

Several formats of educational sessions target physicians in primary care: lectures part of continuing medical education, online interventions and interprofessional team discussions [23]. In a similar way to the education of patients, stewardship activities usually include education of GPs as a reinforcement of other interventions, which makes it difficult to evaluate the impact of GPs education itself on antibiotic prescribing practices.

Recently, the nationwide implementation of an online communication skills training illustrated the difficulty of reaching high participation rate in a real-life setting. In this randomized trial, only 3% and 1% of GPs completed the first and second educational trainings, which was the main reason explaining the lack of impact of the intervention on antibiotic prescription [28].

A recent qualitative study identified factors affecting the uptake of educational interventions by GPs: first, facilitators: (1) having flexible and relevant learning strategies; (2) having easy to access information, resources and reminders; and (3) creating a heightened awareness about AMR; and, second, barriers: time pressure (mostly linked to patient volume) [15].

### 3.2.3. Communication Skills Training

Training physicians in communication skills facilitates shared decision making with patients by clarifying misperceptions and presenting available evidence behind benefits and harms of treatment. A meta-analysis showed moderate quality evidence that interventions aiming to facilitate shared decision making and help physicians communicate with patients reduce antibiotic use for acute RTI in primary care [29]. Communication skills training as a stand-alone intervention seems to reduce antibiotic consumption under trial conditions [30]. Most often though, this strategy is combined with other interventions, such as point-of-care laboratory testing. However, even when acquired, time pressure is the main barrier to the use of communication skills as GPs mention that it takes more time to convince patients they have a viral infection and do not need antibiotics rather than make a prescription [15]. Shorter consultation time results in less reassurance and information given about the lack of utility of antibiotics [15,31,32]. As stated in the previous paragraph, another important barrier to communication skills training is the uptake of training by physicians [28]. Other barriers identified in a qualitative study for the optimization of the management of UTIs in primary care via shared decision making included misunderstanding of depth of knowledge between GP and patient, miscommunication between the patient and the GP and the nature of the consultations (i.e., phone consultation) [33].

### 3.2.4. Delayed Prescribing

Delayed prescribing can be a useful strategy particularly when a patient expects to receive an antibiotic. It is a safe strategy which lead to a reduction in antibiotic use in RTI consultations even in high-risk patients compared to immediate antibiotics [34,35].



Interestingly, delayed prescribing does not affect patient satisfaction compared to immediate antibiotics, as this strategy gives both the GP and patient a sense of security [35]. Perceived as a safety net in case of diagnostic uncertainty or logistic restraints, GPs also view the delayed prescribing as educational and empowering to patients, strengthening the patient-physician relationship even [36]. It further helps patients to avoid after-hours consultation. On the other hand, it can sometimes be experienced as a loss of control over management decisions and is not suitable for patients who are not able to understand the indications for antibiotics [36,37]. GPs note that a factor supporting the use of delayed prescribing was prior experience with this strategy. GPs also mentioned the need of greater uniformity within and between practices on delayed prescribing, which could be reinforced by training, guidelines and feedback [38].

### 3.2.5. Electronic Clinical Decision Support Tools

Computer-assisted clinical decision support tools (CDSS) provide the prescribers an easy and rapid access to information and assist GPs at the point of prescribing. It can do so by creating alerts, proposing antibiotic order sets or by prompting questions to guide the choice, dosage and duration of treatment in the patient's electronic health record.

Under trial conditions and mostly for RTIs in primary care, CDSS induce a marginal to moderate reduction in antibiotic prescribing [39]. Very few interventions report pre-deployment stakeholder analyses or prescriber decision mapping to justify the intervention design [40]. Barriers to implementation are interruption of the workflow causing additional time pressure, and inflexibility of the application [41]. Besides these practicalities, the quality and perceived usefulness of the tool's content to assist local AMS also seem an essential feature [42].

### 3.2.6. Biomarkers at the Point-of-Care: C Reactive Protein and Procalcitonin

Biomarkers of inflammation are elevated in the acute phase response to tissue injury irrespective of its etiology. They can safely guide clinicians prescribing decision by ruling out severe bacterial infections. A recent Cochrane review including 12 trials concludes that point-of-care C reactive protein (POC-CRP) safely reduces antibiotic prescription among patients with acute RTI in primary care [43]. There were differences between studies regarding the CRP cut-off values applied to guide antibiotics. In some studies, the recommendation was rather vague, while, in other studies, the recommendation was based on different numeric cut-offs ( $\geq 40$  mg/L;  $>50$  mg/L;  $>60$  mg/L;  $\geq 100$  mg/L for an immediate prescription of antibiotics). A Swiss cluster-randomized study also showed a decrease in antibiotic prescription among patients with lower RTIs in primary care with the use of point-of-care procalcitonin (POC-PCT) at a cut-off of  $\geq 0.25$   $\mu\text{g/L}$  compared to usual care [44]. POC-CRP and -PCT are attractive tools as they specifically address diagnostic uncertainty, a known barrier to appropriate antibiotic prescription [15,45–47], while the risk of refraining from prescribing antibiotics (e.g., complications) is generally overestimated by the GP [48]. However, the long-lasting effect of POC-CRP was recently questioned when analyzing the impact 12 months after its implementation. Indeed, the early improvement seen with CRP disappeared with time mainly due to reduced use. The time required to perform the test might be a barrier to long-term engagement [49]. However, a qualitative study in high prescribing practices in England about the implementation of POC-CRP guided prescriptions (and delayed prescription) found that GPs deemed this strategy had a limited value as clinical tool, and was useful only in rare instances of clinical uncertainty and/or for clinicians less experienced. However, it was seen as a helpful “social tool” to negotiate treatment while maintaining GP-patient relationships or educating patients [50].

A nationwide prospective web-based survey evaluated GP attitudes towards POC-CRP. It showed that GPs would use lower CRP cut-offs to guide prescribing for more severe RTIs than for uncomplicated RTIs. Faced with intermediate CRP results in non-severe patients, GPs preferred to postpone decision on antibiotic prescription by 3 to 5 days rather than to write a delayed prescription [45]. For both POC-CRP and -PCT, GP's attitudes are

mostly positive, as they feel it allows for safe reduction of antibiotic consumption for RTIs. Reimbursement issues, the need for quality control and the negative impact on work flows were other factors affecting the adoption of POC-CRP and -PCT in primary care [51,52].

In a study analyzing factors associated with overruling of POC-PCT guidance in the setting of a clinical trial, some GPs characteristics (GP's number of years of experience [median of eight year among those who did not overrule PCT guidance versus 10 years among those who overruled PCT guidance] as well as GPs working in an urban setting) were associated with antibiotic prescription in spite of low PCT levels, highlighting the general behavioral problem of overprescription by physicians. (Knüsli 2022—unpublished data, [10]).

### 3.3. Collective (Team, Organization) Level Change

#### 3.3.1. Guideline Dissemination

National and international guidelines exist for the most common infections encountered in primary care. They are mostly developed by professional organizations (for example national infectious diseases societies) or, less often, publicly funded national institutes. The dissemination of guidelines is classically a top-down intervention and is often part of multifaceted interventions [20]. When used as a stand-alone intervention, mixed results were obtained with no effect to a modest overall decrease in antibiotic consumption accompanied by increased use of recommended antibiotics [11,53,54]. Although easily accessible guidelines are an appreciated resource by GPs and GPs are often aware of guidelines on common infections such as upper respiratory tract infections and of the preponderance of viral RTI in their setting, they tend to prescribe antibiotics because of concern of adverse patients outcome without antibiotics [55]. Data on the age of GPs are inconsistent, with some studies showing an association between older age and antibiotics prescription, others not [56]. Guidelines dissemination seems to be less effective to change prescribing of older GPs as demonstrated in the treatment of uncomplicated UTI, where the GP's age was directly associated with prescribing antibiotic therapy not recommended by guidelines (e.g., fluoroquinolones) [57].

Older GPs might distrust guidelines in favor of their own clinical impression because of their extensive expertise or might have kept more liberal prescribing practices. Additionally, impact of guidelines dissemination might be lower among older GPs because of lower knowledge of current and updated guidelines as shown in the treatment of cardiovascular diseases [57,58].

#### 3.3.2. Multifaceted Intervention Deployed by a Large Health Care Organization

The impact, effectiveness, and safety of implementing the CDC core elements [13] as a whole was assessed in a quasi-experimental controlled study focusing on uncomplicated acute RTI. The study showed a safe but modest reduction in antibiotic consumption. A post-implementation survey indicated that while site champions were comfortable delivering the bundle intervention, there were time constraints preventing them from carrying out their tasks properly [59].

#### 3.3.3. Provider Feedback

Provider feedback is a top-down intervention, which does not give extra work to physicians. Usually, such interventions include thousands of physicians. Feedback on prescription habits is usually organized at the level of groups of physician practices sharing a common electronic health record, or by health insurers based on billing data. Personalized feedback interventions are often part of multifaceted interventions (e.g., guidelines diffusion, provider education, peer comparison). Mixed results were obtained, as some studies showed no reduction in antibiotic prescribing [60], while others showed a moderate reduction in antibiotic consumption, especially when designed with insights from behavioral sciences (e.g., peer comparison or accountable justification) [61–64]. Qualitative data suggests 'deep skepticism' about the usefulness of audit data on antimicrobial pre-

scribing [65–67]. Physicians mention that some aspects of the feedback are presented in a complex manner which is difficult to process in a time-constrained environment. They would favor a visual presentation of the data [67]. In nursing homes, physicians feel that audit reports did not impact their prescribing as they were already aware of the problem. They tend to welcome broader interventions where audit is complemented by education of nurses and communication skills training [68].

#### 3.3.4. Quality Circles

Quality circles (QCs) are made up of 6 to 12 primary healthcare professionals who regularly meet to reflect on and improve their standard practice. Different types of QCs exist, e.g., with or without pharmacists. In Switzerland, the interprofessional QCs approach has shown potential to reduce antibiotic prescribing by GPs in private practice and in nursing homes [69]. These physician-pharmacist quality circles are based on open exchange of experiences, new knowledge acquisition and implementation. Regarding antibiotic use, this approach is operationalized through: (1) the analysis of antibiotic prescription data of each GP (annual proportion of patients with at least a prescription, profile of molecules prescribed, etc.) incl. benchmarking within the group and with GPs non active in such QC; (2) the dissemination of good clinical practice recommendations according to evidence-based medicine; (3) the definition of a local drug treatment consensus per common infection with conservative use of antibiotics; if antibiotics are needed, the choice is based on the selection of the spectrum of activity, adverse effects, interaction profile, local resistance and package size adapted to treatment duration; (4) the application of the consensus by the GPs involved; (5) the continuous improvement through revision of the consensus every 1 to 2 years to integrate new evidence and discussion of the effective changes in practice [70,71]. QCs can improve standard practice like prescription patterns and diagnostic habits, enhance professional development and psychological well-being in GPs. However, the results of randomized controlled trials are inconsistent and offer only limited behavioral explanations for these positive effects [72]. In the study by Klepser et al. [73] community pharmacists used rapid POC tests to guide clinical decision making as appropriate under a physician-led evidence-based protocol to treat patients with influenza and group A *Streptococcus* pharyngitis. This model pairing physicians and community pharmacists led to a more prudent use of antibiotics while providing safe and convenient care for patients. A recent GP–pharmacist collaboration showed effective implementation of delayed prescribing, educational programs or by reviewing broad-spectrum antimicrobial prescribing [74]. Readiness is a key factor for changing antibiotic prescribing in primary care. The most notable distinguishing characteristics between the high and low readiness-to-change practices were with regard to the nature and quality of group dynamics including communication, learning climate and cohesion [75]. The importance of group dynamics motivated the conception of QCs in European primary care.

### 3.4. Structural/Policy/Legal Level Change

#### 3.4.1. Education of the Public

Public campaigns are done in many countries to provide information on the appropriate use of antibiotics in outpatients. There is a wide variation in the intensity and type of campaign, some based on paper leaflets or simple internet messages to wide and expensive mass-media campaigns [76]. Most campaigns target the public and the physicians at the same time. Most often, health authorities implement these campaigns as part of their national strategy to reduce antimicrobial resistance. All campaigns focus on respiratory tract infections as they are the reason for most prescriptions of antibiotics.

Most public campaigns seemed to reduce antibiotic use, but there is a lack of evidence demonstrating their impact as a stand-alone intervention [23,77,78]. Multifaceted informational campaigns coupled with GP and pharmacist education repeated over several years appear to have the greatest effect [76,79–81].



### 3.4.2. Governmental Strategies

Many governments have national antimicrobial resistance actions plan, which can be very diverse. Seventeen government policy interventions have been described worldwide: most commonly public awareness campaign, guidelines, changing regulations around prescribing and reimbursement. Unfortunately, because of a lack of rigorous evaluations, their impact on antimicrobial use remains unclear. Most of these policies focus on changing the habits of physicians, rather than targeting other healthcare professionals or altering healthcare structures to reduce antibiotic consumption [82]. However, in Sweden, since the mid-1990s, the governments and health authorities took a bottom-up regulatory approach to the risks of AMR by establishing a program including benchmarking, locally adapted guidelines and restriction in reimbursement, complemented by public awareness campaigns with since then a gradual decrease in antibiotics consumption [83].

Stakeholders' views on different government interventions are rarely described. Notably, a qualitative study in France indicated that GPs prefer government interventions not directly targeting prescribers. GPs rather preferred indirect interventions such as increasing the unit sales price of antibiotics, the restricted reporting of susceptibility tests, or the limitation of the number of molecules available in primary care [24].

Governments should implement specific rules, funding and legislation for the antibiotic dispensing circuit, specifically of unit-dose antibiotics, in community and hospital pharmacies [84].

## 4. Discussion

Overall, this narrative review aimed to identify factors influencing the uptake of AMS activities in primary care. Indeed, several studies showed a low uptake of such activities, which jeopardizes their real-life impact on antibiotics consumption. Several individual, collective and policy AMS activities are usually combined, which makes them difficult to evaluate as a stand-alone intervention. Although there is a paucity of qualitative research on the uptake of AMS interventions in primary care, we still identified some recurrent patterns affecting the implementation of AMS activities. First, the main barrier to successful implementation of most AMS activities is the difficulty to reach primary care physicians. It is mainly due to the heterogeneity of GPs and to time constraints in their daily work, which jeopardizes the uptake of AMS activities. Quality circles seems to be a promising setting to enhance the uptake of AMS activities in primary care. Second, the tight workflow of GPs highlights the need of having flexible, easy-to-access and brief AMS activities. Third, several AMS activities are a unique opportunity of strengthening the physician-patients relationship, mainly through communication skills training and laboratory testing which reduce diagnostic uncertainty and can be used as a facilitator of implementation.

## 5. Conclusions

We identified some elements affecting the implementation of AMS activities, main barriers being the difficulty to reach primary care physicians and the tight workflow of GPs, while a facilitator being the opportunity to strengthen physician-patients relationship.

Failure to take these factors into account can result in stewardship activities resulting in little-to-no real-world impact. A thorough understanding of the individual and contextual factors that drive current behaviors and influence implementation of AMS activities is necessary to inform the systematic, tailored design of approaches to implement AMS in primary care setting.

**Author Contributions:** V.S. and M.V.S. prepared the first draft, literature research and tables. L.C.C. conceptualization and revision. A.N. pharmaceutical aspects and revision. C.P.-S. public health aspects and revision. Y.M. and N.B.B. structure and final revision. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was partly funded by the knowledge transfer activities of the National Funding Program 72 of the Swiss National Science Foundation, and is related to project grant 212429a 8 (to Y.M., L.C.C., C.P.-S., N.B.B.), the Eureka Eurostars grant [grant number E!113595 to N.B.B.] and the Leenaards Foundation academic award to N.B.B.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Goossens, H.; Ferech, M.; Vander Stichele, R.; Elseviers, M. Outpatient antibiotic use in Europe and association with resistance: A cross-national database study. *Lancet* **2005**, *365*, 579–587. [CrossRef]
- Murray, C.J.; Ikuta, K.S.; Sharara, F.; Swetschinski, L.; Aguilar, G.R.; Gray, A.; Han, C.; Bisignano, C.; Rao, P.; Wool, E.; et al. Global burden of bacterial antimicrobial resistance in 2019: A systematic analysis. *Lancet* **2022**, *399*, 629–655. [CrossRef] [PubMed]
- Suda, K.J.; Hicks, L.A.; Roberts, R.M.; Hunkler, R.J.; Matusiak, L.M.; Schumock, G.T. Antibiotic Expenditures by Medication, Class, and Healthcare Setting in the United States, 2010–2015. *Clin. Infect. Dis.* **2018**, *66*, 185–190. [CrossRef]
- Measuring Outpatient Antibiotic Prescribing. 2021. Available online: <https://www.cdc.gov/antibiotic-use/data/outpatient-prescribing/index.html> (accessed on 24 September 2021).
- Swedres—Svarm. *Sales of Antibiotics and Occurrence of Resistance in Sweden*; Swedres—Svarm: Uppsala, Sweden, 2020.
- ESPAUR Report 2019–20. Available online: [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/936199/ESPAUR\\_Report\\_2019-20.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/936199/ESPAUR_Report_2019-20.pdf) (accessed on 24 September 2021).
- Federal Office of Public Health and Federal Food Safety and Veterinary Office. *Swiss Antibiotic Resistance Report 2020. Usage of Antibiotics and Occurrence of Antibiotic Resistance in Switzerland*; Federal Office of Public Health and Federal Food Safety and Veterinary Office: Bern, Switzerland, 2020.
- Hersh, A.L.; King, L.M.; Shapiro, D.J.; Hicks, L.A.; Fleming-Dutra, K.E. Unnecessary Antibiotic Prescribing in US Ambulatory Care Settings, 2010–2015. *Clin. Infect. Dis.* **2021**, *72*, 133–137. [CrossRef]
- Fairlie, T.; Shapiro, D.J.; Hersh, A.L.; Hicks, L.A. National trends in visit rates and antibiotic prescribing for adults with acute sinusitis. *Arch. Intern. Med.* **2012**, *172*, 1513–1514. [CrossRef]
- Lhopitallier, L.; Kronenberg, A.; Meuwly, J.-Y.; Locatelli, I.; Mueller, Y.; Senn, N.; D’Acromont, V.; Boillat-Blanco, N. Procalcitonin and lung ultrasonography point-of-care testing to determine antibiotic prescription in patients with lower respiratory tract infection in primary care: Pragmatic cluster randomised trial. *BMJ* **2021**, *374*, n2132. [CrossRef] [PubMed]
- Slekovec, C.; Leroy, J.; Vernaz-Hegi, N.; Faller, J.P.; Sekri, D.; Hoen, B.; Talon, D.; Bertrand, X. Impact of a region wide antimicrobial stewardship guideline on urinary tract infection prescription patterns. *Int. J. Clin. Pharm.* **2012**, *34*, 325–329. [CrossRef]
- Fishman, N.; Society for Healthcare Epidemiology of America; Infectious Diseases Society of America. Policy statement on antimicrobial stewardship by the Society for Healthcare Epidemiology of America (SHEA), the Infectious Diseases Society of America (IDSA), and the Pediatric Infectious Diseases Society (PIDS). *Infect. Control. Hosp. Epidemiol.* **2012**, *33*, 322–327. [CrossRef]
- Sanchez, G.V.; Fleming-Dutra, K.E.; Roberts, R.M.; Hicks, L.A. Core Elements of Outpatient Antibiotic Stewardship. *Morb. Mortal. Wkly. Rep. Recomm. Rep.* **2016**, *65*, 1–12. [CrossRef] [PubMed]
- Hawes, L.; Buising, K.; Mazza, D. Antimicrobial Stewardship in General Practice: A Scoping Review of the Component Parts. *Antibiotics* **2020**, *9*, 498. [CrossRef] [PubMed]
- Jeffs, L.; McIsaac, W.; Zahradnik, M.; Senthinathan, A.; Dresser, L.; McIntyre, M.; Tannenbaum, D.; Bell, C.; Morris, A. Barriers and facilitators to the uptake of an antimicrobial stewardship program in primary care: A qualitative study. *PLoS ONE* **2020**, *15*, e0223822. [CrossRef] [PubMed]
- Charani, E.; McKee, M.; Ahmad, R.; Balasegaram, M.; Bonaconsa, C.; Merrett, G.B.; Busse, R.; Carter, V.; Castro-Sanchez, E.; Franklin, B.D.; et al. Optimising antimicrobial use in humans—Review of current evidence and an interdisciplinary consensus on key priorities for research. *Lancet Reg. Health Eur.* **2021**, *7*, 100161. [CrossRef]
- Damschroder, L.J.; Reardon, C.M.; Widerquist, M.A.O.; Lowery, J. The updated Consolidated Framework for Implementation Research based on user feedback. *Implement. Sci.* **2022**, *17*, 75. [CrossRef]
- Atkins, L.; Chadborn, T.; Bondaronek, P.; Ashiru-Oredope, D.; Beech, E.; Herd, N.; de La Morinière, V.; González-Iraizoz, M.; Hopkins, S.; McNulty, C.; et al. Content and Mechanism of Action of National Antimicrobial Stewardship Interventions on Management of Respiratory Tract Infections in Primary and Community Care. *Antibiotics* **2020**, *9*, 512. [CrossRef] [PubMed]
- Drekonja, D.; Filice, G.; Greer, N.; Olson, A.; MacDonald, R.; Rutks, I.; Wilt, T.J. *Antimicrobial Stewardship Programs in Outpatient Settings: A Systematic Review*; Department of Veterans Affairs: Washington, DC, USA, 2014. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK274571/> (accessed on 15 October 2022).

20. Tonkin-Crine, S.K.; Tan, P.S.; van Hecke, O.; Wang, K.; Roberts, N.W.; McCullough, A.; Hansen, M.P.; Butler, C.C.; Del Mar, C.B. Clinician-targeted interventions to influence antibiotic prescribing behaviour for acute respiratory infections in primary care: An overview of systematic reviews. *Cochrane Database Syst. Rev.* **2017**, *9*, Cd012252. [\[CrossRef\]](#)
21. Bern Institute of Primary Health Care. Research Projects. Antibiotic Prescription in Primary Care. Available online: [https://www.biham.unibe.ch/research/research\\_projects/antibiotic\\_prescription\\_in\\_primary\\_healthcare\\_practices/index\\_eng.html](https://www.biham.unibe.ch/research/research_projects/antibiotic_prescription_in_primary_healthcare_practices/index_eng.html) (accessed on 10 October 2022).
22. Tran, V.T.; Kisseleva-Romanova, E.; Rigal, L.; Falcoff, H. Impact of a printed decision aid on patients' intention to undergo prostate cancer screening: A multicentre, pragmatic randomised controlled trial in primary care. *Br. J. Gen. Pract.* **2015**, *65*, e295–e304. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Satterfield, J.; Miesner, A.R.; Percival, K.M. The role of education in antimicrobial stewardship. *J. Hosp. Infect.* **2020**, *105*, 130–141. [\[CrossRef\]](#)
24. Mauffrey, V.; Kivits, J.; Pulcini, C.; Boivin, J.M. Perception of acceptable antibiotic stewardship strategies in outpatient settings. *Med. Mal. Infect.* **2016**, *46*, 285–293. [\[CrossRef\]](#)
25. O'Connor, R.; O'Doherty, J.; O'Regan, A.; Dunne, C. Antibiotic use for acute respiratory tract infections (ARTI) in primary care; what factors affect prescribing and why is it important? A narrative review. *Ir. J. Med. Sci.* **2018**, *187*, 969–986. [\[CrossRef\]](#)
26. McKay, R.; Mah, A.; Law, M.R.; McGrail, K.; Patrick, D.M. Systematic Review of Factors Associated with Antibiotic Prescribing for Respiratory Tract Infections. *Antimicrob. Agents Chemother.* **2016**, *60*, 4106–4118. [\[CrossRef\]](#)
27. Blanchette, L.; Gauthier, T.; Heil, E.; Klepser, M.; Kelly, K.M.; Nailor, M.; Wei, W.; Suda, K. The essential role of pharmacists in antibiotic stewardship in outpatient care: An official position statement of the Society of Infectious Diseases Pharmacists. *J. Am. Pharm. Assoc.* **2018**, *58*, 481–484. [\[CrossRef\]](#) [\[PubMed\]](#)
28. D'Hulster, L.; Abrams, S.; Bruyndonckx, R.; Anthierens, S.; Adriaenssens, N.; Butler, C.C.; Verheij, T.; Goossens, H.; Little, P.; Coenen, S. Nationwide implementation of online communication skills training to reduce overprescribing of antibiotics: A stepped-wedge cluster randomized trial in general practice. *JAC Antimicrob. Resist.* **2022**, *4*, dlac070. [\[CrossRef\]](#) [\[PubMed\]](#)
29. Coxeter, P.; Del Mar, C.B.; McGregor, L.; Beller, E.M.; Hoffmann, T.C. Interventions to facilitate shared decision making to address antibiotic use for acute respiratory infections in primary care. *Cochrane Database Syst. Rev.* **2015**, *2015*, Cd010907. [\[CrossRef\]](#) [\[PubMed\]](#)
30. Strumann, C.; Steinhäuser, J.; Emcke, T.; Sönnichsen, A.; Goetz, K. Communication training and the prescribing pattern of antibiotic prescription in primary health care. *PLoS ONE* **2020**, *15*, e0233345. [\[CrossRef\]](#) [\[PubMed\]](#)
31. Fletcher-Lartey, S.; Yee, M.; Gaarslev, C.; Khan, R. Why do general practitioners prescribe antibiotics for upper respiratory tract infections to meet patient expectations: A mixed methods study. *BMJ Open* **2016**, *6*, e012244. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Kumar, S.; Little, P.; Britten, N. Why do general practitioners prescribe antibiotics for sore throat? Grounded theory interview study. *BMJ* **2003**, *326*, 138. [\[CrossRef\]](#) [\[PubMed\]](#)
33. Lecky, D.M.; Howdle, J.; Butler, C.C.; McNulty, C.A. Optimising management of UTIs in primary care: A qualitative study of patient and GP perspectives to inform the development of an evidence-based, shared decision-making resource. *Br. J. Gen. Pract.* **2020**, *70*, e330–e338. [\[CrossRef\]](#)
34. Stuart, B.; Hounkpatin, H.; Becque, T.; Yao, G.; Zhu, S.; Alonso-Coello, P.; Altiner, A.; Arroll, B.; Böhning, D.; Bostock, J.; et al. Delayed antibiotic prescribing for respiratory tract infections: Individual patient data meta-analysis. *BMJ* **2021**, *373*, n808. [\[CrossRef\]](#)
35. Spurling, G.K.; Del Mar, C.B.; Dooley, L.; Foxlee, R.; Farley, R. Delayed antibiotic prescriptions for respiratory infections. *Cochrane Database Syst. Rev.* **2017**, *9*, Cd004417. [\[CrossRef\]](#)
36. Dallas, A.; Davey, A.; Mulquiney, K.; Davis, J.; Glasziou, P.; Van Driel, M.; Magin, P. Delayed prescribing of antibiotics for acute respiratory infections by GP registrars: A qualitative study. *Fam. Pract.* **2020**, *37*, 406–411. [\[CrossRef\]](#)
37. Høye, S.; Frich, J.; Lindboek, M. Delayed prescribing for upper respiratory tract infections: A qualitative study of GPs' views and experiences. *Br. J. Gen. Pract.* **2010**, *60*, 907–912. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Ryves, R.; Eyles, C.; Moore, M.; McDermott, L.; Little, P.; Leydon, G.M. Understanding the delayed prescribing of antibiotics for respiratory tract infection in primary care: A qualitative analysis. *BMJ Open* **2016**, *6*, e011882. [\[CrossRef\]](#) [\[PubMed\]](#)
39. Holstiege, J.; Mathes, T.; Pieper, D. Effects of computer-aided clinical decision support systems in improving antibiotic prescribing by primary care providers: A systematic review. *J. Am. Med. Inform. Assoc.* **2015**, *22*, 236–242. [\[CrossRef\]](#) [\[PubMed\]](#)
40. Rawson, T.M.; Moore, L.S.P.; Hernandez, B.; Charani, E.; Castro-Sanchez, E.; Herrero, P.; Hayhoe, B.; Hope, W.; Georgiou, P.; Holmes, A.H. A systematic review of clinical decision support systems for antimicrobial management: Are we failing to investigate these interventions appropriately? *Clin. Microbiol. Infect.* **2017**, *23*, 524–532. [\[CrossRef\]](#) [\[PubMed\]](#)
41. Forrest, G.N.; Van Schooneveld, T.C.; Kullar, R.; Schulz, L.T.; Duong, P.; Postelnick, M. Use of electronic health records and clinical decision support systems for antimicrobial stewardship. *Clin. Infect. Dis.* **2014**, *59* (Suppl. 3), S122–S133. [\[CrossRef\]](#) [\[PubMed\]](#)
42. Kortteisto, T.; Komulainen, J.; Mäkelä, M.; Kunnamo, I.; Kaila, M. Clinical decision support must be useful, functional is not enough: A qualitative study of computer-based clinical decision support in primary care. *BMC Health Serv. Res.* **2012**, *12*, 349. [\[CrossRef\]](#) [\[PubMed\]](#)

43. Smedemark, S.A.; Aabenhus, R.; Llor, C.; Fournaise, A.; Olsen, O.; Jørgensen, K.J. Biomarkers as point-of-care tests to guide prescription of antibiotics in people with acute respiratory infections in primary care. *Cochrane Database Syst. Rev.* **2022**, *10*, Cd010130. [\[CrossRef\]](#)
44. Lhopitallier, L.; Kronenberg, A.; Meuwly, J.Y.; Locatelli, I.; Dubois, J.; Marti, J.; Mueller, Y.; Senn, N.; D'Acremont, V.; Boillat-Blanco, N. Procalcitonin and lung ultrasonography point-of-care testing to decide on antibiotic prescription in patients with lower respiratory tract infection in primary care: Protocol of a pragmatic cluster randomized trial. *BMC Pulm. Med.* **2019**, *19*, 143. [\[CrossRef\]](#)
45. Martínez-González, N.A.; Plate, A.; Jäger, L.; Senn, O.; Neuner-Jehle, S. The Role of Point-of-Care C-Reactive Protein Testing in Antibiotic Prescribing for Respiratory Tract Infections: A Survey among Swiss General Practitioners. *Antibiotics* **2022**, *11*, 543. [\[CrossRef\]](#)
46. Lecky, D.M.; Granier, S.; Allison, R.; Verlander, N.Q.; Collin, S.M.; McNulty, C.A.M. Infectious Disease and Primary Care Research-What English General Practitioners Say They Need. *Antibiotics* **2020**, *9*, 265. [\[CrossRef\]](#)
47. Lopez-Vazquez, P.; Vazquez-Lago, J.M.; Figueiras, A. Misprescription of antibiotics in primary care: A critical systematic review of its determinants. *J. Eval. Clin. Pract.* **2012**, *18*, 473–484. [\[CrossRef\]](#) [\[PubMed\]](#)
48. Petersen, I.; Johnson, A.M.; Islam, A.; Duckworth, G.; Livermore, D.M.; Hayward, A.C. Protective effect of antibiotics against serious complications of common respiratory tract infections: Retrospective cohort study with the UK General Practice Research Database. *BMJ* **2007**, *335*, 982. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Little, P.; Stuart, B.; Francis, N.; Douglas, E.; Tonkin-Crine, S.; Anthierens, S.; Cals, J.W.L.; Melbye, H.; Santer, M.; Moore, M.; et al. Antibiotic Prescribing for Acute Respiratory Tract Infections 12 Months After Communication and CRP Training: A Randomized Trial. *Ann. Fam. Med.* **2019**, *17*, 125–132. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Borek, A.J.; Campbell, A.; Dent, E.; Butler, C.C.; Holmes, A.; Moore, M.; Walker, A.S.; McLeod, M.; Tonkin-Crine, S. Implementing interventions to reduce antibiotic use: A qualitative study in high-prescribing practices. *BMC Fam. Pract.* **2021**, *22*, 25. [\[CrossRef\]](#) [\[PubMed\]](#)
51. Cals, J.W.; Chappin, F.H.; Hopstaken, R.M.; van Leeuwen, M.E.; Hood, K.; Butler, C.C.; Dinant, G.J. C-reactive protein point-of-care testing for lower respiratory tract infections: A qualitative evaluation of experiences by GPs. *Fam. Pract.* **2010**, *27*, 212–218. [\[CrossRef\]](#)
52. Geis, D.; Canova, N.; Lhopitallier, L.; Kronenberg, A.; Meuwly, J.Y.; Senn, N.; Mueller, Y.; Fasseur, F.; Boillat-Blanco, N. Qualitative exploration of the perception and acceptance of the use of procalcitonin point-of-care testing and lung ultrasonography by general practitioners to decide on antibiotic prescriptions for lower respiratory infections. 2022, *unpublished data*.
53. Venekamp, R.P.; Rovers, M.M.; Verheij, T.J.; Bonten, M.J.; Sachs, A.P. Treatment of acute rhinosinusitis: Discrepancy between guideline recommendations and clinical practice. *Fam. Pract.* **2012**, *29*, 706–712. [\[CrossRef\]](#)
54. Weiss, K.; Blais, R.; Fortin, A.; Lantin, S.; Gaudet, M. Impact of a multipronged education strategy on antibiotic prescribing in Quebec, Canada. *Clin. Infect. Dis.* **2011**, *53*, 433–439. [\[CrossRef\]](#)
55. Md Rezal, R.S.; Hassali, M.A.; Alrasheedy, A.A.; Saleem, F.; Md Yusof, F.A.; Godman, B. Physicians' knowledge, perceptions and behaviour towards antibiotic prescribing: A systematic review of the literature. *Expert Rev. Anti Infect. Ther.* **2015**, *13*, 665–680. [\[CrossRef\]](#)
56. Martínez-González, N.A.; Di Gangi, S.; Pichierri, G.; Neuner-Jehle, S.; Senn, O.; Plate, A. Time Trends and Factors Associated with Antibiotic Prescribing in Swiss Primary Care (2008 to 2020). *Antibiotics* **2020**, *9*, 837. [\[CrossRef\]](#)
57. Plate, A.; Kronenberg, A.; Risch, M.; Mueller, Y.; Di Gangi, S.; Rosemann, T.; Senn, O. Treatment of urinary tract infections in Swiss primary care: Quality and determinants of antibiotic prescribing. *BMC Fam. Pract.* **2020**, *21*, 125. [\[CrossRef\]](#) [\[PubMed\]](#)
58. Hoorn, C.; Crijns, H.; Dierick-van Daele, A.T.M.; Dekker, L.R.C. Review on Factors Influencing Physician Guideline Adherence in Cardiology. *Cardiol. Rev.* **2019**, *27*, 80–86. [\[CrossRef\]](#) [\[PubMed\]](#)
59. Madaras-Kelly, K.; Hostler, C.; Townsend, M.; Potter, E.M.; Spivak, E.S.; Hall, S.K.; Goetz, M.B.; Nevers, M.; Ying, J.; Haaland, B.; et al. Impact of Implementation of the Core Elements of Outpatient Antibiotic Stewardship Within Veterans Health Administration Emergency Departments and Primary Care Clinics on Antibiotic Prescribing and Patient Outcomes. *Clin. Infect. Dis.* **2021**, *73*, e1126–e1134. [\[CrossRef\]](#) [\[PubMed\]](#)
60. Hürlimann, D.; Limacher, A.; Schabel, M.; Zanetti, G.; Berger, C.; Mühlemann, K.; Kronenberg, A. Improvement of antibiotic prescription in outpatient care: A cluster-randomized intervention study using a sentinel surveillance network of physicians. *J. Antimicrob. Chemother.* **2015**, *70*, 602–608. [\[CrossRef\]](#)
61. Meeker, D.; Linder, J.A.; Fox, C.R.; Friedberg, M.W.; Persell, S.D.; Goldstein, N.J.; Knight, T.K.; Hay, J.W.; Doctor, J.N. Effect of Behavioral Interventions on Inappropriate Antibiotic Prescribing Among Primary Care Practices: A Randomized Clinical Trial. *JAMA* **2016**, *315*, 562–570. [\[CrossRef\]](#)
62. Gulliford, M.C.; Prevost, A.T.; Charlton, J.; Juszczak, D.; Soames, J.; McDermott, L.; Sultana, K.; Wright, M.; Fox, R.; Hay, A.D.; et al. Effectiveness and safety of electronically delivered prescribing feedback and decision support on antibiotic use for respiratory illness in primary care: REDUCE cluster randomised trial. *BMJ* **2019**, *364*, 1236. [\[CrossRef\]](#)



63. Gerber, J.S.; Prasad, P.A.; Fiks, A.G.; Localio, A.R.; Grundmeier, R.W.; Bell, L.M.; Wasserman, R.C.; Keren, R.; Zaoutis, T.E. Effect of an outpatient antimicrobial stewardship intervention on broad-spectrum antibiotic prescribing by primary care pediatricians: A randomized trial. *JAMA* **2013**, *309*, 2345–2352. [\[CrossRef\]](#)
64. Halbeisen, A.S.; Sacilotto, F.S.; Godet, R.; Signorell, P.; Sigrist, A.; Glinz, S.; Moffa, D.; Zeller, G.; Widmer, A.; Kronenberg, A.; et al. *Antibiotic Prescription Audit and Feedback to Reduce Antibiotic Consumption in Primary Care: A Nationwide Pragmatic Randomized Controlled Trial*; Department of Veterans Affairs, Veterans Health Administration, Quality Enhancement Research Initiative, Health Services Research & Development Service: Washington, DC, USA. Available online: <https://ssrn.com/abstract=4106536> (accessed on 16 October 2022).
65. Szymczak, J.E.; Feemster, K.A.; Zaoutis, T.E.; Gerber, J.S. Pediatrician perceptions of an outpatient antimicrobial stewardship intervention. *Infect. Control. Hosp. Epidemiol.* **2014**, *35* (Suppl. 3), S69–S78. [\[CrossRef\]](#)
66. Zetts, R.M.; Stoesz, A.; Garcia, A.M.; Doctor, J.N.; Gerber, J.S.; Linder, J.A.; Hyun, D.Y. Primary care physicians' attitudes and perceptions towards antibiotic resistance and outpatient antibiotic stewardship in the USA: A qualitative study. *BMJ Open* **2020**, *10*, e034983. [\[CrossRef\]](#)
67. Roche, K.F.; Morrissey, E.C.; Cunningham, J.; Molloy, G.J. The use of postal audit and feedback among Irish General Practitioners for the self—Management of antimicrobial prescribing: A qualitative study. *BMC Prim. Care* **2022**, *23*, 86. [\[CrossRef\]](#)
68. Laur, C.; Sribaskaran, T.; Simeoni, M.; Desveaux, L.; Daneman, N.; Mulhall, C.; Lam, J.; Ivers, N.M. Improving antibiotic initiation and duration prescribing among nursing home physicians using an audit and feedback intervention: A theory-informed qualitative analysis. *BMJ Open Qual.* **2021**, *10*, e001088. [\[CrossRef\]](#) [\[PubMed\]](#)
69. Plüss-Suard, C.; Niquille, A.; Héquet, D.; Krähenbühl, S.; Pichon, R.; Zanetti, G.; Bugnon, O.; Petignat, C. Decrease in Antibacterial Use and Facility-Level Variability After the Introduction of Guidelines and Implementation of Physician-Pharmacist-Nurse Quality Circles in Swiss Long-term Care Facilities. *J. Am. Med. Dir. Assoc.* **2020**, *21*, 78–83. [\[CrossRef\]](#) [\[PubMed\]](#)
70. Niquille, A.; Ruggli, M.; Buchmann, M.; Jordan, D.; Bugnon, O. The nine-year sustained cost-containment impact of swiss pilot physicians-pharmacists quality circles. *Ann. Pharmacother.* **2010**, *44*, 650–657. [\[CrossRef\]](#) [\[PubMed\]](#)
71. Mombelli, M.; Plüss-Suard, C.; Niquille, A.; Zanetti, G.; Boillat-Blanco, N. Antimicrobial stewardship in primary care setting. *Rev. Med. Suisse* **2016**, *12*, 744–748. [\[PubMed\]](#)
72. Rohrbasser, A.; Wong, G.; Mickan, S.; Harris, J. Understanding how and why quality circles improve standards of practice, enhance professional development and increase psychological well-being of general practitioners: A realist synthesis. *BMJ Open* **2022**, *12*, e058453. [\[CrossRef\]](#) [\[PubMed\]](#)
73. Klepser, M.E.; Adams, A.J.; Klepser, D.G. Antimicrobial stewardship in outpatient settings: Leveraging innovative physician-pharmacist collaborations to reduce antibiotic resistance. *Health Secur.* **2015**, *13*, 166–173. [\[CrossRef\]](#)
74. Saha, S.K.; Thursky, K.; Kong, D.C.M.; Mazza, D. A Novel GPPAS Model: Guiding the Implementation of Antimicrobial Stewardship in Primary Care Utilising Collaboration between General Practitioners and Community Pharmacists. *Antibiotics* **2022**, *11*, 1158. [\[CrossRef\]](#)
75. Elango, S.; Szymczak, J.E.; Bennett, I.M.; Beidas, R.S.; Werner, R.M. Changing Antibiotic Prescribing in a Primary Care Network: The Role of Readiness to Change and Group Dynamics in Success. *Am. J. Med. Qual.* **2018**, *33*, 154–161. [\[CrossRef\]](#)
76. Huttner, B.; Goossens, H.; Verheij, T.; Harbarth, S. Characteristics and outcomes of public campaigns aimed at improving the use of antibiotics in outpatients in high-income countries. *Lancet Infect. Dis.* **2010**, *10*, 17–31. [\[CrossRef\]](#)
77. Arnold, S.R.; Straus, S.E. Interventions to improve antibiotic prescribing practices in ambulatory care. *Cochrane Database Syst. Rev.* **2005**, *2005*, Cd003539. [\[CrossRef\]](#)
78. Cusumano, J.A.; Klinker, K.P.; Huttner, A.; Luther, M.K.; Roberts, J.A.; LaPlante, K.L. Towards precision medicine: Therapeutic drug monitoring-guided dosing of vancomycin and  $\beta$ -lactam antibiotics to maximize effectiveness and minimize toxicity. *Am. J. Health Syst. Pharm.* **2020**, *77*, 1104–1112. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Sabuncu, E.; David, J.; Bernède-Bauduin, C.; Pépin, S.; Leroy, M.; Boëlle, P.Y.; Watier, L.; Guillemot, D. Significant reduction of antibiotic use in the community after a nationwide campaign in France, 2002–2007. *PLoS Med.* **2009**, *6*, e1000084. [\[CrossRef\]](#) [\[PubMed\]](#)
80. Bauraind, I.; Lopez-Lozano, J.M.; Beyaert, A.; Marchal, J.L.; Seys, B.; Yane, F.; Hendrickx, E.; Goossens, H.; Tulkens, P.M.; Verbist, L. Association between antibiotic sales and public campaigns for their appropriate use. *JAMA* **2004**, *292*, 2468–2470. [\[PubMed\]](#)
81. Formoso, G.; Paltrinieri, B.; Marata, A.M.; Gagliotti, C.; Pan, A.; Moro, M.L.; Capelli, O.; Magrini, N. Feasibility and effectiveness of a low cost campaign on antibiotic prescribing in Italy: Community level, controlled, non-randomised trial. *BMJ* **2013**, *347*, f5391. [\[CrossRef\]](#) [\[PubMed\]](#)
82. Rogers Van Katwyk, S.; Grimshaw, J.M.; Nkangu, M.; Nagi, R.; Mendelson, M.; Taljaard, M.; Hoffman, S.J. Government policy interventions to reduce human antimicrobial use: A systematic review and evidence map. *PLoS Med.* **2019**, *16*, e1002819. [\[CrossRef\]](#) [\[PubMed\]](#)



- 
83. Hawkins, O.; Scott, A.M.; Montgomery, A.; Nicholas, B.; Mullan, J.; van Oijen, A.; Degeling, C. Comparing public attitudes, knowledge, beliefs and behaviours towards antibiotics and antimicrobial resistance in Australia, United Kingdom, and Sweden (2010–2021): A systematic review, meta-analysis, and comparative policy analysis. *PLoS ONE* **2022**, *17*, e0261917. [[CrossRef](#)] [[PubMed](#)]
  84. Costa, T.; Pimentel, A.C.; Mota-Vieira, L.; Castanha, A.C. The benefits of a unit dose system in oral antibiotics dispensing: Azorean hospital pharmacists tackling the socioeconomic problem of leftovers in Portugal. *Drugs Ther. Perspect.* **2021**, *37*, 212–221. [[CrossRef](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.