

Table S1. Minimum inhibitory concentration (MIC) values of antimicrobials assessed against 27 strains of *Alarcobacter butzleri* isolated from commercial chicken meat from São Paulo, Brazil.

Antimicrobial	Number of strains according to MIC (µg/mL)														MIC ₅₀	MIC ₉₀	%R
	≤0.015	0.03	0.06	0.12	0.25	0.5	1	2	4	8	16	32	64	>64			
Azithromycin	0	0	0	0	0	8	8	4	4	0	1	2	0	0	1.0	16	11.1
Ciprofloxacin	0	0	14	6	3	2	0	0	0	0	1	1	0	0	0.06	0.5	7.4
Erythromycin		0	0	0	0	0	2	13	10	1	0	1	0	0	2	4	3.7
Gentamicin				1	25	1	0	0	0	0	0	0	0	0	0.25	0.25	0
Tetracycline			0	0	0	0	5	13	8	0	1	0	0	0	2	4	3.7
Florfenicol		0	0	0	0	0	0	0	10	1	6	10	0	0	16	32	62.9
Nalidixic acid									0	2	3	6	16	0	64	64	59.2
Telithromycin	0	0	0	0	0	0	7	18	1	1	0	0	0	0	2	2	3.7
Clindamycin		0	0	0	0	0	0	1	5	5	8	8	0	0	16	32	77.2

*Gray cells are untested values; the sidebar indicates the breakpoints. Resistant strains were in bold. MIC₅₀ and MIC₉₀ values are those that inhibit 50% and 90% of tested strains. %R = % of resistant strains