

Brief Report

Generation of Herbicide-Resistant Soybean by Base Editing

Tao Wei ^{1,2}, Linjian Jiang ³, Xiang You ¹, Pengyu Ma ², Zhen Xi ^{2,*} and Ning Ning Wang ^{1,*} 

¹ Tianjin Key Laboratory of Protein Sciences, Department of Plant Biology and Ecology, College of Life Sciences, Nankai University, Tianjin 300071, China; weitao88888@163.com (T.W.); xiang.you@pku-iaas.edu.cn (X.Y.)

² State Key Laboratory of Elemento-Organic Chemistry, Department of Chemical Biology, National Engineering Research Center of Pesticide, College of Chemistry, Nankai University, Tianjin 300071, China; mapengyu0216@163.com

³ Key Laboratory of Pest Monitoring and Green Management, MOA, Department of Plant Pathology, College of Plant Protection, China Agricultural University, Beijing 100193, China; jianglinjian@cau.edu.cn

* Correspondence: zhenxi@nankai.edu.cn (Z.X.); wangnn@nankai.edu.cn (N.N.W.)

Simple Summary: Soybean is a prominent grain and oil crop in the world. The production of soybean is threatened by weed competition. Recently, CRISPR/Cas9-mediated base editing, which can cause irreversible nucleotide changes in target loci, has been widely used in basic plant science and breeding. However, until now, herbicide-resistant traits in soybean have not been created by the base editor. The objective of this study was to modify different AHAS homologous alleles by base editor to generate a novel AHAS-inhibiting herbicide-resistant soybean. Our results showed that the GmAHAS4 P180S mutants have apparent resistance to AHAS-inhibiting herbicides, which will be beneficial for weed control in soybean production.

Abstract: Weeds cause the largest yield loss in soybean production. The development of herbicide-resistant soybean germplasm is of great significance for weed control and yield improvement. In this study, we used the cytosine base editor (BE3) to develop novel herbicide-resistant soybean. We have successfully introduced base substitutions in *GmAHAS3* and *GmAHAS4* and obtained a heritable transgene-free soybean with homozygous P180S mutation in *GmAHAS4*. The GmAHAS4 P180S mutants have apparent resistance to chlorsulfuron, flucarbazone-sodium, and flumetsulam. In particular, the resistance to chlorsulfuron was more than 100 times that of wild type TL-1. The agronomic performance of the GmAHAS4 P180S mutants showed no significant differences to TL-1 under natural growth conditions. In addition, we developed allele-specific PCR markers for the GmAHAS4 P180S mutants, which can easily discriminate homozygous, heterozygous mutants, and wild-type plants. This study demonstrates a feasible and effective way to generate herbicide-resistant soybean by using CRISPR/Cas9-mediated base editing.

Keywords: soybean; *GmAHAS4*; herbicide-resistant; base editing



Citation: Wei, T.; Jiang, L.; You, X.; Ma, P.; Xi, Z.; Wang, N.N. Generation of Herbicide-Resistant Soybean by Base Editing. *Biology* **2023**, *12*, 741. <https://doi.org/10.3390/biology12050741>

Academic Editors: Zhaofeng Huang and Lang Pan

Received: 18 April 2023

Revised: 15 May 2023

Accepted: 17 May 2023

Published: 19 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

In fields, weeds not only compete with crops for light, growth space, and nutrients, but also spread pests and diseases and release toxic substances to reduce food production [1]; therefore, effective weed control is critical for world food security [2]. Currently, herbicides are extensively used to control weeds. However, improper use of herbicides can cause serious problems such as crop injury and the development of herbicide-resistant weeds [3]. Generating herbicide-resistant cultivars provides an effective solution [4]. On the one hand, herbicide-resistant varieties effectively reduced the risk of herbicide injury on crops; on the other hand, herbicide-resistant varieties with new mutations could introduce new herbicides to control recalcitrant weeds.

Soybean (*Glycine max* (L.) Merr.) is a prominent grain and oil crop in the world. Soybean is easily disturbed by weeds because the seeds are planted at a wide interval to

form branches and to allow the canopy to expand fully in the later stage of growth. The late canopy closure makes soybean more prone to weeds than other crops [5]. Weeds cause the largest yield loss in soybean production. On a global basis, there is evidence that 37% of soybean yields is threatened by weed competition [6]. Therefore, creating herbicide resistant soybean germplasm is of great significance for weed control and yield improvement. However, the efficiency of using traditional breeding tools to create herbicide-resistant varieties is very low. Recently, a CRISPR/Cas9-mediated base editor which could efficiently and precisely perform C to T base editing in mammalian cells has been developed [7]. This tool was quickly applied to different plant species, generating several novel traits with significant agricultural values [8].

Acetohydroxy acid synthase (AHAS), a key enzyme for the biosynthesis of branched-chain amino acids including valine, leucine, and isoleucine, is the target of many herbicides with different chemical properties [9]. These types of herbicides, including imidazolinones (IMI), pyrimidinyl thiobenzoates (PTB), sulfonaminocarbonyl triazolinones (SCT), sulfonlureas (SU), and triazolopyrimidines (TP), are widely used for weed control in crop fields, including soybean [10]. Thus, the creation of a herbicide-insensitive AHAS enzyme is of great value in agricultural application. Mutations in key amino acids of target enzymes can cause target site resistance, which is the most common mechanism leading to resistance to AHAS-inhibiting herbicides [11]. At present, in weed biotypes with field evolutionary resistance, at least 28 amino acid substitutions conferring resistance have been identified in eight conserved positions of AHAS enzymes [12]. Among them, the most common mutation site is the P197 codon (numbered according to the corresponding sequence of *A. thaliana*) [13]. Recently, the P197 site has been modified by the base editor to generate herbicide-resistant germplasm in many crops [4]. However, until now, herbicide-resistant traits have not been created by base editor in soybean.

In soybean, four homologs of AHAS have been identified, including *GmAHAS1* (Glyma04g37270), *GmAHAS2* (Glyma06g17790), *GmAHAS3* (Glyma13g31470), and *GmAHAS4* (Glyma15g07860). C to T mutations were once successfully introduced to the P178 codon of *GmAHAS1* via chemical mutagenesis [14], which conferred a moderate resistant level to sulfonlurea herbicides. However, it is unknown whether mutations at the P197 site (numbered according to the corresponding sequence of *A. thaliana*) of other homologous alleles will endow soybean resistance to herbicides. The objective of the present study was to modify different AHAS homologous alleles by base editing technology to create a novel herbicide-resistant soybean.

2. Materials and Methods

2.1. Vector Construction and Soybean Transformation

Close analysis revealed that the sequences of *GmAHAS3* and *GmAHAS4* in the P197 region are identical, and the sequence of *GmAHAS2* is different from that of *GmAHAS3* and *GmAHAS4* with only one base (Figure 1), so the P197 region of *GmAHAS2* and *GmAHAS3/4* were selected as the target sites for creating herbicide-resistant soybean by using a base editing system. These two selected target sequences (AGGTCCCCCGGCGCATGAT and AGGTCCCCCGCCGCATGAT) were cloned into pBSE901, respectively [15], in which sgRNA transcription was driven by the *Arabidopsis* U6 promoter and BE3 was driven by double 35S promoter. The *bar* gene, encoding phosphinothricin acetyltransferase (PAT) and conferring tolerance to phosphinothricin (PPT), was used as a selectable marker. These two binary vectors were then transformed into *Glycine max* (L.) cultivar Tianlong 1 (TL-1), as described previously [16].

	Pro197
<i>AtAHAS</i>	GGACAAGTCCCTCGTCGTATGATTGG
<i>GmAHAS1</i>	GGCCAGGTCCCCCGCCGGATGATCGG
<i>GmAHAS2</i>	GGCCAGGTCCCCGCGCATGATTGG
<i>GmAHAS3</i>	GGCCAGGTCCCCCGCCGCATGATCGG
<i>GmAHAS4</i>	GGCCAGGTCCCCCGCCGCATGATCGG
	<u>Target</u> <u>PAM</u>

Figure 1. The alignment of target regions of *AtAHAS* and *GmAHAS* genes. The P197 of *AtAHAS* are listed, and corresponding codons of *GmAHAS* genes are shown in red. The target sites of *GmAHAS* genes were marked with underscores.

2.2. Mutation and Transgene-Free Detection

For mutation detection, genomic DNA from the leaf tissue of T0 transgenic plants and their progenies was extracted using a modified CTAB method [17]. The target regions of *GmAHAS1-4* were amplified by PCR using gene-specific primers and the products were subjected to Sanger sequencing.

To identify transgene-free plants, the gene-specific primer pairs U6-26p-F/U6-26t-R, Cas9-F/Cas9-R, and Bar-F/Bar-R were used to amplify the sgRNA, Cas9, and Bar genes, respectively. Plants-lacking all three genes concurrently, were considered to be free of transgenes.

2.3. Off-Target Detection

Potential off-target sites, which contained 1–5 nucleotide mismatches relative to the *GmAHAS3/4* target sites, were identified by CRISPR-GE “<http://skl.scau.edu.cn/targetdesign/>” (accessed on 18 October 2020) [18]. The potential off-target sites were amplified with the gene-specific-primers (Table S1) and detected by Sanger sequencing.

2.4. Herbicide Tolerance Assay

Soybean seedlings were cultured in a greenhouse with 14 h light at 250 to 300 $\mu\text{mol photons/m}^2/\text{s}$ and 10 h dark at 25 °C with 60% humidity. Five representative herbicides from five AHAS inhibitors families, including bispyribac-sodium, imazapyr, chlorsulfuron, flucarbazone-sodium, and flumetsulam, were sprayed once at the V2 stage. The concentration gradient of each herbicide was 0, 1, 10, 100, and 1000 mg/L, respectively. Phenotypes and plant height of *GmAHAS4* P180S mutants and wild-type plants were recorded 10 days after herbicide treatment. Each treatment was repeated at least three times.

2.5. Development of Allele-Specific PCR Markers

To distinguish *GmAHAS4* P180S mutant from WT plants, two pairs of primers were designed. A common forward primer (*GmAHAS4*-P180-F) was *GmAHAS4*-specific and could differentiate between *GmAHAS4* and other homologous genes. The two base editing sites were chosen to design at the 3' end of the reverse primers. The wild type *GmAHAS4* gene was amplified by *GmAHAS4*-P180-F/*GmAHAS4*-180P-R, and the P180S mutant *GmAHAS4* gene was amplified by *GmAHAS4*-P180-F/*GmAHAS4*-180S-R.

3. Results

3.1. Targeted Base Editing of *GmAHAS* Genes

A total of 8 T0 transgenic plants with *GmAHAS2*-targeting vector and 15 T0 transgenic plants with *GmAHAS3/4*-targeting vector were obtained after screening by the PAT testing strips. Sanger sequencing results revealed that no gene editing occurred in 8 transgenic

plants with *GmAHAS2*-targeting vector, 5 out of 15 T0 transgenic plants with *GmAHAS3/4*-targeting vector exhibited editing events, among which 3 plants (Line 1, 5, and 11) were edited at the target sites of *GmAHAS3* and *GmAHAS4* simultaneously, 1 plant (Line 10) only mutated at the target site of *GmAHAS3*, and the other plant (Line 8) only mutated at the target site of *GmAHAS4* (Figure 2). Interestingly, most of these mutations are C to T, and in Line10, in addition to C to T, we also found C to A mutation. Furthermore, we found that these plants are all chimeric mutants, and the mutation levels at different sites in different plants are different. To assess the possible off-target events of the five base editing mutations, a total of nine putative off-target sites (including *GmAHAS1* and *GmAHAS2*) were predicted by CRISPR-GE. After sequencing the PCR products of the potential off-target sites, we did not detect any off-target events in these gene-edited plants (Figure S1).




Figure 2. Base editing of *GmAHAS* genes. Chromatograms of Sanger sequencing results of 5 T0 plants with editing events. Red lines indicate mutation locations.

3.2. Identification of *GmAHAS4* P180S Mutants

To test whether the base-edited mutations could be passed to the next generation efficiently, and to obtain heritable homozygous mutants, T1 progeny of five T0-edited plants were analyzed. Based on the sequencing results, only the mutation of Line 8 was stably inherited to the T1 generation, and a total of 23 wild-type plants (P180P/P180P), 18 heterozygous mutants (P180P/P180S), and 14 homozygous mutants (P180S/P180S) were obtained (Figure 3A). PCR amplification of *Cas9*, *sgRNA*, and *Bar* genes showed that a total of 15 T1 mutants lacked the exogenous T-DNA due to genetic segregation (Figure 3B). To examine the stable transmission of P180S allele, T1 homozygous P180S

mutants without transgenic components were cultured to T3 generation. Sequencing results showed that these T3 plants all contained homozygous P180S mutations (Figure S2), and PCR amplification results showed that they did not contain transgenic components (Figure S3). Meanwhile, no new editing events were detected in the target region of *GmAHAS1-3* and seven putative off-target sites of the T3 P180S mutants (Figure S4).




Figure 3. Identification of transgene-free *GmAHAS4* P180S homozygous mutants. (A) The mutation types of *GmAHAS4* in the T1 generation of line 8 according to the Sanger sequencing results. (B) Isolation of T1 plants without T-DNA insert. The presence and absence of individual genes were detected by PCR-amplification with gene-specific primers (Cas9, sgRNA, Bar).

3.3. Herbicide Tolerance of *GmAHAS4* P180S Mutants

To determine whether the base editing mutants could endow soybean with herbicide resistance, the wild-type controls (TL-1), together with the transgene-free T3 homozygous *GmAHAS4* P180S plants, were treated with different concentrations of bispyribac-sodium, imazapyr, chlorsulfuron, flucarbazone-sodium, and flumetsulam at the V2 stage. As shown in Figure 4, there was no phenotypic difference between the *GmAHAS4* P180S mutants and the wild-type controls (TL-1) without herbicide treatment. Compared with the wild-type TL-1, the *GmAHAS4* P180S mutants showed slight resistance to bispyribac-sodium and no apparent resistance to imazapyr. As for chlorsulfuron, the wild-type TL-1 cannot grow at a concentration of 1 mg/L, while the *GmAHAS4* P180S mutant can still maintain growth at 100 mg/L. As for flucarbazone-sodium, the wild-type plants died quickly at 100 mg/L, while the mutant survived at 1000 mg/L, although its growth was severely inhibited. As

for flumetsulam, both TL-1 and mutants could maintain growth at the concentration of 100 mg/L. When the concentration increased to 1000 mg/L, the wild-type TL-1 plants died rapidly, while the GmAHAS4 P180S mutant maintained growth.




Figure 4. Resistance of GmAHAS4 P180S to five AHAS-inhibiting herbicides. (A) Phenotype of GmAHAS4 P180S after herbicides treatment for 10 days. (B) Plant height of GmAHAS4 P180S after herbicides treatment for 10 days.

3.4. Agronomic Traits of *GmAHAS4* P180S Mutants

To evaluate the effect of the P180S allele on agronomic traits, the transgene-free T3 homozygous *GmAHAS4* P180S soybean were grown in the greenhouse under natural growth conditions in Tianjin, China. As shown in Figure 5, the agronomic traits, including the morphology of plant and seed, seed number per plant, seed yield per plant, and hundred-grain weight, showed no significant differences between TL-1 and the P180S plants.




Figure 5. Agronomic traits of *GmAHAS4* P180S mutants. (A) Morphology of plant (bar = 10 cm). (B) Morphology of seed. (C) Seed number per plant. (D) Seed yield per plant. (E) Hundred-grain weight.

Furthermore, we designed allele-specific PCR markers to distinguish *GmAHAS4* P180S mutants from the wild-type TL-1 plants. The upstream primer was designed according to the specific sequence of *GmAHAS4*, and two base editing sites were designed to the 3' end of the downstream primer (Figure 6A). The results showed that these primers could easily discriminate homozygous mutants, heterozygous mutants and wild-type plants (Figure 6B).




Figure 6. Allele-specific primer design (A) and PCR amplification of *GmAHAS4* P180S homozygous, heterozygous mutants and wild type plants (B).

4. Discussion

Due to its simplicity, specificity, efficiency, and versatility, the CRISPR/Cas system has quickly become the preferred tool for plant genome engineering [19–21]. The application of the CRISPR/Cas system in plant genome editing mainly includes gene knockout, gene targeted insertion or replacement, and gene expression regulation [22]. Recently, as a new gene-editing strategy, CRISPR/Cas-mediated base editing, which can cause irreversible nucleotide changes in target loci without double stranded DNA cleavage or any donor template, has been widely used in basic plant science and breeding [23,24]. For example, C to G substitution with frequency of 1.6% to 3.9% and C to T substitution with frequency of 1.4% to 11.5% were detected in rice by using base editing. In addition, C to T substitution in SLR1 (S97L) resulted in a distinct semi dwarf phenotype [25]. To facilitate weed control, herbicide-resistant germplasm has been generated via the base editor in many crops, including rice [26], wheat [2], canola [27], tomato [26], and watermelon [15]. In the present study, we used the cytosine base editor BE3 to generate a novel AHAS-inhibiting herbicide-resistant soybean mutant. Our results showed that the *GmAHAS4* P180S mutants have apparent resistance to chlorsulfuron, flucarbazone-sodium, and flumetsulam.

Since different types of AHAS-inhibiting herbicides bind to various sites of the herbicide binding domain, particular point mutation of the AHAS gene could confer tolerance to a specific range of AHAS-inhibiting herbicides [28]. Meanwhile, the herbicide resistance conferred by AHAS mutation has a dose effect [29]. In rice, the four different mutations in the P197 codon—P197A, P197S, P197Y, and P197F—exhibited different resistance patterns towards different AHAS-inhibiting herbicides. P197S and P197A showed themselves to

be more sensitive to bispyribac than P197Y and P197F [28]. In wheat, the P197S conferred higher resistance to nicosulfuron than P197A [2]. In this study, the *GmAHAS4* P180S mutants showed no apparent resistance to imazapyr, slight resistance to bispyribac–sodium, and apparent resistance to flumetsulam. The resistance to flucarbazone-sodium was more than 10 times and the resistance to chlorsulfuron was more than 100 times.

Gene editing efficiency and inheritance of genome modifications are two important considerations when using CRISPR/Cas9-mediated gene editing technology for basic research and molecular breeding [16]. Michno et al. reported that the integration and inheritance of CRISPR/Cas9 soybean lines include three patterns, namely, no transmission of mutations and transgenes, inheritance of transgenes located within the target sites but no mutations, and transmission of mutations but segregation of transgenes [30]. In CRISPR/Cas9-induced mutations, the loss of mutations and transgenes in subsequent generations seems to be common. Shi et al. reported that neither transgene nor mutations of two T0 generation soybean PDS gene editing lines were inherited by T1 progenies [17]. They found that the indel frequencies of the T0 generation were low in both strains, so the T0 plants were likely chimeric, and the mutation only existed in somatic cells, so it was not inherited into the T1 generation. In the present study, we did not obtain *GmAHAS2* gene editing mutants in T0 transgenic plants. Meanwhile, although five *GmAHAS3/4* gene editing mutants were obtained in the T0 transgenic plants, only one mutation was stably inherited to the next generation. The possible reasons for these phenomena mainly include the following aspects: (i) the genetic transformation efficiency of soybean was low [31], and the obtained transgenic plants were too few; (ii) the vector used in this study has low editing efficiency in soybean [32]; (iii) the T0 transgenic regenerated plants obtained by tissue culture were chimeric, and the mutations were only present in somatic cells [17]; thus, they cannot be inherited to the next generation. Therefore, systematic optimization of the soybean transgenic regeneration system and the gene-editing system to reduce the generation of chimeras and false-positive plants is of great significance for improving the efficiency of heritable gene editing mutation in soybean.

5. Conclusions

In summary, we have successfully introduced base substitutions in *GmAHAS3* and *GmAHAS4* by CRISPR/Cas-mediated base-editing and obtained a transgene-free herbicide-resistant soybean with homozygous P180S mutation in *GmAHAS4*. This P180S mutation has no adverse effects on agronomic traits. Moreover, we successfully designed allele-specific markers to identify P180S mutations of *GmAHAS4*, which will help breed herbicide-resistant soybean. In the future, multi-point mutation of the AHAS gene and simultaneous modification of different homologs will be beneficial to obtain soybean with multiple herbicide resistance, thus contributing to weed management in soybean production.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/biology12050741/s1>. Table S1 Primers used in this study. Figure S1 Off-target detection of 5 T0 mutants. Figure S2 Chromatograms of Sanger sequencing results of T3 generation of *GmAHAS4* P180S plants. Figure S3 PCR identification T3 generation of transgene-free *GmAHAS4* P180S plants. Figure S4 Off-target detection of T3 *GmAHAS4* P180S mutants.

Author Contributions: Conceptualization, T.W., Z.X. and N.N.W.; Methodology, T.W. and L.J.; Software, T.W.; Validation, T.W., L.J., X.Y. and P.M.; Formal Analysis, T.W., Z.X. and N.N.W.; Investigation, T.W., X.Y. and P.M.; Resources, T.W. and L.J.; Data Curation, T.W.; Writing—Original Draft Preparation, T.W.; Writing—Review and Editing, T.W., L.J., Z.X. and N.N.W.; Visualization, T.W. and L.J.; Supervision, Z.X. and N.N.W.; Project Administration, T.W., Z.X. and N.N.W.; Funding Acquisition, T.W., Z.X. and N.N.W. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by grants from the National Natural Science Foundation of China (22007050), and the Fundamental Research Funds for the Central Universities, Nankai University (63191320) to T.W., the Major S&T Projects on the Cultivation of New Varieties of Genetically Modified Organisms (Nos. 2016ZX08004005-004 and 2016ZX08010002-007) to N.N.W., and the National Key R&D Program of China (2017YFD0200501) to Z.X.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: All data used to support the findings of this study are included within the article and they are also available from the corresponding author upon request.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Colbach, N.; Chauvel, B.; Darmency, H.; Délye, C.; Corre, V.L. Choosing the best cropping systems to target pleiotropic effects when managing single-gene herbicide resistance in grass weeds. A blackgrass simulation study. *Pest Manag. Sci.* **2016**, *72*, 1910–1925. [CrossRef] [PubMed]
- Zhang, R.; Liu, J.; Chai, Z.; Chen, S.; Bai, Y.; Zong, Y.; Chen, K.; Li, J.; Jiang, L.; Gao, C. Generation of herbicide tolerance traits and a new selectable marker in wheat using base editing. *Nat. Plants* **2019**, *5*, 480–485. [CrossRef] [PubMed]
- Mackelprang, R.; Lemaux, P.G. Genetic engineering and editing of plants: An analysis of new and persisting questions. *Annu. Rev. Plant Biol.* **2020**, *71*, 659–687. [CrossRef]
- Hussain, A.; Ding, X.; Alariqi, M.; Manghwar, H.; Hui, F.; Li, Y.; Cheng, J.; Wu, C.; Cao, J.; Jin, S. Herbicide Resistance: Another Hot Agronomic Trait for Plant Genome Editing. *Plants* **2021**, *10*, 621. [CrossRef] [PubMed]
- Song, J.S.; Chung, J.H.; Lee, K.J.; Kwon, J.; Kim, J.W.; Im, J.H.; Kim, D.S. Herbicide-based weed management for soybean production in the far eastern region of Russia. *Agronomy* **2020**, *10*, 1823. [CrossRef]
- Kanatas, P.; Travlos, I.; Papastylianou, P.; Gazoulis, I.; Kakabouki, I.; Tsekoura, A. Yield, quality and weed control in soybean crop as affected by several cultural and weed management practices. *Not. Bot. Horti Agrobot.* **2020**, *48*, 329–341. [CrossRef]
- Komor, A.C.; Kim, Y.B.; Packer, M.S.; Zuris, J.A.; Liu, D.R. Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature* **2016**, *533*, 420–424. [CrossRef]
- Komatsu, A.; Ohtake, M.; Shimatani, Z.; Nishida, K. Production of herbicide-sensitive strain to prevent volunteer rice infestation using a CRISPR-Cas9 cytidine deaminase fusion. *Front. Plant Sci.* **2021**, *11*, 925. [CrossRef]
- Lonhienne, T.; Low, Y.S.; Garcia, M.D.; Croll, T.; Gao, Y.; Wang, Q.; Brillault, L.; Williams, C.M.; Fraser, J.A.; McGeary, R.P.; et al. Structures of fungal and plant acetohydroxyacid synthases. *Nature* **2020**, *586*, 317–321. [CrossRef]
- Li, Y.; Zhu, J.; Wu, H.; Liu, C.; Huang, C.; Lan, J.; Zhao, Y.; Xie, C. Precise base editing of non-allelic acetolactate synthase genes confers sulfonylurea herbicide resistance in maize. *Crop. J.* **2020**, *8*, 449–456. [CrossRef]
- Yu, Q.; Powles, S.B. Resistance to AHAS inhibitor herbicides: Current understanding. *Pest Manag. Sci.* **2014**, *70*, 1340–1350. [CrossRef] [PubMed]
- Zhao, N.; Yan, Y.; Du, L.; Zhang, X.; Liu, W.; Wang, J. Unravelling the effect of two herbicide resistance mutations on acetolactate synthase kinetics and growth traits. *J. Exp. Bot.* **2020**, *71*, 3535–3542. [CrossRef] [PubMed]
- Tranel, P.J.; Wright, T.R.; Heap, I.M. Mutations in Herbicide-Resistant Weeds to ALS Inhibitors. 2023. Available online: <http://www.weedscience.com> (accessed on 1 June 2022).
- Ghio, C.; Ramos, M.L.; Altieri, E.; Bulos, M.; Sala, C.A. Molecular characterization of Als1, an acetohydroxyacid synthase mutation conferring resistance to sulfonylurea herbicides in soybean. *Theor. Appl. Genet.* **2013**, *126*, 2957–2968. [CrossRef] [PubMed]
- Tian, S.; Jiang, L.; Cui, X.; Zhang, J.; Guo, S.; Li, M.; Zhang, H.; Ren, Y.; Gong, G.; Zong, M.; et al. Engineering herbicide-resistant watermelon variety through CRISPR/Cas9-mediated base-editing. *Plant Cell Rep.* **2018**, *37*, 1353–1356. [CrossRef]
- Zhang, P.; Du, H.; Wang, J.; Pu, Y.; Yang, C.; Yan, R.; Yang, H.; Cheng, H.; Yu, D. Multiplex CRISPR/Cas9-mediated metabolic engineering increases soya bean isoflavone content and resistance to soya bean mosaic virus. *Plant Biotechnol. J.* **2020**, *18*, 1384–1395. [CrossRef]
- Shi, Q.; Lu, M.; Tian, L. An efficient and specific CRISPR-Cas9 genome editing system targeting soybean phytoene desaturase genes. *BMC Biotechnol.* **2022**, *22*, 7.
- Wang, F.; Xu, Y.; Li, W.; Chen, Z.; Wang, J.; Fan, F.; Tao, Y.; Jiang, Y.; Zhu, Q.-H.; Yang, J. Creating a novel herbicide-tolerance OsALS allele using CRISPR/Cas9-mediated gene editing. *Crop. J.* **2021**, *9*, 305–312. [CrossRef]
- Wang, M.; Xu, Z.; Gosavi, G.; Ren, B.; Cao, Y.; Kuang, Y.; Zhou, C.; Spetz, C.; Yan, F.; Zhou, X.; et al. Targeted base editing in rice with CRISPR/ScCas9 system. *Plant Biotechnol. J.* **2020**, *18*, 1645–1647. [CrossRef]
- Li, Y.; Li, W.; Li, J. The CRISPR/Cas9 revolution continues: From base editing to prime editing in plant science. *J. Genet. Genom.* **2021**, *48*, 661–670. [CrossRef]
- Zhang, D.; Hussain, A.; Manghwar, H.; Xie, K.; Xie, S.; Zhao, S.; Larkin, R.M.; Qing, P.; Jin, S.; Ding, F. Genome editing with the CRISPR-Cas system: An art, ethics and global regulatory perspective. *Plant Biotechnol. J.* **2020**, *18*, 1651–1669. [CrossRef]

22. Zhan, X.; Lu, Y.; Zhu, J.K.; Botella, J.R. Genome editing for plant research and crop improvement. *J. Integr. Plant Biol.* **2021**, *1*, 3–33. [[CrossRef](#)] [[PubMed](#)]
23. Kuang, Y.; Li, S.; Ren, B.; Yan, F.; Spetz, C.; Li, X.; Zhou, X.; Zhou, H. Base-Editing-Mediated Artificial Evolution of OsALS1 In Planta to Develop Novel Herbicide-Tolerant Rice Germplasms. *Mol. Plant* **2020**, *13*, 565–572. [[CrossRef](#)] [[PubMed](#)]
24. Liu, L.; Kuang, Y.; Yan, F.; Li, S.; Ren, B.; Gosavi, G.; Spetz, C.; Li, X.; Wang, X.; Zhou, X.; et al. Developing a novel artificial rice germplasm for dinitroaniline herbicide resistance by base editing of OsTubA2. *Plant Biotechnol. J.* **2020**, *19*, 5–7. [[CrossRef](#)] [[PubMed](#)]
25. Lu, Y.; Zhu, J. Precise editing of a target base in the rice genome using a modified CRISPR/Cas9 system. *Mol. Plant* **2017**, *10*, 523–525. [[CrossRef](#)] [[PubMed](#)]
26. Shimatani, Z.; Kashojiya, S.; Takayama, M.; Terada, R.; Arazoe, T.; Ishii, H.; Teramura, H.; Yamamoto, T.; Komatsu, H.; Miura, K.; et al. Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion. *Nat. Biotechnol.* **2017**, *35*, 441–443. [[CrossRef](#)]
27. Wu, J.; Chen, C.; Xian, G.; Liu, D.; Lin, L.; Yin, S.; Sun, Q.; Fang, Y.; Zhang, H.; Wang, Y. Engineering herbicide-resistant oilseed rape by CRISPR/Cas9-mediated cytosine base-editing. *Plant Biotechnol. J.* **2020**, *18*, 1857–1859. [[CrossRef](#)]
28. Zhang, R.; Chen, S.; Meng, X.; Chai, Z.; Wang, D.; Yuan, Y.; Chen, K.; Jiang, L.; Li, J.; Gao, C. Generating broad-spectrum tolerance to ALS-inhibiting herbicides in rice by base editing. *Sci. China Life Sci.* **2021**, *64*, 1624–1633. [[CrossRef](#)]
29. Cheng, H.; Hao, M.; Ding, B.; Mei, D.; Wang, W.; Wang, H.; Zhou, R.; Liu, J.; Li, C.; Hu, Q. Base editing with high efficiency in allotetraploid oilseed rape by A3A-PBE base editing system. *Plant Biotechnol. J.* **2020**, *19*, 87–97. [[CrossRef](#)]
30. Michno, J.M.; Virdi, K.; Stec, A.O.; Liu, J.; Wang, X.; Xiong, Y.; Stupar, R.M. Integration, abundance, and transmission of mutations and transgenes in a series of CRISPR/Cas9 soybean lines. *BMC Biotechnol.* **2020**, *20*, 10. [[CrossRef](#)]
31. Li, Z.; Cheng, Q.; Gan, Z.; Hou, Z.; Zhang, Y.; Li, Y.; Li, H.; Nan, H.; Yang, C.; Chen, L.; et al. Multiplex CRISPR/Cas9-mediated knockout of soybean LNK2 advances flowering time. *Crop. J.* **2021**, *9*, 767–776. [[CrossRef](#)]
32. Bai, M.; Yuan, J.; Kuang, H.; Gong, P.; Li, S.; Zhang, Z.; Liu, B.; Sun, J.; Yang, M.; Yang, L.; et al. Generation of a multiplex mutagenesis population via pooled CRISPR-Cas9 in soya bean. *Plant Biotechnol. J.* **2010**, *18*, 721–731. [[CrossRef](#)] [[PubMed](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.