

Antimicrobial Activity of Novel Deep Eutectic Solvents

Noor Akbar ^{1,2}, Naveed Ahmed Khan ^{2,3,*}, Taleb Ibrahim ⁴, Mustafa Khamis ¹, Amir Sada Khan ⁴, Ahmad M. Alharbi ⁵, Hasan Alfahemi ⁶ and Ruqaiyyah Siddiqui ^{1,3}

¹ College of Arts and Sciences, American University of Sharjah, University City, Sharjah 26666, United Arab Emirates; noor@yahoo.com (N.A.); mkhamis@aus.edu (M.K.); ruqaiyyahsiddiqui35@gmail.com (R.S.)

² Department of Clinical Sciences, College of Medicine, University of Sharjah, University City, Sharjah P.O. Box 27272, United Arab Emirates

³ Department of Medical Biology, Faculty of Medicine, Istinye University, Istanbul 34010, Turkey

⁴ Department of Chemical Engineering, American University of Sharjah, Sharjah P.O. Box 26666, United Arab Emirates; taleb@yahoo.com (T.I.); amir@yahoo.com (A.S.K.)

⁵ Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, Taif University, Taif 21944, Saudi Arabia; a.alharbi@tu.edu.sa

⁶ Department of Medical Microbiology, Faculty of Medicine, Al-Baha University, P. O. Box. 1988, Al-Baha 65799, Saudi Arabia; halfahmi@bu.edu.sa

* Correspondence: naveed5438@gmail.com; Tel.: +971-6505-7722

Supplementary data.




Figure S1 DES 2.




Figure S2 DES 3.




Figure S3 DES 7




Figure S4 DES 8