

Review

Dysbiosis of Gut Microbiota from the Perspective of the Gut–Brain Axis: Role in the Provocation of Neurological Disorders

Meenakshi Kandpal ¹, Omkar Indari ¹, Budhadev Baral ¹, Shweta Jakhmola ¹, Deeksha Tiwari ¹, Vasundhra Bhandari ², Rajan Kumar Pandey ³ , Kiran Bala ⁴, Avinash Sonawane ⁵  and Hem Chandra Jha ^{1,*} 

¹ Infection Bioengineering Group, Department of Biosciences and Biomedical Engineering, Indian Institute of Technology Indore, Simrol, Indore 453552, Madhya Pradesh, India

² Department of Pharmacoinformatics, National Institute of Pharmaceutical Education and Research, Hyderabad 500037, Telengana, India

³ Department of Medical Biochemistry and Biophysics, Karolinska Institute, 17165 Stockholm, Sweden

⁴ Algal Ecotechnology & Sustainability Group, Department of Biosciences and Biomedical Engineering, Indian Institute of Technology Indore, Simrol, Indore 453552, Madhya Pradesh, India

⁵ Disease Biology & Cellular Immunology Lab, Department of Biosciences and Biomedical Engineering, Indian Institute of Technology Indore, Simrol, Indore 453552, Madhya Pradesh, India

* Correspondence: hemcjha@iiti.ac.in; Tel.: +91-99-7165-3189



Citation: Kandpal, M.; Indari, O.; Baral, B.; Jakhmola, S.; Tiwari, D.; Bhandari, V.; Pandey, R.K.; Bala, K.; Sonawane, A.; Jha, H.C. Dysbiosis of Gut Microbiota from the Perspective of the Gut–Brain Axis: Role in the Provocation of Neurological Disorders. *Metabolites* **2022**, *12*, 1064. <https://doi.org/10.3390/metabo12111064>

Academic Editors: Richa Batra and Priyanka Baloni

Received: 29 September 2022

Accepted: 1 November 2022

Published: 3 November 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The gut–brain axis is a bidirectional communication network connecting the gastrointestinal tract and central nervous system. The axis keeps track of gastrointestinal activities and integrates them to connect gut health to higher cognitive parts of the brain. Disruption in this connection may facilitate various neurological and gastrointestinal problems. Neurodegenerative diseases are characterized by the progressive dysfunction of specific populations of neurons, determining clinical presentation. Misfolded protein aggregates that cause cellular toxicity and that aid in the collapse of cellular proteostasis are a defining characteristic of neurodegenerative proteinopathies. These disorders are not only caused by changes in the neural compartment but also due to other factors of non-neural origin. Mounting data reveal that the majority of gastrointestinal (GI) physiologies and mechanics are governed by the central nervous system (CNS). Furthermore, the gut microbiota plays a critical role in the regulation and physiological function of the brain, although the mechanism involved has not yet been fully interpreted. One of the emerging explanations of the start and progression of many neurodegenerative illnesses is dysbiosis of the gut microbial makeup. The present understanding of the literature surrounding the relationship between intestinal dysbiosis and the emergence of certain neurological diseases, such as Alzheimer's disease, Parkinson's disease, Huntington's disease, and multiple sclerosis, is the main emphasis of this review. The potential entry pathway of the pathogen-associated secretions and toxins into the CNS compartment has been explored in this article at the outset of neuropathology. We have also included the possible mechanism of undelaying the synergistic effect of infections, their metabolites, and other interactions based on the current understanding.

Keywords: gut–brain axis; neurodegenerative disease; gut microbiota; gut dysbiosis; vagus nerve; neuroinflammation

1. Introduction

Microbial diversity colonizes nearly every nook and corner of the human body accessible to outer surroundings, including the skin as well as the gastrointestinal, genitourinary, and respiratory tracts. The number of microbial entities in the human body is believed to be $\sim 10^{13}$ – 10^{14} , which corresponds to a 1:1 ratio of human and microbial cells [1]. This consortium of microbial communities resides in a specific biological niche, collectively known as the “microbiota” [2]. The human microbiota includes a variety of fungi, viruses, archaeobacteria, protozoa, and, predominantly, bacteria [3]. Although some are toxic, most

microbial diversity is innocuous or even helpful to humans. These microorganisms residing inside/upon the human body play a remarkable role in maintaining the homeostasis of human health [4]. The human microbiota contributes significantly to developing the immune system and to building a physical barrier against pathogenic organisms [5]. However, microbial homeostasis perturbations also correspond to various diseased states, for example, inflammatory bowel disease (IBD) [6], Crohn's disease (CD) [7], Alzheimer's disease (AD) [8] and Parkinson's disease (PD) [9], non-alcoholic fatty liver disease [10], atopic eczema allergies [11], gastric cancer [12], etc. Therefore, it is of the utmost importance to investigate the function of the microbiota in normal physical and pathological conditions to aid the development of therapeutics against them.

Furthermore, the discovery of microbial diversity in different anatomical locations has led to the detailed study and characterization of the microorganisms inhabiting varied microenvironments in the human body [13]. The Human Microbiome Project (HMP) is one such initiative taken up by the Common Fund-NIH to explore the job of gut microbiota in human health and diseases [14]. The term "microbiome" refers to a collection of microbial populations that inhabit a specific environment, including their ecosystem, collective genome, and the surrounding environmental conditions [15]. The HMP has analyzed the microbial communities inhabiting various human body niches, such as the skin, oral cavity, gastrointestinal tract, and urogenital tract, and their role in healthy and diseased states [16].

The gut microbiota aids in a wide range of functions in human health; due to its prominent role in the neuroendocrine functions [17], metabolism [18], and immunity of an individual [19], the gastrointestinal microbiome has piqued the scientific community's curiosity in recent years [20]. The gastrointestinal tract (GIT) is one of the most heavily colonized organs. The human gut has the largest surface area and is rich in nutrients that bacteria can use as a substrate, making it an ideal location for colonization [21]. Over 70% of all microorganisms in the human body are thought to reside in the colon.

The role of the gut microbiota in regulating various neuroendocrine functions has been widely studied [22]. However, there is increasing agreement among human and animal studies that the disruption of the gut microbiota impacts brain development, neurological outcomes, and disorders, resulting in long-term behavioral changes [23]. Hence, gut dysbiosis is reported to be associated with several neurological disorders, including Alzheimer's [24], Parkinson's [25], Huntington's disease (HD) [26], and multiple sclerosis (MS) [27].

2. Composition of Gut Microbiota and Its Associated Multifarious Function in Host Health

The human gut microbiota weighs approximately 2.1 kg and contains about 50–100 times more information than the human genome, making it a distinct organ of the human body [28]. These bacteria predominantly live on the human body as symbionts, assisting the host in maintaining homeostasis in steady-state conditions. The gut microbiota makes significant contributions to (i) gut permeability; (ii) preventing pathogen colonization and invasion; (iii) facilitating nutrient metabolism; (iv) assisting in the synthesis of vitamins such as vitamin K, vitamin B complexes, and folate; (v) expediting essential intestinal epithelial roles such as absorption and secretion [29]; and (vi) regulating innate and acquired immune responses that locally (GI mucosa level) and systemically influence distant organs [30].

Firmicutes, *Bacteroidetes*, *Proteobacteria*, *Actinobacteria*, *Verrucomicrobia*, and *Fusobacteria* comprise the predominant gut microbial diversity [31]. Ninety percent of the microbial population comprises *Firmicutes* and *Bacteroidetes*. *Bacteroidetes* constitute prominent genera such as *Bacteroides*, *Parabacteroides*, and *Prevotella* [32]. *Firmicutes*, the other dominant group of bacteria, includes several genera, such as *Clostridium*, *Lactobacillus*, *Streptococcus*, *Enterococcus*, *Eubacterium*, and *Ruminococcus*, among others [33]. *Bifidobacterium* and *Collinsella* are two significant genera of the phylum *Actinobacteria*, accounting for less than 10% of the total gut microbiota. *Helicobacter* and *Escherichia* are the dominant genera of *Proteobacteria* (less than 2%). *Fusobacterium* and *Akkermansia* dominate the genera of the phyla *Fusobacteria* and *Verrucomicrobia*, respectively, and constitute less than 3% of total gut microbial diversity [34].

2.1. Metabolic Function—Nutrient and Other Dietary Component Metabolism

The gut microbiota has been linked to various GIT activities, notably nutritional needs, energy storage, and diverse physiological pathways [35]. Dietary carbohydrates impart a significant portion of the microbiota's nutrition. Short-chain fatty acids (SCFAs) are produced in the colon by bacteria such as *Bacteroides*, *Bifidobacterium*, *Roseburia*, *Faecalibacterium*, and *Enterobacteria* [36]. SCFAs such as butyrate, propionate, and acetate provide tremendous energy to hosts [37]. *Bacteroides* species are the most predominant organisms engaged in carbohydrate metabolism, as they encode for specific classes of enzymes, such as glycosyltransferases, glycoside hydrolases, and polysaccharide lyases, among others (Figure 1). For example, *Bacteroides thetaiotaomicron* has a genome that codes for over 260 hydrolases, significantly more than the human genome [38].

Bacteroides have been demonstrated to synthesize conjugated linoleic acid, a fatty acid possessing anti-diabetic, hypolipidemic, anti-obesogenic, immunomodulatory, and anti-atherogenic properties. A study published by the Velagapudi group showed that gut microbiota performed a crucial role in the regulation of lipogenesis when compared to conventionally raised mice (CONV-R) and germ-free mice (which lack a whole microbial community) [39]. The study reported higher levels of microbial-derived metabolites (citric acid, propionic acid, and fumaric acid) and lowered triglycerides and cholesterol levels in the serum metabolome of CONV-R mice compared to the metabolome of germ-free mice, thus demonstrating the effect of gut microbial diversity on host energy and lipid metabolism [40] (Figure 1). Furthermore, the gut microbiota has been associated with a reduced risk of renal oxalate stone formation. Because the production of oxalate in the intestine is a byproduct of bacterial metabolism and carbohydrate fermentation, it is quelled by organisms such as *Lactobacillus* species, *Bifidobacterium* species, and *Oxalobacter formigenes* [41].

The gut microbiota has been considered a cornerstone of maintaining the health status of its human host because it not only facilitates the harvesting of nutrients and energy from ingested food, but also produces numerous metabolites that can regulate host metabolism [42]. One such class of metabolites is the bile acids. Primary bile acids (BAs), such as cholic acid (CA) and chenodeoxycholic acid (CDCA), which are by-products of cholesterol metabolism and clearance, are synthesized by the liver [43] and are subsequently modified by the gut microbiota. The gut microbiota transform these compounds into an assortment of forms that significantly expand their biological value and diversity [44]. Primary bile acids have been reported to be modified by the gut microbiota in four distinct ways, including via the deconjugation of taurine or glycine amino acids as well as via the dehydrogenation, dihydroxylation, and epimerization of the cholesterol core [45]. A novel class of "microbially conjugated bile acids", which have the ability to conjugate amino acids to bile acid, is synthesized by our gut bacteria in addition to the previously reported alterations [46].




Figure 1. Summary of the gut microbiota-associated functioning in the human body. The gut microbiota leads to enhanced lipid clearance by repressing angiopoietin-like protein 4 (Angptl4), an inhibitor of LPL, due to which energy metabolites are elevated in the serum, which is demonstrated by an increase in associated genes in the liver transcriptome. Triglyceride levels in the serum are reduced while they rise in adipose tissue and in the liver. Dietary carbohydrates are metabolized into SCFAs, a rich energy source for the host by colonic bacteria by employing a special class of enzyme, hydrolases. Furthermore, these SCFAs, especially butyrate, can provoke the GPCR 43 expressed by intestinal epithelial cells and regulate the development, differentiation, and maturation of Treg cells via epigenetic regulation, resulting in the inhibition of Th17 cell development and the reduction of colonic inflammation. Selected gut microbiota can also act as vitamin suppliers to the host, as they synthesize vitamin B complexes and vitamin K and supplies them to the host.

2.2. Vitamins Supplier—Bacteria as a Source of Vitamins for Their Hosts

Among numerous associated functions, one of the beneficial activities of microorganisms is the production of multiple vitamins [47]. Another critical metabolic activity of the gut microbiota is the biosynthesis of vitamin K and various components of vitamin B

(Figure 1). *Lactobacilli* appear to be unable to synthesize folate de novo, but some *Bifidobacterial* species belonging to the phylum *Actinobacteria* can synthesize folate [48]. *Firmicutes* and *Proteobacteria* species, *Bacillus subtilis*, and *Escherichia coli* can produce riboflavin [49]. Vitamin B12 is commercially produced utilizing *Propionibacterium freudenreichii*, a member of the phylum *Actinobacteria* [50]. Another *Firmicutes* phylum member, *Lactobacillus reuteri*, is thought to possess the potential for vitamin B12 production [51].

2.3. Immunomodulatory Role of Gut Microbiota

The gut microbiota helps to influence both the innate and adaptive immune systems in the gut [52]. The effector and regulatory T cells (Tregs), gut-associated lymphoid tissues (GALT), group 3 innate lymphoid cells, IgA-producing B (plasma) cells, dendritic cells in the lamina propria, and resident macrophages are immune response constituents and cell types that actively engage in the immunoregulatory process [53]. The gut microbiota is also necessary for Foxp3⁺ Treg cell development, differentiation, and function. However, the mechanism by which this task is facilitated is yet to be explored. For example, in specific *Clostridium* clusters, Treg induction may be independent of pattern recognition receptors (PRRs) or based on MyD88-dependent mechanisms [54], while in the case of *Bacillus fragilis*, the process appears to be mediated by Toll-like receptor-2 (TLR-2) signaling by polysaccharide A [55]. Furthermore, SCFAs derived from microbial metabolism, particularly butyrate, have indeed been coupled with Treg development and function [56]. SCFAs have been visualized to stimulate G-protein-coupled receptors by IECs and have been shown to govern Tregs via epigenetic regulation (enhanced acetylation) of the Foxp3 locus [57] (Figure 1).

3. Host Factors Controlling Gut Microbiota

The host selects its gut microbiota by generating various molecular signals and effector molecules that regulate the framework of microbiota-colonized surfaces and thus influence their composition [58]. Numerous associated host factors regulate the constitution of gut microbiota, and some of these factors are mentioned below.

3.1. Delivery Pattern

The mode of delivery is regarded as an essential influential factor in developing the gut microbiota [59]. The composition and architecture of the gut microbial community vary significantly among infants born via cesarean section and those born vaginally [60]. The maternal/vaginal microbiota is denied to Caesarean-born babies, and the first exposure is marked by the absence of strict anaerobes and facultative anaerobes such as *Clostridium species* [61]. After birth, the delivery method influences gut microbial development in the early stages of life. *Prevotella* and *Lactobacillus* from the mother's vaginal microbial community predominate the foremost gut microbiota of newborns delivered vaginally [62]. On the other hand, those born via cesarean section receive their gut microbiota from their skin, which tends to result in the dominance of the following microbial populations: *Streptococcus*, *Corynebacterium*, and *Propionibacterium*. [63].

3.2. Infant Feeding

Infant feeding is another crucial factor that regulates gut microbial assembly, as the mother's milk is rich in different prebiotic microorganisms that shape the infant's gut microbiota [64]. More than 700 species of bacteria are present in a planktonic state in milk that maintains a balance with the immune cells present in milk [65]. Though the microbial diversity varies with lactation time [66], *Streptococcus*, *Staphylococcus*, *Pseudomonas*, and *Acinetobacter* are the predominant species found in breast milk. The average bacterial load is $\sim 10^6$ microorganisms/mL, which increases by about 100 times as the infant receives 8×10^8 bacterial cells with each 800 mL of breast milk consumed [67]. Colostrum samples taken at 1 and 6 months show a different diversity of bacteria, including *Prevotella*, *Leptotrichia*, and *Veillonella*, along with other microorganisms [68]. During breastfeeding, the skin

interface allows some bacteria to become mixed with healthy milk microbiota [69]. The milk microbiome plays a vital role in infant growth and development. Obese mothers have lower diversity and a distinct microbiota composition in their breast milk compared to normal-weight mothers [70]. Immune cells that are interchanged through feeding, such as IgA, are essential for the inert immunity of infants [71].

3.3. Medication: Antibiotics

Antibiotics are a two-edged sword, as their inappropriate utilization kills both morbid and beneficial microbial flora indiscriminately, permitting dysbacteriosis and the growth of undesired microbes [72]. Antibiotic therapy diminishes the overall diversity of the intestinal flora, including the deprivation of some essential taxa, resulting in metabolic shifts, increasing the susceptibility to colonization, and the evolution of antibiotic-resistant bacteria [73]. In adults, a combination of gentamicin, meropenem, and vancomycin increased the popularity of *Enterobacteriaceae* and certain other pathobionts while decreasing the prevalence of *Bifidobacterium* and butyrate-producing organisms [74]. The inappropriate intake of antibiotics disrupts the crucial mechanism of the intestinal flora's "competitive exclusion strategy," [75] which exerts nutrient competition, niche exclusion, and competitive metabolic interaction to hamper the growth of pathogenic microbial diversity [76].

3.4. Genetics: mi-RNAs

Small non-coding RNA and mi-RNAs are another essential host factor that might be used to appraise the intestinal microbiota's composition, shape, and structure [77]. Micro RNAs are single-stranded, short-length (approximately 18–23 nucleotides) endogenous RNA molecules that are synthesized within the nucleus and that are carried out to the cytoplasm afterward to regulate gene expression (gene silencing) [78]. As specific mi-RNA has a high degree of sequence complementarity with the target mRNA, mi-RNA hybridizes with the 3' untranslated region of the target gene, facilitating mRNA degradation or impeding translation [79].

Liu et al. demonstrated that inter-species gene regulation via fecal microRNAs (miRNA) facilitates host control of the gut microbiota [80]. Fecal miRNA is predominantly synthesized by Hopx-positive cells and intestinal epithelial cells, as these two cells are found to secrete mi-RNA-containing exosomes. A direct correlation between mi-RNA-deficient IEC and associated gut dysbacteriosis has been studied in detail, and how fecal transplantation can rehabilitate intestinal flora has also been addressed as well, suggesting the potential role of mi-RNA in the regulation of the gut microbiota. Mi-RNAs can enter gut microbial cells; target bacterial genes precisely; and govern the growth, microbial gene transcripts, and prevalence of a particular gut microbiota [81]. Evidence that miRNAs may play a role in gut bacterial development comes from the improved in vitro growth of the bacteria *Fusobacterium nucleatum* and *E. coli* via hsa-miRNA-515-5P and hsa-miRNA-1226-5p, respectively. This was accomplished by cultivating the strains using synthetic miRNAs [82]. The administration of fecal miRNAs has effects that shape the gut microbiota [83].

4. Microbiota–Gut Brain Interconnection

The gastrointestinal tract and brain are interconnected via the paracrine effect of the signaling molecules secreted in the gut and vice versa. This reciprocal transmission of messages between the gut and brain, thereby performing functional regulation, is termed the "Gut–brain axis" (GBA) [84]. The GBA allows information to be exchanged back and forth between the digestive system and the central nervous system [85]. Interestingly, the gut microbiota and its secretions may modulate brain physiology by governing numerous processes such as development, aging, maturation, homeostasis, and various brain functions [86]. Brain behavior is directly impacted by microbial secretions [87]. According to the research on germ-free (GF) mice, the gut microbiota synthesizes many metabolites in the bloodstream; however, most of them are then modified by the host. These substances have a significant impact on mammalian behavior and neuroendocrine responses [88].

The microbes can influence neural activity by modulating various cerebral biochemical signaling pathways, potentially triggering cognitive impairments [89], thus suggesting their significant impact on brain functioning and physiology. Recent studies have focused on the intellectual effect of gut-on-brain and vice versa. The gut microbial community is vital for proper brain growth and mature cognitive function. The microbiota has been known to regulate critical neuroplasticity-related mechanisms in the adult brain, such as neurogenesis [90] and microglial activation [91]. The brain, on either hand, can alter the composition and different activities of the gut microbiota, such as its role in food absorption and metabolism [92]. Through the autonomic nervous system, the brain appears to significantly influence the structure, composition, and activity of the microbial community in the intestine. It can potentially influence regional gut motility, microbial gene expression via luminal hormone secretion, intestinal transit and secretion, and gut permeability [93]. Previous studies from our research group have demonstrated the pivotal role of pathogenic co-infection in making a conducive environment in the succession of multifactorial diseases such as cancer and parasitic diseases such as malaria [94]. Our investigations have corroborated previous reports that exposure to a bacterium (*H. pylori*) is associated with EBV reactivation, ultimately contributing to the aggravation of cancer pathology. When co-cultured with EBV, the I10 strain of *H. pylori* resulted in the elevated expression of various bacterium-associated pathogenic genes such as *cagA* and *babA* [95]. At the same time, EBV-associated lytic (*gp350* and *bzlf1*) and latent (*ebna1*, *ebna3c*, *Imp1*, *Imp2a*, and *Imp2b*) genes were also found to be elevated in co-infected cells compared to single pathogen infection. This study provides evidence that co-infection creates a favorable microenvironment that promotes the growth of both pathogens, synergistically contributing to exacerbating disease severity [96]. Similarly, one pathogen may promote the production of another pathogenic microbe's morbid gene, resulting in disease pathology aggression. Therefore, we took up the task of summarizing the consequences of gut microbial diversity on the brain and vice versa, focusing on pathogens such as *Helicobacter pylori* and Epstein–Barr virus. The microbiota–gut–brain axis is an arising and evolving concept for studying the influence of two distantly separated biological systems on one another [97]. This new discipline focuses on the induction, etiology, and progression of various metabolic and mental dysfunctions. Hence, understanding the microbial diversity of the gut–brain axis could also be utilized to develop novel therapeutic strategies to overcome neurological disorders. Therefore, in the current article, we mainly focused on reviewing the underexplored synergetic effects of pathogens and their interactions on the onset and progression of numerous pathologies, including cancers and neurological disorders. Additionally, the present article has considered the outcome of various gut microbial secretions in the advancement and modulation of AD, MS, PD, and HD.

5. Routeways for the Bidirectional Communication between the Gut and Brain

The GBA allows multimodal interactions between cerebral and intestinal functions, including the neuronal, immune signaling, and microbial-derived secretory signaling molecules, via the production of various neurotransmitters (GABA, serotonin, and dopamine) from the gut microbiota [98] (Figure 2). The gut microbiota produces neuroactive metabolites such as neurotransmitters or their precursors, which might alter the associated neurotransmitters' concentrations [99]. This demonstrates that the neurotransmitter synthesis mechanism in the gut may impact the brain's neuronal activity and cognitive capabilities, either directly or indirectly [100]. GBA's vast communication system comprises a cascading program that integrates the CNS (brain and spinal cord), enteric nervous system (ENS), autonomic nervous system (ANS), and hypothalamic–pituitary–adrenal (HPA) axis [101]. The parasympathetic and sympathetic limbs of the ANS drive both afferent and efferent signals originating in the lumen and CNS, respectively [102]. The signals released from the lumen are sent to the CNS through the spinal, enteric, and vagal pathways, while signals from the CNS are directed to the intestinal wall [103].




Figure 2. Illustrative diagram depicting the interconnection between gut and brain. The proposed bidirectional communication is firmly affected by various pathways, including the autonomic nervous system (ANS), hypothalamic–pituitary–adrenal (HPA), immune pathways, enteric nervous system (ENS), endocrine pathways, and neural pathways. Gut microbiota can produce microbial metabolites that activate the neuroenteric plexus, stimulate neuropeptide production in the brain, and increase gut–blood barrier and blood–brain barrier (BBB) permeability. The brain releases molecules that stimulate the function of the gut and neuroendocrine plexus.

5.1. Neuronal Pathway: Activation of the Vagus Nerve

The human gut has an enormous network of nerves intertwined with it, referred to as the ENS [104]. The vagus nerve, which ties the body’s visceral organs to the central nervous system, has been considered a central line of communication for the gut microbiota to monitor and control the brain and behavior. The tenth cranial nerve, the vagus nerve, conveys signals to and from the intestinal system to other organs (including the brain). It forms the most crucial part of the ANS and is the longest composite nerve that mediates communication between these two systems. The sensory neuron, which contributes 80% of the overall communication (transmits microbial-derived metabolites from gut to brain), and the motor neuron, which contributes roughly 20% of the total communication (carries the signal from the brain to the gut), are responsible for the information exchange [105]. Even though the gut microbiota can interact via endocrine and immunological routes, hijacking vagus nerve signals is undoubtedly the swiftest and most straightforward way for the microbiota to influence the brain [106]. The vagus nerve, which disseminates informative signals from the gastric mucosa to the CNS, appears to be involved in microbial communication with the brain. Indeed, no neurochemical or behavioral effects were observed in the vasectomized mice group, indicating that the vagus nerve is the pivotal modulatory fundamental communication medium between the microbiota and the brain [107]. Animal studies have established that the gut microbiota may stimulate the vagus nerve and that this activation is important in exposing effects on the brain and, as a result, behavior [108]. Early on, such information was available through the examination of pathogen-infected animals. C-fos expression has been lowered in the PVN of rats infected with *Salmonella typhimurium* after subdiaphragmatic vagotomy [109] even though *S. typhimurium* infection has been linked to intestinal inflammation; new research suggests that microbial diversity in the gastrointestinal system can effectively trigger brain circuitry regardless of whether or

not there is an immune response [110]. The anxiolytic effect observed after *Bifidobacterium longum* therapy in a model of chronic colitis-related anxiety-like behavior was missing in mice vasectomized before colitis induction [111]. In studies, the gut microbiota has been found to alter these vagal-mediated effects. Certain bacterial strains have been shown to use vagus nerve signals to interact with the brain and to influence behavior [112]. For example, a subclinical dose of the diarrhea-causing bacteria *Campylobacter jejuni* enhanced anxiety-related behavior and Fos immunoreactivity in vagal afferent cell bodies and in nucleus tractus solitaries (NTS), the brain's major projection side of the gut-related vagal afferents [113]. The vagus nerve is the chief constituent of the parasympathetic nervous system, which regulates a wide range of biological functions such as mood, immune response, metabolism, and cardiac regulation [114].

5.2. Microbial Signaling Molecules as a Pathway of Communication

The generation of gut microbial metabolites exhibits a vital role in the crosstalk between the gut and brain [115]. Through the immunological and endocrine pathways, gut microbiota metabolites influence the nervous system's function [116]. The gut microbiota, for example, can emit neuro-signaling molecules such as catecholamines, GABA, melatonin, and acetylcholine (ACh) to control the CNS via the vagus nerve [117]. SCFAs (acetate, butyrate, and propionate) and other signaling molecules such as LPS, serotonin, and GABA produced by gut microbiota seem to affect the activity of the ENS that eventually modulates the afferent neural pathway, which carries the signal further to the brain. SCFAs might boost the sympathetic nervous system and mucosal serotonin secretion, modulating the brain's memory and learning processes [118]. Apart from SCFAs, several other microbially induced compounds, such as secondary bile acids (2BAs) and tryptophan metabolites, are also involved in GBA communication [119]. These molecules predominantly transmit signals via interaction with enterochromaffin cells (ECCs), enteroendocrine cells (EECs), and the mucosal immune system. In contrast, some of them may also penetrate to enter the systemic circulation and intestinal barrier and may perhaps cross the blood–brain barrier [120].

5.3. Immune Signaling Pathway

The gut microbiota extensively regulates the formation, function, and maturation of the mucosal immune system, implying its possible involvement in mood and behavior control [121]. Segmented filamentous bacteria (SFB) are effective gut B and T lymphocyte stimulators in the gut [122]. The gut microbiota is also reported to use TLRs to interact with the host [123]. Interestingly, TLR-10 is expressed in various cells in the human body, such as intestinal epithelial cells, mast cells, macrophages, neutrophils, dendritic cells, lymphocytes, glial cells, and neuronal cells [124]. Microbial components can activate TLR 1–10, producing IL-1 β , TNF- α , IL-6, and IL-8 [125]. Another study on TLR-4 knockout mice reported that TLR-4 facilitates inflammatory reactions and gastrointestinal problems via gut dysbiosis and leaky gut in a Gulf War disease model [126]. Additionally, the microbiota can influence hormone peptide signaling by synthesizing peptide-like antigenic proteins derived from the gut microbiota [127].

6. Correlation between the Gut Microbiota and Cerebral Function

The gut microbiota is known to have various health benefits in humans [128]. Moreover, recent analysis has shown that the gut microbiota can influence CNS health and diseases [129]. Thus, the exploration of the microflora residing in the brain–intestinal axis has gained popularity, garnering the concern of both gastroenterologists and neuroscientists. Recently published research articles have revealed that the gut microbiota regulates various basic neurodevelopmental and cognitive processes, such as the formation and maintenance of the integrity of the blood–brain barrier (BBB), microglia maturation, neurogenesis, and myelination, as well as the production of neurotrophins, neuromodulators,

and their respective receptors [130]. These findings show that the gut microbiota plays a significant role in influencing normal human neurodevelopment [131].

The BBB could be impaired by excessive quantities of superoxide, the stimulation of MMPs, and the elevation of inflammatory mediators in the CNS [132]. Destruction of the BBB significantly increases permeability and leakage, eventually resulting in immune cell infiltration to the CNS and eventually to neuroinflammation [133]. The finding that the gastrointestinal microbiota governs neurodevelopment is absolute proof of the gut microbial population as a cerebral peacekeeper [134]. Furthermore, in rats and mice, the gut microbiota influences postnatal and adult ENS development [135]. It is worth highlighting that in mice, the gut microbiota modulates the permeability of the BBB [136]. Mono-colonization with *Bacteroides thetaiotaomicron* or *Clostridium tyrobutyricum*, as well as sodium butyrate therapy, alleviates BBB permeability in germ-free (GF) mice compared to in control animals by intensifying the biologically functional molecule of tight junction proteins [137]. Tight junction integrity is critical for sustaining BBB function [138]. These data suggest that the gut microbial community and its products may be required for appropriate BBB permeability [139].

The bacterial population residing in the ileum part of the intestine can govern both the primary immigration and the homeostatic circulation of neuroglial cells in the intestinal epithelium of mice [140]. GF mice have a much-reduced quantity and density of mucosal enteric glial cells than normal mice [141]. This discovery implies that microbiota and secretory compounds may influence gut homeostasis through enteric glial cells. Moreover, these cells in the ileum connect bacterial signals to the neurological system of the host [142].

The human body's second genome is assumed to be gut microbiota [143]. Its makeup and variety are altered regularly depending on the circumstances. The concentration levels of numerous chemical compounds differ significantly between GF and wild-type (WT) mice in the early detection of the CNS metabolome, suggesting that the microbial community is firmly related to brain health, disease, and functions such as learning, memory, development, and behavior patterns [144]. The hippocampus serves as a learning and memory center, regulating memory encoding, spatial navigation, and memory consolidation, and is associated with mental diseases such as dementia [145]. All prior research has shown a connection between gut microbiota and hippocampal plasticity, neurochemicals, and function [146]. In a study conducted by Matsumoto and colleagues, neurotransmitters and hippocampal amino acids in GF mice at postnatal week seven were found to be significantly different from those in specific pathogen-free (SPF) mice, with decreased amounts of several amino acids, including "A, R, L, Q, I, F, V" and GABA, and increased amounts of "S". GF mice had significantly higher levels of N-acetyl-aspartate, creatine, taurine, and lactate compared to SPF mice, while succinate levels were much lower [147]. Furthermore, the GF mice hippocampus revealed a significant increase in synaptic function via the upregulation of synapse-promoting genes, responsive microglia markers, and synaptic density, which could altogether be recovered by conventional murine microbiota or human *Bifidobacterium* species colonization, implying that *Bifidobacteria* are associated with the initiation of the functional mechanisms in the neural pathway in the hippocampus [148].

However, another study conducted by Luck's group showed that the gut microbiota modulates brain-derived neurotrophic factor (BDNF) and cAMP response element-binding protein (CREB) in the hippocampus [149]. BDNF governs activity-mediated synaptic plasticity and psychological disorders [150], whereas CREB regulates genes involved in neural development, synaptic plasticity, acquisition, and memory [151]. Additionally, antitumor flavonoid quercetin, a phytochemical metabolite, has been demonstrated to improve cognitive performance by increasing gut microbiota and relative abundance of *Facklamia*, *Glutamicibacter*, and *Aerococcus*, resulting in enhanced hippocampal BDNF, thus improving learning and memory [152]. Furthermore, investigations have demonstrated that the gut microbiota modulates the shape and neurogenesis of the hippocampus. Studies on GF animals provide compelling data that the hippocampus of GF mice was noticeably

larger, with pyramidal neurons that were shorter, stubby, and far less branched with mushroom spines and granule cells compared to the control mice [153].

7. Gut Dysbacteriosis: Consequences, Diagnostic and Therapeutic Options

7.1. Microbial Imbalance Leads to Several Neurological Disorders

Dysbiosis or dysbacteriosis is a disorder originating from the disruption or reorganization of the gut microbiota, resulting in a severe imbalance between beneficial and harmful microbes [154]. The following are the main factors that contribute to gut microbiota dysbiosis: (a) lack of helpful microorganisms, (b) excessive proliferation of potentially pathogenic bacteria, and (c) a decrease in the overall microbial diversity in the gut [155]. The disruption of microbial balance within the gut has been incriminated in many pathologies, including in gut-related diseases and neurodegenerative disorders (Figure 3 and Table 1) [156–158]. Previous studies have manifested mounting evidence that gut microbiota dysbiosis is functionally linked to brain immunological dysfunctions, contributing to poor mental health [159]. Digestion, immunopotentiality, the encouragement of microvilli development, the fermentation of dietary fibers, and pathogens required for gastrointestinal tract colonization depend on the gut microbiota [160]. According to several observational and animal studies, the gut microbiota appears to have an essential role in the neuropathogenesis of CNS disorders by GBA function alteration [161]. Dysbiosis can result in neuroinflammation by elevating bacterial metabolites and inflammatory cytokines in the gut and BBB [162]. This could play a role in the development of numerous neurodegenerative diseases, including AD, PD, MS, and amyotrophic lateral sclerosis (ALS) [163].




Figure 3. Variation in the composition of microbial diversity at the phylum level in numerous neurological disorders (AD, MS, PD, and HD). Changes in the occurrence of gut microbial population, dominant gut phyla, including *Actinobacteria*, *Bacteroidetes*, *Firmicutes*, *Proteobacteria*, *Verrucomicrobia*, and *Fusobacteria*, in neurological illnesses compared to in healthy controls.

Table 1. Variation in the composition of microbial diversity at the genus level in numerous neurological disorders. Abundance or scarceness of specific genera in a particular neurological disease compared to in healthy controls, where ↑ represents higher in number compared to healthy ones, ↓ represents lower in number compared to healthy ones, and - represents not reported yet.

GENUS	Alzheimer's Disease	Parkinson's Disease	Huntington's Disease	Multiple Sclerosis
<i>Bifidobacterium</i>	↓	↑	↑	-
<i>Clostridium</i>	↓	↓	↓	↑
<i>Dialister</i>	↓	-	-	-
<i>Turicibacter</i>	↓	-	-	-
<i>Bacteroides</i>	↑	↓	↓	-

Table 1. Cont.

GENUS	Alzheimer's Disease	Parkinson's Disease	Huntington's Disease	Multiple Sclerosis
<i>Blautia</i>	↑	↓	-	↑
<i>Bilophila</i>	↑	-	-	-
<i>Lactobacillus</i>	-	↓	-	↓
<i>Faecalibacterium</i>	-	↓	↑	-
<i>Coprococcus</i>	-	↓	-	-
<i>Prevotella</i>	-	↓	↑	↓
<i>Akkermansia</i>	-	↑	-	↑
<i>Methanobrevibacter</i>	-	-	-	↑
<i>Butyricimonas</i>	-	-	-	↓
<i>Collinsella</i>	-	-	-	↓
<i>Slackia</i>	-	-	-	↓
<i>Megamonas</i>	-	-	↓	-
<i>Gemmiger</i>	-	-	↑	-
<i>Allistipes</i>	↑	-	↓	-

An SCFA-producing gut microbiota is essential to the host's health and well-being [164]. SCFA deficiency can have various negative consequences, including inflammation, eventually contributing to neurological illnesses, namely AD, PD, and MS [165]. The comparably low abundance of these microbes might cause neurological illness etiologies or may just be an indicator of disease development [166]. These discoveries potentially pave the way for new treatment methods based on microbiota modifications. Certain compounds secreted by the gastrointestinal microbiota have subsequently been associated with cognitive and cerebrovascular illnesses [167]. New therapy techniques, including prebiotics and probiotics, may help to restore the gut microbiota, alter the gut–brain barrier, and reduce the risk of certain pathologies [168].

7.2. Strategies to Prevent Dysbiosis of Microbiota

The maintenance of a balanced gut microbiota is positively correlated with the host's health status [169]. Therefore, restoring the original gut microbiota is essential to escape the cascading effect of disturbed microbial diversity, resulting in neurodegenerative disorders [170]. Several strategies have been studied and used to maintain equilibrium between beneficial and pathogenic bacteria in the gut [171]. One of the famous and effective strategies is the supplementation of probiotics and prebiotics in colonizing a healthy microbial community [172,173] (Table 2).

Table 2. Modulation of the gut microbiota as a therapeutic approach for several neurological deformities. Direct modulation of the gut microbiota can be a potential therapeutic target for treating neurological disorders. The table summarizes recent research findings on the reformation of the gut microbiota by probiotic therapies and fecal microbiota transplantation to overcome the neuropathologic condition.

S. No.	Therapeutic Approach	Phylum/Genus or the Name of Particular Bacteria	Neurological Disorder	References
1.	Recolonization of beneficial bacteria	Oral administration of <i>Bacteroides fragilis</i>	Experimental autoimmune encephalomyelitis	[174]
2.	Fecal microbiota transplantation	Fecal microbiota suspension was injected through a TET tube.	Parkinson's disease	[175]

Table 2. Cont.

S. No.	Therapeutic Approach	Phylum/Genus or the Name of Particular Bacteria	Neurological Disorder	References
3.	Bacteriotherapy	Transcolonoscopic infusion of 13 non-pathogenic enteric bacteria	Chronic Fatigue Syndrome (CFS)	[176]
4.	Administration of human commensal bacteria	<i>Bacteroides fragilis</i>	MIA mouse model of ASD (autism spectrum disorder)	[177]
5.	Probiotic supplementation	<i>Lactobacillus acidophilus</i> , <i>L. fermentum</i> , <i>Bifidobacterium lactis</i>	Alzheimer's disease	[178]
6.	Fecal microbiota transplantation	FMT from healthy uninjured mice	Spinal cord injury (SCI) mice model	[179]
7.	Probiotic supplementation	<i>Bifidobacterium bifidum</i> BGN4 and <i>Bifidobacterium longum</i> BORI	Alzheimer's disease	[180]
8.	Probiotic supplementation	<i>Bifidobacterium infantis</i>	Rat maternal separation (MS) model of depression	[181]
9.	Fecal microbiota transplantation	Fecal suspension injected into the colon through the catheters	Traumatic brain injury (TBI) in male Sprague Dawley rats	[182]
10.	Probiotic supplementation	<i>Streptococcus thermophilus</i> , <i>Bifidobacterium lactis</i> , <i>Lactobacillus acidophilus</i> , <i>Lactobacillus helveticus</i> ,	Alzheimer's disease triple-transgenic mice	[183]

Probiotics are live bacteria that confer a health benefit to the host when taken in the requisite amount [184]. They can be used both to stave off the onset of dysbiosis when an individual comes across a predisposing condition (such as the inappropriate use of antibiotics, aging, chronic gut inflammation, imbalanced diet, etc.) [185] and as a curative agent to restore the healthy microorganisms to balance the ongoing condition of dysbiosis [186]. Probiotics should fall in the category of the strains resembling our normal gut microbiota, designated as GRAS (generally regarded as safe). GRAS is known as “health-friendly bacteria” that exhibit beneficial properties for the host’s health. These live bacteria can maintain good viability, are not toxic or pathogenic to the host, are good enough to extract nutrients from a regular diet, and do not interfere with the body’s homeostasis [187]. Some other immunological benefits of probiotics include activating the local immune cells (macrophages), helping in boosting the immune response, and modulating cytokine production [188]. Commonly used probiotics include *lactic acid bacteria*, *Bifidobacteria*, yeast *Saccharomyces boulardii*, *enterococci*, and Gram-negative bacteria *Escherichia coli*, [189]. *Bifidobacterium* is a well-known microorganism that produces essential vitamins, enzymes, and some acids. It also has immune activation properties resulting from lowered gut pH, thus inhibiting the growth of certain pathogens [190]. The central idea from previous studies about the benefits of probiotics leads to the conclusion that (i) probiotics can help reduce the ubiquity and severity of infectious diseases [191,192]; (ii) their unique ability to restore the gut microbiota allows them to be used as the sole treatment for many intestinal disorders and neurological disorders [193]; and (iii) if antibiotics are required, probiotics can be used in conjunction with antibiotics to decrease treatment duration and side effects [194].

Prebiotics, defined as a non-digestible fermented food ingredient, provoke the activity or growth of a number of healthy/good bacteria in the gut that play a pivotal role in human health and that naturally occur in different food products [195]. They are also widely used to rebalance the intestinal microflora. According to the International Scientific Association of Probiotics and Prebiotics (ISAPP), “dietary prebiotics are the selectively fermented ingredient that results in specific changes in the composition and/or activity of the gastrointestinal microbiota, thus conferring a benefit upon host health” [196]. By definition, they may also include a substrate consumed by the host-microbial community to confer a health benefit to the host. Prebiotic substances must have the following characteristics: they must be fermentable, capable of stimulating the activity of gut microbial diversity, and

be resistant to stomach acid and mammalian enzymes [197]. Prebiotics can revitalize the gut microbiota in the intestine. Fructo-oligosaccharides (FOS) and galacto-oligosaccharide (GOS) are the most classified forms of prebiotics [198]. Some microorganisms such as *Aspergillus* sp., *Penicillium* sp., *Arthrobacter* sp., and *Aureobasidium spare* are excellent sources of a crucial key enzyme known as fructosyl-transferase (FTase) for the production of FOS [199]. Interestingly, the prebiotics fructooligosaccharides (FOS) and galactooligosaccharides (GOS) boosted *Lactobacilli* and *Bifidobacteria* development in the intestines and increased hippocampus BDNF and NR1 subunit expression in comparison to controls [200]. GOS also increases NR2A subunits in the hippocampus nucleus and NR1 expression in the frontal cortex, as well as plasma D-alanine, an NMDA receptor agonist.

Predacious employs a range of intercommunication mechanisms, one of which is predation. The term “predator” refers to bacteria that actively seek out and destroy their prey, eating their macromolecules for nourishment [201]. Predatory bacteria are important for regulating and changing bacterial populations in various settings. Even though these bacteria are nearly everywhere, only a few species have been thoroughly investigated for their potential use [202]. Because predatory bacteria are frequently smaller than their prey, they can penetrate the prey, kill it from the inside, and multiply. Interestingly, numerous predatory behaviors have emerged in the bacterial community; epibiotic predation does not require intracellular proliferation. Predatory bacteria are used to re-equilibrate a dysbiotic gut microbiota characterized by Gram-negative bacteria. In birds, the use of *B. bacteriovorus* to reestablish eubiosis in the gut microbiota has proven successful [203]. It is noteworthy that any potential therapeutic use of predatory bacteria should be based on appropriate and reliable experimental data to define the dosages to be supplied accurately. Otherwise, predatory bacteria, such as antibiotics, might become harmful in the long run if administered inappropriately.

7.3. Potential Biomarker for Dysbiosis and Its Implications

It is thought to be a difficult undertaking to identify “ideal biomarkers” for many diseases, including metabolic disturbances and neurological disorders [204]. Furthermore, the discovery of microbiome-based biomarkers can improve the precision of illness classification when paired with clinical data and other biomarkers [205].

One recent analysis found that urolithin detection in urine is a practical, non-invasive, and quick method that can reveal dysbiosis of the gut microbiota and intestinal inflammation in Parkinson’s disease patients [206]. *Roseburia species* is at a reduced level in Parkinson’s disease and hence can be a potential marker [207].

Several microbial metabolites may be used as predictive biomarkers to track illness conditions in association with gut microbial dysbiosis [208]. A peripheral gut microbiota-derived metabolite, indoxyl sulfate, has the potential to serve as a biomarker for metabolite profiling and diagnostic suitability for dysbiosis and neurologic signature [209]. The relative abundance of *Enterobacteriaceae*, which are expected to be used as clinical biomarkers of post-stroke cognitive impairment (PSCI), may have the ability to predict PSCI in post-stroke patients [210].

8. Effect of Intestinal Microbiota-Derived Metabolites on Neurological Disorders

The altered composition of the gut microbiota and its various microbial metabolites can lead to numerous brain disorders by different mechanisms. The present review article describes the effects of gut microbial-derived biomolecules on the onset and progression of most common neurological disorders, such as AD, PD, HD, and MS.

8.1. Aggregate-Forming Tendency of Gut Bacterial Proteins in Alzheimer’s Disease

AD is a common progressive neurodegenerative condition that causes brain shrinkage and cell death [211]. It is defined by the development of two unusual formations termed plaques and tangles, which are the main suspects in nerve cell damage [212]. Tangles are intracellular twisted fibers of a protein called tau that builds up inside cells, whereas

plaques are the extracellular aggregation of a protein fragment called amyloid (A β) that builds up in the spaces between nerve cells [213].

Recent studies have suggested an association between gut microbiota-related biomolecules with the progression of Alzheimer's. Elevated levels of numerous Gram-negative bacteria in the intestinal tract and the cerebrum of AD patients resulted in the idea of the pathogenesis and contribution of these microbial communities in respective neurodegenerative disorders [214]. LPS, a major characteristic component of Gram-negative bacteria such as *Helicobacter pylori*, *E. coli*, and *Salmonella*, was found to play a vital role in the aggregation of amyloid fibers extracellularly [215]. LPS has a potential role in the fibrillogenesis of beta-amyloid, leading to the analytical deficiency studied in in vitro conditions [216]. Meanwhile, various gut bacterial strains generate a remarkable amount of effective amyloid-like proteins. According to a discovery by Chapman's group, extracellular fibers termed "curli", which are formed by *E. coli* and other gut microbiota, have structural and physicochemical features with amyloids [217]. Curli was the first amyloid to be characterized in a new class of "functional" amyloids that is quickly expanding. According to scientific analysis, *Staphylococcus*, *Streptococcus*, *Mycobacteria*, *Salmonella*, *Citrobacter*, *Klebsiella*, and *Bacillus* species can construct extracellular amyloids [218].

Based on previous studies, a possible hypothesis could be drawn for the potential neurotropic role of bacterially secreted pathogenic peptides (such as cagA, vacA, and babA-H. *pylori*) and their virulent outer membrane proteins (*Hop Q*) in seeding amyloid β accumulation, which may further contribute to neuroinflammation. By employing bioinformatics tools, the aggregation tendency of these peptides could be deduced and compared with the positive control, and then inference regarding the contribution of the aggregates forming peptides could be determined [219].

8.2. Impact of Microbiota on the Induction of Parkinson's Disease

PD is a neurodegenerative disorder characterized by the degeneration of dopamine-producing neurons in a specific region of the brain called the substantia nigra pars compacta and the accumulation of protein α -synuclein [220]. These pathophysiological conditions lead to movement-related disorders in Parkinson's patients [221]. Alterations in the gut microbiota composition is strongly associated with PD. The gut microbial diversity with certain phyla, including *Firmicutes*, *Bacteroidetes*, and *Fusobacteria*, was deformed in patients with Parkinson's symptoms compared to the control [222].

Brain behavior's chief mediator and controller is the enrichment of specific pathogenic microbial phyla and less abundant beneficial microbial communities. SCFA, which functions as an anti-inflammatory agent and helps to prevent neuroinflammatory microenvironment creation, was found scantily in the GI tract of Parkinson's patients due to the lower number of SCFA-producing microorganisms (*Bacteroidetes* and *Firmicutes*) [223]. A lower abundance of bacteria such as *Clostridium tyrobutyricum*, which produces high levels of butyrate, has an intended role in the progression of PD [224]. Another potential function of SCFA is maintaining the integrity of BBB and being capable of ameliorating a dysfunctional BBB in germ-free mice associated with an upregulation of tight junction protein expression [225]. Additionally, in Parkinson's patients, LPS from the innumerable gut-residing Gram-negative bacteria were found capable of crossing BBB by modulating tight junction protein expression (similar to occludin and claudin) [226]. These alterations in the expression of tight junction proteins help to accelerate the extravasation of immune cells in the brain [227], the microglial cells result in the higher production of pro-inflammatory molecules, and neuroinflammation is one of the hallmarks of initiation of neurodegenerative disorders (Figure 4) [228].




Figure 4. Probable mechanism of gut microbial population and their metabolites to cause neuroinflammation. Intensification and accumulation of pathogenic gut microbiota due to dysbacteriosis lead to the formation of an inflamed and leaky gut. Microbes and their associated metabolites can enter the peripheral circulatory system, where they further lower the expression of tight junction proteins in brain endothelial cells and lead to the disintegration of the blood–brain barrier that promotes the

entrance of pathogenic microbes, some of them carrying prions into the brain. Meanwhile, microbial metabolites stimulate the extravasation of immune cells into the brain, all of which together trigger cytokine storm, the cause of neuroinflammation, one of the important hallmarks of neurological disorders characterized by degenerated and demyelinated neurons.

The administration of L-DOPA (levodopa), a dopamine precursor, is the most common and effective treatment to manage PD [229]. The widespread presence of *H. pylori* infection has been widely studied in this neurodegenerative disease associated with L-DOPA malabsorption, which results in a higher motor impairment than in uninfected Parkinson's patients [230]. Hence, eradicating *H. pylori* could lead to a remarkable refinement in motor fluctuations by improving the absorption of the Parkinson's medication levodopa [231]. Previous studies focused on this bacterium's ability to decrease the bioavailability of L-DOPA to the dopaminergic neuronal cells, but the mechanism is still unexplored [232].

One of the possible mechanisms could be the utilization of L-DOPA as a nutrient source by *H. pylori*, making it unavailable to be utilized by the host. Consuming L-DOPA *H. pylori* enhances its virulence, thus contributing further to aggravating this neurological disorder. L-DOPA is derived from the amino acid phenylalanine, which has a significant role in the growth and motility of bacteria [233]. As motility plays one significant role in determining a bacterium's virulence, a hypothesis could be drawn that the consumption of L-DOPA might enhance the expression of the motility (flagellin) gene, consequently inducing motility and resulting in motility in the severity of symptoms of *H. pylori* infection in PD. Another probable mechanism is the secretion of an enzyme by the gut microbiota, which might be responsible for converting L-DOPA into an anomalous product. The biotransformation of L-DOPA into another impractical form could be one of the reasons for making it worthless for Parkinson's therapy and reducing its availability [234].

8.3. Huntington's Disease Association with Gut Dysbiosis

A trinucleotide repeat amplification in the Huntington (HTT) gene, which is distributed extensively across the brain and peripheral tissues, causes HD, a chronic neurodegenerative illness [235]. The major phyla that seem to be altered in HD patients are *Bacteroidetes* and *Actinobacteria*; meanwhile, the *Firmicutes* population was found to be diminished compared to the healthy control (HC) [236]. At the genus level, *Intestinimonas*, *Lactobacillus*, and *Bilophila* were higher in number, and comparatively, the genera *Clostridium* is less abundant than in normal individuals [237]. Huntington's patients' gut microbiota differed significantly from those of healthy subjects. The patients' gut microbiota seemed to be occupied by fewer microbial species, resulting in a less diverse ecosystem [238]. A bacterial species named *Eubacterium hallii* has been linked to various clinical symptoms of Huntington's. Symptomatic individuals with low *E. hallii* counts had more severe motor symptoms [239]. The abundance of *E. hallii* and the timing of symptom manifestation revealed a high and negative connection in pre-symptomatic patients. Intensification of the pathogenic gut microbiota (*Lactobacillus*, *Intestinimonas*), as well as various strains of viruses, may result in the production of harmful secretions that interfere with the production of anti-inflammatory molecules (IL-4) by specific immune cells, allowing bacteria to evade the immune response and contribute to a variety of neurological disorders [240].

A recent study on HD patients reported exclusively repressed levels of lymphokine IL-4 compared to healthy controls, whereas other cytokines remain unaltered. In the HD group, a significant correlation between systemic levels of cytokines and relative abundances of fecal microbiota has been found [241]. Interestingly, A recent study demonstrates a connection between host cytokine response in diseased and normal individuals; as a result, the level of IL-4 was found to be repressed in the HD patient compared to that of controls, while there were no significant changes in the rest of the cytokine molecules in both cases.

However, the previous study did not focus on the linkage between IL-4 secretion and the progression of HD. One of the possible reasons could be the central role of IL-4 in regulating normal brain function and a positive effect on cognitive behavior by synchronizing

the activation of various cerebral growth factors such as BDNF and NGF. Simultaneously, IL-4 acts as an anti-inflammatory agent and ultimately staves off the degeneration or death of neurons [242].

8.4. Substantial Alteration of Human Gut Microbiota in Multiple Sclerosis

MS is a demyelinating autoimmune disease of the central nervous system (CNS) that influences the brain and spinal cord [243]. Autoimmunity against CNS antigens is hypothesized to cause immune cell infiltration, which results in CNS lesions that are diagnostic of MS [244]. Both MS and experimental autoimmune encephalomyelitis (EAE), an animal model of MS, demonstrate BBB impairment [245]. Disruption of the gut microbiota encourages permeability of the intestine, referred to as “leaky gut”, which displays one of the major contributors to the consequence of MS, an autoimmune disease [246]. The alteration of several microorganisms is associated with MS. The gut microbiota was found to be modified in different stages and types of MS, namely RRMS (Relapsing-Remitting Multiple Sclerosis), PRMS (Progressive-Relapsing Multiple Sclerosis), SPMS (Secondary Progressive Multiple Sclerosis), and PPMS (Primary Progressive Multiple Sclerosis) [247]. MS is strongly correlated with the depletion of major metabolite-producing microbes such as SCFA-producing microorganisms (Genera *Clostridium*). Butyrate- and propionate-biosynthesizing microorganisms are alleviated in RRMS compared to HC, which represents great evidence of a positive correlation between SCFA and the regression of MS [248].

Nevertheless, the mechanism of action of SCFA regarding MS plaques is underexplored. One of the possible reasons could be the neuroprotective consequence of SCFA on MS that might have a considerable role in reinstating the integrity of BBB [249]. Various SCFAs might be responsible for the higher expression level of tight junction proteins to maintain the BBB. That is how the depletion of SCFA-producing bacteria influences the cohesive property of BBB and allows the pathogenic myelin-specific T cell to enter the CNS. Additionally, SCFAs have an anti-inflammatory property that regulates the unnecessary generation of immune response by promoting Treg cell differentiation, which further reduces inflammation [250]. Another hypothesis could be drawn by the presence of detrimental bacterial and viral secretion or peptides as a result of dysbiosis, which might mimic the myelin antigen when exposed to the gut and may be responsible for the activation of myelin-specific peripheral T cells that promote their penetration to the CNS via disrupted BBB. This is how T cells could utilize one mechanism to execute the autoimmune attack on oligodendrocytes.

Rumah et al. detected *Clostridium perfringens* type B in a patient’s feces three months following the beginning of MS symptoms [251]. The epsilon toxin (ETX), released by *C. perfringens*, can pass the BBB and cause oligodendrocyte damage, suggesting a plausible mechanism for demyelination in MS. They also discovered a diminished population of *C. perfringens* A in the GI tract of MS patients and enhanced ETX responsiveness by ten times compared to HC. Another analysis revealed that MS patients had a considerably higher Archaeal population (*Methanobrevibacteraceae*) than the controls. *Methanobrevibacter smithii* has high immunogenicity and may cause inflammation in the patient [252]. It has also been observed that many anti-inflammatory microbes were found in decreased abundance in MS patients [253]. Significant changes in *Proteobacteria* microbiota, such as *Shigella* and *Escherichia* abundance, were also reported in pediatric MS when compared with controls.

9. Altered Bile Acid Profile Associates with Neurological Dysfunction

The expression of numerous bile acid receptors as well as changes in bile acid metabolism have been reported to be promising biomarkers for prognostic tools in a plethora of neurodegenerative disorders [254]. In a recent clinical study, an extensive bile acid test was carried out on the plasma of 30 healthy controls, 20 people with moderate cognitive impairment, and 30 people with clinical AD. In contrast to moderate cognitive impairment patients, AD patients had significantly higher levels of glycochenodeoxycholic acid, glycodeoxycholic

acid, and glycolithocholic acid. lithocholic acid (LCA) levels were also significantly higher in AD patients than in controls. These glycine-conjugated bile acids and the presence of LCA provide valuable diagnostic biomarker features [255]. A surgical mouse prodromal PD model was used in a subsequent investigation to further explicate the existence of bile acids. Three of these acids—omega-muricholic acid, tauroursodeoxycholic acid (TUDCA), and ursodeoxycholic acid (UDCA), were found to be substantially reduced in the serum of the group treated with α -synuclein fibrils. UDCA and TUDCA, both neuroprotective secondary bile acids that can pass through the BBB, were markedly affected, with a 17- and 14-fold decrease from the control group [256].

The presence of bile acids continues to have increased relevance in maladies of the CNS beyond neurodegenerative diseases. Several bile acids have been shown to exhibit neuroprotective benefits in a variety of neurodegenerative illnesses, including AD, PD, HD, and retinal degeneration, both in cellular and animal models as well as in human clinical studies [257]. Bile acids have been studied in animal models of AD. Tauroursodeoxycholic acid (TUDCA), an endogenous bile acid, exhibits potent neuroprotective effects in a variety of disease-related experimental paradigms, including neuronal exposure to A β . Treatment of double transgenic mice (APP/PS1) expressing the human amyloid precursor protein with the KM670/671NL Swedish double mutation and the human presenilin 1 L166P mutation under the control of a neuron-specific promoter with 0.4% (*wt/wt*) of TUDCA resulted in the significantly reduced accumulation of A β deposits in the brain and markedly ameliorated memory deficits [258]. Additionally, the effects of bile acids on chemical and genetic models of Parkinson's disease (PD) have been also documented. In C57BL/6 glutathione S-transferase pi (GSTP)-deficient mice, MPTP-induced degeneration of dopaminergic neurons in the nigrostriatal axis was reported to be ameliorated by TUDCA [259].

10. The Role of Dysbiosis in the Aging Process

A negative shift in microflora is a characteristic feature of dysbiosis. An imbalance in the body's natural microflora can lead to physiological alterations. Geriatric alterations can also lead to changes in the macrofloral diversity and, in turn, accelerates aging-related problems [260].

Cross-sectional and longitudinal studies involving various age groups have found an alteration in gut microbial diversity with increasing age [261] (Figure 5). Researchers have found that age-related gut dysbiosis generally triggers the enhanced growth and proliferation of facultative anaerobes [262]. Meanwhile, a massive reduction in the diversity of probiotic bacteria was further followed by the altered *firmicutes/bacteroidetes* ratio. According to the latest analysis, the architecture and composition of the gut microbiota is a predictor of an individual's survival. Moreover, the abundance of *Bacteroides* or a lack of originality in the microbiome is linked to shorter life expectancies and higher morbidity [263]. For instance, dysbiosis in the commensal microbial community has been shown to shorten the lifespan of *Drosophila* [264], but populating the gut of middle-aged African turquoise killifish with bacteria isolated from young donors led to lifetime extension and delayed behavioral deterioration [265].

One of the major risk factors contributing to the emergence of neurodegenerative disorders is aging [266]. The onset of age-related neurodegenerative diseases is facilitated by alterations in the gut microbiota [267]. An upsurge in bacterial LPS in plasma and the brain is coupled with aging [268]. The increase in LPS level is correlated with the increased expression of TLR4, myeloid differential protein-88, and the nuclear translocation of nuclear factor κ B in both intestinal and brain tissues [269]. Additionally, LPS-induced ulcerative colitis mediated systemic inflammation, increasing BBB permeability, leading to inflammation in the substantia nigra region of the brain, and causing dopaminergic loss [270].




Figure 5. Composition of gut microbiota in the healthy aging process. The age-related sequential change in gut microbiota composition and metabolic function from infant to centenarian.

11. Synergetic Effect of Co-Infection and Microbial Interaction on a Neurological Disorder

The onset and progression of various infectious diseases are mainly associated with a specific pathogen, yet it is not always the case. Co-infection with two or more pathogens is now being reported to modulate the severity of neurological diseases aggressively [271]. The human host may become coinfecting with multiple pathogens concurrently or may be simultaneously infected with two or more pathogens. Reactivation of latently inhabiting virulent pathogens upon infection with a new pathogen may also lead to coinfection conditions [272]. Commensal gut microbiota has been found to regulate and activate viral pathogenic genes through various mechanisms that either provoke or repress numerous diseases [273].

The gut microbiota enhances viral infection through various mechanisms, including through facilitating virus genetic recombination. Many viral genomes undergo recombination, which improves their environmental fitness and thus their infectivity [274]. After exposure to the particular commensal gut microbiota, certain RNA viruses gain an advantage in delivering their genomic content into target cells. The interaction of viruses with microbiota aids the viral entry into the host cell. It ensures genetic recombination, which further results in the generation of progeny with increased resistance to the restrictive condition. The resistive capacity of the recombinant population has a more severe effect on infectious disease outcomes. In addition, the gut microbiota also seems to enhance viral stability as it modulates viral replication [275].

Moreover, viral infection has also been associated with the disruption of the gut microbiota, resulting in colon inflammation and ultimately promoting the pathogenesis of various neurological disorders. Viral infection resulting from dysbiosis leads to the enhancement of pathogenic microbial populations such as LPS and cytokine-producing bacteria. This facilitates the aggregation of specific proteins, such as α -synuclein in the gut (ENS), that can translocate to the CNS via the vagus nerve and aggravate neurological pathologies [276]. The studies mentioned above establish that infection with a single pathogen can make the environment conducive to other pathogens' activation. The expression of the morbid genes of co-existing pathogens may become enhanced during co-infection conditions and can be best studied in co-infection models. However, co-infection scenarios are still underexplored regarding the progression of neurodegenerative disorders. A possible hypothesis could be drawn regarding direct infection of one pathogen followed by a

second infection to the neuronal cells and could be used for the study of their pathogenic gene expression, and further specific markers concerning neurological disease could be studied in the co-infected model to analyze the collaborative effect of pathogens on various neurodegenerative markers (Figure 6).




Figure 6. Schematic diagram depicting the effect of a lone bacterial or viral infection and pathogenic coinfection in engendering neuroinflammation. The role of extracellular vesicles from bacteria-infected cells in virus reactivation via PBMC in contributing to neuroinflammation has to be explored. Meanwhile, the analysis of the combined effect of pathogens on various neurodegenerative markers and inflammation could be contemplated in the co-infection models. NB: ? signifies the nonavailability of direct evidence and study needs to be done in this aspect.

12. Modulation of Gut Microbiota for Neurological Disorders in the Perspective of Host-Directed Therapy: Microbiota-Targeted Technique “Fecal Microbiota Transplantation”

The proximal relationship between factors such as gut dysbacteriosis, elevated intestinal permeability, and associated neurocognitive impairment recommends that manipulating the gut microbiota may provide a promising therapeutic option in a group of neuro patients [277]. Fecal microbiota transplantation (FMT) is an approach for directly modifying the victim’s gut microbiota to normalize the proportion and to acquire therapeutic efficacy (Table 2) [278]. FMT is a biological therapy that involves the transmission of fecal matter from one fit individual to another to cure a disease [279]. It is acknowledged as the “ultimate probiotic” because it provides a significantly larger richness and variety of bacterial strains than any readily accessible probiotic. Recently, there has been a surge of interest in the potential advantages of FMT in both gastrointestinal and non-gastrointestinal diseases (Table 2) Lately, case studies of patients with MS [280], myoclonus-dystonia, autism, depression, and chronic fatigue syndrome who have been successfully treated with FMT have opened up new avenues for more promising trials in characterizing FMT as a potential therapy for such conditions. FMT has aided functional ability, favored neuronal axonal regeneration, ameliorated animal metabolic profiling and weight gain, and substantially improved gut barrier integrity and gastrointestinal motility in SCI (spinal cord injury) mice (Table 2).

A study by Yingli Jing's group demonstrated that FMT-mediated gut microbial transformation strengthens motor, cognitive, and GI functions in SCI mice, potentially via the anti-inflammatory functions of SCFA. Elevated neurotransmitter levels of dopamine in PD mice that already had acquired FMT from healthy mice revealed the significant application of the microbiome modulation technique in various nervous system-related disorders [281].

Numerous animal studies and case reports from humans emphasize that FMT has a beneficial impact on PD, MS, AD, and stroke. The possible health benefits of FMT for patients with PD may be ameliorated by impaired α -syn accumulation in the intestinal wall and, consequently, in the brain, as a result of reduced inflammation-induced oxidative stress [282]. Greater availability and potency of levodopa following FMT with feces from donors with reduced levels of bacterial tyrosine decarboxylases in their feces could strengthen the beneficial effects on Parkinson's symptoms [283]. A rise in Treg cell numbers after FMT in MS patients may constrict autoimmunity with demyelination and possibly the progression of the disease. Furthermore, feces from young healthy FMT donors may slow the advancement of AD by decreasing the translocation of pro-inflammatory gut microbiota from the gut to the brain and, as a result, the neuroinflammation processes facilitated by them. In conclusion, fecal microbiota transplantation can restore a drastically changed gut microbiota in neurodegenerative disorders.

13. Function of Modulated Microbial Communities in Healthy Aging and Rejuvenation

Age-related microbiome and gastrointestinal disorders are reversible and treatable to some extent. To change the gut microbiota and encourage healthy aging, a number of rejuvenation techniques have been used, including probiotic medication and fecal microbiota transplantation [284]. Previous research has consistently demonstrated a positive association between the abundance of beneficial microflora and centenarian longevity. For example, *Akkermansia muciniphila* stimulates the production of mucus in the gut, which is essential for maintaining intestinal integrity and other advantageous symbioses [285]. The oral treatment of *Akkermansia* also improves senescence-related phenotypes in intestinal integrity, muscle function, and immune response in aged mice and thus extends the health span, which is further supported by the understanding of the relationship between aging and the gut microbiota [286]. Additionally, the relative abundance of the butyrate producer *Oscillospira* in rejuvenated mice was dramatically raised by all rejuvenation techniques [287].

Besides probiotics, the consumption of some of the chemical compounds has been proven to be helpful for the maintenance of gut microbial health [288]. *N*-Acetylcysteine (NAC) is an FDA-approved drug primarily associated with its anti-inflammatory and antioxidant activity, which supports the maintenance of a cellular redox imbalance [289]. It is a potential drug to prevent glucose metabolic disturbances by reshaping the structure of the gut microbiota.

In HFD-fed mice, NAC significantly ameliorates gut barrier disruption, glucose intolerance, and inflammatory responses [290]. The potential regulatory mechanism may help to reshape the altered gut microbial structure by intensifying the growth of beneficial bacteria such as *Akkermansia*, *Bifidobacterium*, *Lactobacillus*, and *Allobaculum* and by reducing the abundances of harmful bacteria such as *Desulfovibrio* and *Blautia*, which are concurrent with the repair of disrupted metabolic pathways. These investigations reveal the prominent role of NAC in alleviating gut dysbiosis and associated inflammation [291].

14. Conclusions

The architecture and composition of the gut microbiota undergo significant changes throughout time, and these alterations are typically associated with or followed by adverse health consequences. Numerous factors, such as the way of life, stress, nutritional challenges, antibiotics, and the aging process may alter the gut microbiota. It has been established that strategies to counteract these detrimental fluctuations contribute to alleviating symptoms and recovery from certain conditions. The gut–brain axis is an important field of study, with several studies associating changes in gut microbiota composition with a range

of clinical disorders. The discovery of a biological connection between the microbiota, immune signaling, and the CNS shows that the microbiota's microbial metabolites or systemic signals may alter neurological and immune function in the brain. Cancer, autoimmune illnesses, and neurological conditions such as MS, PD, and AD have all been linked to dysbiosis in the gut microbiota. Bacteria impact the immunological responses of the host in part by producing metabolites. By regulating and reactivating each other's pathogenic genes, synergistic interactions between many pathogens may play a role in the onset and progression of various neurodegenerative disorders. The dysbiosis of the gut microbiota is caused by a rise in pathogenic microorganisms, which may be restored by probiotics, prebiotics, predatory bacteria, and fecal microbial transplantation from healthy individuals. These methods may assist in the battle against neurological symptoms resulting from microbial dysbiosis. Furthermore, the discovery of microbiome-based biomarkers can improve the precision of illness, including metabolic disturbances and neurological disorders. In conclusion, the microbial population inhabiting and dwelling on the human body serves a significant function in preserving the human host's health. Any imbalance between "good" and "bad" microflora may cause illness and reduce the quality of life of a person. Therefore, it is essential that the function of microbiota in many diseases, particularly neurological disorders, be highlighted and thoroughly researched.

Author Contributions: Conceptualization, M.K. and H.C.J.; analysis, and interpretation of data in the work, M.K., H.C.J., O.I., B.B., D.T., S.J.; Formal analysis and visualization, M.K., V.B., R.K.P., K.B., A.S., H.C.J.; resources, M.K.; writing—original draft preparation, M.K., writing—review and editing, M.K., V.B., R.K.P., H.C.J.; supervision and validation, H.C.J.; Analysis and interpretation of data in the work, H.C.J. All authors have read and agreed to the published version of the manuscript.

Funding: Indian Council of Medical Research, grant no: BMI/12(82)/2021.

Acknowledgments: We are thankful to the Indian Council of Medical Research (ICMR) for providing a research grant (No: BMI/12(82)/2021). We are thankful to UGC, the Ministry of Human Resources, and DBT for providing fellowships to Meenakshi Kandpal, Budhadev Baral, Shweta Jakhmola, Deeksha Tewari, and Omkar Indari in the form of research stipends. We appreciate our lab colleagues for their insightful discussions and advice.

Conflicts of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as potential conflicts of interest. The authors have reported no potential conflicts of interest.

References

1. Kho, Z.Y.; Lal, S.K. The Human Gut Microbiome—A Potential Controller of Wellness and Disease. *Front. Microbiol.* **2018**, *9*, 1835. [[CrossRef](#)] [[PubMed](#)]
2. Thursby, E.; Juge, N. Introduction to the human gut microbiota. *Biochem. J.* **2017**, *474*, 1823–1836. [[CrossRef](#)] [[PubMed](#)]
3. Matijašić, M.; Meštrović, T.; Paljetak, H.; Perić, M.; Barešić, A.; Verbanac, D. Gut Microbiota beyond Bacteria—Mycobiome, Virome, Archaeome, and Eukaryotic Parasites in IBD. *Int. J. Mol. Sci.* **2020**, *21*, 82668. [[CrossRef](#)] [[PubMed](#)]
4. Ogunrinola, G.A.; Oyewale, J.O.; Oshamika, O.O.; Olasehinde, G.I. The Human Microbiome and Its Impacts on Health. *Int. J. Microbiol.* **2020**, *2020*, 8045646. [[CrossRef](#)] [[PubMed](#)]
5. Belkaid, Y.; Hand, T.W. Role of the Microbiota in Immunity and Inflammation. *Cell* **2014**, *157*, 121–141. [[CrossRef](#)]
6. Pickard, J.M.; Zeng, M.Y.; Caruso, R.; Núñez, G. Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease. *Immunol. Rev.* **2017**, *279*, 70–89. [[CrossRef](#)]
7. Schaubeck, M.; Clavel, T.; Calasan, J.; Lagkouvardos, I.; Haange, S.B.; Jehmlich, N.; Basic, M.; Dupont, A.; Hornef, M.; von Bergen, M.; et al. Dysbiotic gut microbiota causes transmissible Crohn's disease-like ileitis independent of failure in antimicrobial defence. *Gut* **2015**, *65*, 225–237. [[CrossRef](#)]
8. Liu, S.; Gao, J.; Zhu, M.; Liu, K.; Zhang, H.-L. Gut Microbiota and Dysbiosis in Alzheimer's Disease: Implications for Pathogenesis and Treatment. *Mol. Neurobiol.* **2020**, *57*, 5026–5043. [[CrossRef](#)]
9. Wang, Q.; Luo, Y.; Chaudhuri, K.R.; Reynolds, R.; Tan, E.-K.; Pettersson, S. The role of gut dysbiosis in Parkinson's disease: Mechanistic insights and therapeutic options. *Brain* **2021**, *144*, 2571–2593. [[CrossRef](#)]
10. Gkolfakis, P.; Dimitriadis, G.; Triantafyllou, K. Gut microbiota and non-alcoholic fatty liver disease. *Hepatobiliary Pancreat. Dis. Int.* **2015**, *14*, 572–581. [[CrossRef](#)]
11. Rodarte-Acosta, J.S.; Morales-Sánchez, M.A. El rol de la microbiota intestinal en la dermatitis atópica. *Rev. Del Cent. Derm. Pascua* **2021**, *30*, 76–83. [[CrossRef](#)]

12. Yu, C.; Su, Z.; Li, Y.; Li, Y.; Liu, K.; Chu, F.; Liu, T.; Chen, R.; Ding, X. Dysbiosis of gut microbiota is associated with gastric carcinogenesis in rats. *Biomed. Pharmacother.* **2020**, *126*, 110036. [[CrossRef](#)]
13. Dekaboruah, E.; Suryavanshi, M.V.; Chettri, D.; Verma, A.K. Human microbiome: An academic update on human body site specific surveillance and its possible role. *Arch. Microbiol.* **2020**, *202*, 2147–2167. [[CrossRef](#)]
14. The Integrative HMP (iHMP) Research Network Consortium. The Integrative Human Microbiome Project. *Nature* **2019**, *569*, 641–648. [[CrossRef](#)]
15. Berg, G.; Rybakova, D.; Fischer, D.; Cernava, T.; Vergès, M.-C.C.; Charles, T.; Chen, X.; Cocolin, L.; Eversole, K.; Corral, G.H.; et al. Microbiome definition re-visited: Old concepts and new challenges. *Microbiome* **2020**, *8*, 103. [[CrossRef](#)]
16. Lloyd-Price, J.; Mahurkar, A.; Rahnavard, G.; Crabtree, J.; Orvis, J.; Hall, A.B.; Brady, A.; Creasy, H.H.; McCracken, C.; Giglio, M.G.; et al. Strains, functions and dynamics in the expanded Human Microbiome Project. *Nature* **2017**, *550*, 61–66. [[CrossRef](#)]
17. Galland, L. The Gut Microbiome and the Brain. *J. Med. Food* **2014**, *17*, 1261–1272. [[CrossRef](#)]
18. Rowland, I.; Gibson, G.; Heinken, A.; Scott, K.; Swann, J.; Thiele, I.; Tuohy, K. Gut microbiota functions: Metabolism of nutrients and other food components. *Eur. J. Nutr.* **2018**, *57*, 1–24. [[CrossRef](#)]
19. Wu, H.J.; Wu, E. The role of gut microbiota in immune homeostasis and autoimmunity. *Gut Microbes* **2012**, *3*, 4–14. [[CrossRef](#)]
20. Arnold, J.W.; Roach, J.; Azcarate-Peril, M.A. Emerging Technologies for Gut Microbiome Research. *Trends Microbiol.* **2016**, *24*, 887–901. [[CrossRef](#)]
21. Sekirov, I.; Russell, S.L.; Antunes, L.C.M.; Finlay, B.B. Gut Microbiota in Health and Disease. *Physiol. Rev.* **2010**, *90*, 859–904. [[CrossRef](#)] [[PubMed](#)]
22. Wang, S.-Z.; Yu, Y.-J.; Adeli, K. Role of Gut Microbiota in Neuroendocrine Regulation of Carbohydrate and Lipid Metabolism via the Microbiota-Gut-Brain-Liver Axis. *Microorganisms* **2020**, *8*, 527. [[CrossRef](#)] [[PubMed](#)]
23. Park, J.; Kim, C.H. Regulation of common neurological disorders by gut microbial metabolites. *Exp. Mol. Med.* **2021**, *53*, 1821–1833. [[CrossRef](#)] [[PubMed](#)]
24. Korf, J.M.; Ganesh, B.P.; McCullough, L.D. Gut dysbiosis and age-related neurological diseases in females. *Neurobiol. Dis.* **2022**, *168*, 105695. [[CrossRef](#)] [[PubMed](#)]
25. Baizabal-Carvalho, J.F.; Alonso-Juarez, M. The Link between Gut Dysbiosis and Neuroinflammation in Parkinson’s Disease. *Neuroscience* **2020**, *432*, 160–173. [[CrossRef](#)] [[PubMed](#)]
26. Wasser, C.I.; Mercieca, E.-C.; Kong, G.; Hannan, A.J.; McKeown, S.J.; Glikmann-Johnston, Y.; Stout, J.C. Gut dysbiosis in Huntington’s disease: Associations among gut microbiota, cognitive performance and clinical outcomes. *Brain Commun.* **2020**, *2*, 110. [[CrossRef](#)]
27. Miyake, S.; Kim, S.; Suda, W.; Oshima, K.; Nakamura, M.; Matsuoka, T.; Chihara, N.; Tomita, A.; Sato, W.; Kim, S.-W.; et al. Dysbiosis in the Gut Microbiota of Patients with Multiple Sclerosis, with a Striking Depletion of Species Belonging to Clostridia XIVa and IV Clusters. *PLoS ONE* **2015**, *10*, e0137429. [[CrossRef](#)]
28. Anwar, H.; Irfan, S.; Hussain, G.; Naeem Faisal, M.; Muzaffar, H.; Mustafa, I.; Mukhtar, I.; Malik, S.; Irfan Ullah, M. Gut Microbiome: A New Organ System in Body. In *Parasitology and Microbiology Research*; Antonio Bastidas Pacheco, G., Ali Kamboh, A., Eds.; IntechOpen: London, UK, 2020; ISBN 978-1-78985-901-0.
29. Jandhyala, S.M.; Talukdar, R.; Subramanyam, C.; Vuyyuru, H.; Sasikala, M.; Nageshwar Reddy, D. Role of the normal gut microbiota. *World J. Gastroenterol.* **2015**, *21*, 8787–8803. [[CrossRef](#)]
30. Boziki, M.K.; Mentis, A.-F.A.; Shumilina, M.; Makshakov, G.; Evdoshenko, E.; Grigoriadis, N. COVID-19 Immunopathology and the Central Nervous System: Implication for Multiple Sclerosis and Other Autoimmune Diseases with Associated Demyelination. *Brain Sci.* **2020**, *10*, 345. [[CrossRef](#)]
31. Magne, F.; Gotteland, M.; Gauthier, L.; Zazueta, A.; Poeso, S.; Navarrete, P.; Balamurugan, R. The *Firmicutes/Bacteroidetes* Ratio: A Relevant Marker of Gut Dysbiosis in Obese Patients? *Nutrients* **2020**, *12*, 1474. [[CrossRef](#)]
32. Johnson, E.L.; Heaver, S.L.; Walters, W.A.; Ley, R.E. Microbiome and metabolic disease: Revisiting the bacterial phylum *Bacteroidetes*. *Klin. Wochenschr.* **2016**, *95*, 1–8. [[CrossRef](#)]
33. Mariat, D.; Firmesse, O.; Levenez, F.; Guimaraes, V.D.; Sokol, H.; Dore, J.; Corthier, G.; Furet, J.-P. The *Firmicutes/Bacteroidetes* ratio of the human microbiota changes with age. *BMC Microbiol.* **2009**, *9*, 123. [[CrossRef](#)]
34. Rinninella, E.; Raoul, P.; Cintoni, M.; Franceschi, F.; Miggiano, G.A.D.; Gasbarrini, A.; Mele, M.C. What Is the Healthy Gut Microbiota Composition? A Changing Ecosystem across Age, Environment, Diet, and Diseases. *Microorganisms* **2019**, *7*, 14. [[CrossRef](#)]
35. Devaraj, S.; Hemarajata, P.; Versalovic, J. The Human Gut Microbiome and Body Metabolism: Implications for Obesity and Diabetes. *Clin. Chem.* **2013**, *59*, 617–628. [[CrossRef](#)]
36. Deleu, S.; Machiels, K.; Raes, J.; Verbeke, K.; Vermeire, S. Short chain fatty acids and its producing organisms: An overlooked therapy for IBD? *eBioMedicine* **2021**, *66*, 103293. [[CrossRef](#)]
37. Singh, R.K.; Chang, H.-W.; Yan, D.; Lee, K.M.; Ucmak, D.; Wong, K.; Abrouk, M.; Farahnik, B.; Nakamura, M.; Zhu, T.H.; et al. Influence of diet on the gut microbiome and implications for human health. *J. Transl. Med.* **2017**, *15*, 73. [[CrossRef](#)]
38. Cantarel, B.L.; Lombard, V.; Henrissat, B. Complex Carbohydrate Utilization by the Healthy Human Microbiome. *PLoS ONE* **2012**, *7*, e28742. [[CrossRef](#)]
39. Velagapudi, V.R.; Hezaveh, R.; Reigstad, C.S.; Gopalacharyulu, P.; Yetukuri, L.; Islam, S.; Felin, J.; Perkins, R.; Borén, J.; Orešič, M.; et al. The gut microbiota modulates host energy and lipid metabolism in mice. *J. Lipid Res.* **2010**, *51*, 1101–1112. [[CrossRef](#)]

40. Shreiner, A.B.; Kao, J.Y.; Young, V.B. The gut microbiome in health and in disease. *Curr. Opin. Gastroenterol.* **2015**, *31*, 69–75. [[CrossRef](#)]
41. Sidhu, H.; Hoppe, B.; Hesse, A.; Tenbrock, K.; Bromme, S.; Rietschel, E.; Peck, A.B. Absence of *Oxalobacter formigenes* in cystic fibrosis patients: A risk factor for hyperoxaluria. *Lancet* **1998**, *352*, 1026–1029. [[CrossRef](#)]
42. Gonzalez-Regueiro, J.A.; Moreno-Castaneda, L.; Uribe, M.; Chavez-Tapia, N.C. The Role of Bile Acids in Glucose Metabolism and Their Relation with Diabetes. *Ann. Hepatol.* **2017**, *16*, 16–21. [[CrossRef](#)] [[PubMed](#)]
43. Hofmann, A.F. The Continuing Importance of Bile Acids in Liver and Intestinal Disease. *Arch. Intern. Med.* **1999**, *159*, 2647–2658. [[CrossRef](#)] [[PubMed](#)]
44. Kakiyama, G.; Muto, A.; Takei, H.; Nittono, H.; Murai, T.; Kurosawa, T.; Hofmann, A.F.; Pandak, W.M.; Bajaj, J.S. A simple and accurate HPLC method for fecal bile acid profile in healthy and cirrhotic subjects: Validation by GC-MS and LC-MS. *J. Lipid Res.* **2014**, *55*, 978–990. [[CrossRef](#)] [[PubMed](#)]
45. Guzior, D.V.; Quinn, R.A. Review: Microbial transformations of human bile acids. *Microbiome* **2021**, *9*, 140. [[CrossRef](#)]
46. Quinn, R.A.; Melnik, A.V.; Vrbanac, A.; Fu, T.; Patras, K.A.; Christy, M.P.; Bodai, Z.; Belda-Ferre, P.; Tripathi, A.; Chung, L.K.; et al. Global chemical effects of the microbiome include new bile-acid conjugations. *Nature* **2020**, *579*, 123–129. [[CrossRef](#)]
47. Gu, Q.; Li, P. Biosynthesis of Vitamins by Probiotic Bacteria. In *Probiotics and Prebiotics in Human Nutrition and Health*; Rao, V., Rao, L.G., Eds.; IntechOpen: London, UK, 2016; ISBN 978-953-51-2475-7.
48. Engevik, M.; Morra, C.N.; Röth, D.; Engevik, K.; Spinler, J.K.; Devaraj, S.; Crawford, S.E.; Estes, M.K.; Kalkum, M.; Versalovic, J. Microbial Metabolic Capacity for Intestinal Folate Production and Modulation of Host Folate Receptors. *Front. Microbiol.* **2019**, *10*, 2305. [[CrossRef](#)]
49. Magnusdottir, S.; Ravcheev, D.; De Crécy-Lagard, V.; Thiele, I. Systematic genome assessment of B-vitamin biosynthesis suggests co-operation among gut microbes. *Front. Genet.* **2015**, *6*, 148. [[CrossRef](#)]
50. De Assis, D.A.; Matte, C.; Aschidamini, B.; Rodrigues, E.; Ayub, M.A.Z. Biosynthesis of vitamin B12 by *Propionibacterium freudenreichii* subsp. *shermanii* ATCC 13673 using liquid acid protein residue of soybean as culture medium. *Biotechnol. Prog.* **2020**, *36*, e3011. [[CrossRef](#)]
51. LeBlanc, J.G.; Milani, C.; de Giori, G.S.; Sesma, F.; van Sinderen, D.; Ventura, M. Bacteria as vitamin suppliers to their host: A gut microbiota perspective. *Curr. Opin. Biotechnol.* **2012**, *24*, 160–168. [[CrossRef](#)]
52. Zheng, D.; Liwinski, T.; Elinav, E. Interaction between microbiota and immunity in health and disease. *Cell Res.* **2020**, *30*, 492–506. [[CrossRef](#)]
53. Shrestha, U.K. Immunology of the gut and oral tolerance. *J. Adv. Intern. Med.* **2015**, *4*, 16–24. [[CrossRef](#)]
54. Geuking, M.B.; Cahenzli, J.; Lawson, M.; Ng, D.C.; Slack, E.; Hapfelmeier, S.; McCoy, K.D.; Macpherson, A.J. Intestinal Bacterial Colonization Induces Mutualistic Regulatory T Cell Responses. *Immunity* **2011**, *34*, 794–806. [[CrossRef](#)]
55. Round, J.L.; Lee, S.M.; Li, J.; Tran, G.; Jabri, B.; Chatila, T.A.; Mazmanian, S.K. The Toll-Like Receptor 2 Pathway Establishes Colonization by a Commensal of the Human Microbiota. *Science* **2011**, *332*, 974–977. [[CrossRef](#)]
56. Van der Hee, B.; Wells, J.M. Microbial Regulation of Host Physiology by Short-chain Fatty Acids. *Trends Microbiol.* **2021**, *29*, 700–712. [[CrossRef](#)]
57. Smith, P.M.; Howitt, M.R.; Panikov, N.; Michaud, M.; Gallini, C.A.; Bohlooly-Y, M.; Glickman, J.N.; Garrett, W.S. The Microbial Metabolites, Short-Chain Fatty Acids, Regulate Colonic Treg Cell Homeostasis. *Science* **2013**, *341*, 569–573. [[CrossRef](#)]
58. White, J.R.; Daurios-Singorenko, P.; Hong, J.; Vanholsbeeck, F.; Phillips, A.; Swift, S. The role of host molecules in communication with the resident and pathogenic microbiota: A review. *Med. Microecol.* **2020**, *4*, 100005. [[CrossRef](#)]
59. Reyman, M.; Van Houten, M.A.; Van Baarle, D.; Bosch, A.A.T.M.; Man, W.H.; Chu, M.L.J.N.; Arp, K.; Watson, R.L.; Sanders, E.A.M.; Fuentes, S.; et al. Impact of delivery mode-associated gut microbiota dynamics on health in the first year of life. *Nat. Commun.* **2019**, *10*, 4997. [[CrossRef](#)]
60. Neu, J.; Rushing, J. Cesarean Versus Vaginal Delivery: Long-term Infant Outcomes and the Hygiene Hypothesis. *Clin. Perinatol.* **2011**, *38*, 321–331. [[CrossRef](#)]
61. Grönlund, M.-M.; Lehtonen, O.-P.; Eerola, E.; Kero, P. Fecal Microflora in Healthy Infants Born by Different Methods of Delivery: Permanent Changes in Intestinal Flora After Cesarean Delivery. *J. Pediatr. Gastroenterol. Nutr.* **1999**, *28*, 19–25. [[CrossRef](#)]
62. Coelho, G.D.P.; Ayres, L.F.A.; Barreto, D.S.; Henriques, B.D.; Prado, M.R.M.C.; Dos Passos, C.M. Acquisition of microbiota according to the type of birth: An integrative review. *Rev. Lat.-Am. Enferm.* **2021**, *29*, e3446. [[CrossRef](#)]
63. Dominguez-Bello, M.G.; Costello, E.K.; Contreras, M.; Magris, M.; Hidalgo, G.; Fierer, N.; Knight, R. Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc. Natl. Acad. Sci. USA* **2010**, *107*, 11971–11975. [[CrossRef](#)] [[PubMed](#)]
64. Boix-Amorós, A.; Collado, M.C.; Mira, A. Relationship between Milk Microbiota, Bacterial Load, Macronutrients, and Human Cells during Lactation. *Front. Microbiol.* **2016**, *7*, 492. [[CrossRef](#)] [[PubMed](#)]
65. Rodríguez, J.M. The Origin of Human Milk Bacteria: Is There a Bacterial Entero-Mammary Pathway during Late Pregnancy and Lactation? *Adv. Nutr.* **2014**, *5*, 779–784. [[CrossRef](#)] [[PubMed](#)]
66. Marin-Gómez, W.; Grande, M.J.; Pérez-Pulido, R.; Galvez, A.; Lucas, R. Changes in the Bacterial Diversity of Human Milk during Late Lactation Period (Weeks 21 to 48). *Foods* **2020**, *9*, 1184. [[CrossRef](#)] [[PubMed](#)]
67. Obermajer, T.; Pogačić, T. Commentary: Relationship between Milk Microbiota, Bacterial Load, Macronutrients, and Human Cells during Lactation. *Front. Microbiol.* **2016**, *7*, 1281. [[CrossRef](#)]

68. Nuriel-Ohayon, M.; Neuman, H.; Koren, O. Microbial Changes during Pregnancy, Birth, and Infancy. *Front. Microbiol.* **2016**, *7*, 1031. [\[CrossRef\]](#)
69. Toscano, M.; De Grandi, R.; Grossi, E.; Drago, L. Role of the Human Breast Milk-Associated Microbiota on the Newborns' Immune System: A Mini Review. *Front. Microbiol.* **2017**, *8*, 2100. [\[CrossRef\]](#)
70. Panagos, P.G.; Vishwanathan, R.; Penfield-Cyr, A.; Matthan, N.R.; Shivappa, N.; Wirth, M.D.; Hebert, J.R.; Sen, S. Breastmilk from obese mothers has pro-inflammatory properties and decreased neuroprotective factors. *J. Perinatol.* **2016**, *36*, 284–290. [\[CrossRef\]](#)
71. Wen, L.; Duffy, A. Factors Influencing the Gut Microbiota, Inflammation, and Type 2 Diabetes. *J. Nutr.* **2017**, *147*, 1468S–1475S. [\[CrossRef\]](#)
72. Klingensmith, N.J.; Coopersmith, C.M. The Gut as the Motor of Multiple Organ Dysfunction in Critical Illness. *Crit. Care Clin.* **2016**, *32*, 203–212. [\[CrossRef\]](#)
73. Becattini, S.; Taur, Y.; Pamer, E.G. Antibiotic-Induced Changes in the Intestinal Microbiota and Disease. *Trends Mol. Med.* **2016**, *22*, 458–478. [\[CrossRef\]](#)
74. Palleja, A.; Mikkelsen, K.H.; Forslund, S.K.; Kashani, A.; Allin, K.H.; Nielsen, T.; Hansen, T.H.; Liang, S.; Feng, Q.; Zhang, C.; et al. Recovery of gut microbiota of healthy adults following antibiotic exposure. *Nat. Microbiol.* **2018**, *3*, 1255–1265. [\[CrossRef\]](#)
75. Hibbing, M.E.; Fuqua, C.; Parsek, M.R.; Peterson, S.B. Bacterial competition: Surviving and thriving in the microbial jungle. *Nat. Rev. Microbiol.* **2010**, *8*, 15–25. [\[CrossRef\]](#)
76. Hehemann, J.-H.; Correc, G.; Barbeyron, T.; Helbert, W.; Czjzek, M.; Michel, G. Transfer of carbohydrate-active enzymes from marine bacteria to Japanese gut microbiota. *Nature* **2010**, *464*, 908–912. [\[CrossRef\]](#)
77. Li, M.; Chen, W.-D.; Wang, Y.-D. The roles of the gut microbiota-miRNA interaction in the host pathophysiology. *Mol. Med.* **2020**, *26*, 101. [\[CrossRef\]](#)
78. Djuranovic, S.; Nahvi, A.; Green, R. miRNA-Mediated Gene Silencing by Translational Repression Followed by mRNA Deadenylation and Decay. *Science* **2012**, *336*, 237–240. [\[CrossRef\]](#)
79. Wahid, F.; Shehzad, A.; Khan, T.; Kim, Y.Y. MicroRNAs: Synthesis, mechanism, function, and recent clinical trials. *Biochim. Biophys. Acta* **2010**, *1803*, 1231–1243. [\[CrossRef\]](#)
80. Liu, S.; da Cunha, A.P.; Rezende, R.M.; Cialic, R.; Wei, Z.; Bry, L.; Comstock, L.E.; Gandhi, R.; Weiner, H.L. The Host Shapes the Gut Microbiota via Fecal MicroRNA. *Cell Host Microbe* **2016**, *19*, 32–43. [\[CrossRef\]](#)
81. Link, A.; Becker, V.; Goel, A.; Wex, T.; Malfertheiner, P. Feasibility of Fecal MicroRNAs as Novel Biomarkers for Pancreatic Cancer. *PLoS ONE* **2012**, *7*, e42933. [\[CrossRef\]](#)
82. Wortelboer, K.; Bakker, G.J.; Winkelmeijer, M.; van Riel, N.; Levin, E.; Nieuwdorp, M.; Herrema, H.; Davids, M. Fecal microbiota transplantation as tool to study the interrelation between microbiota composition and miRNA expression. *Microbiol. Res.* **2022**, *257*, 126972. [\[CrossRef\]](#)
83. Viennois, E.; Chassaing, B.; Tahsin, A.; Pujada, A.; Wang, L.; Gewirtz, A.T.; Merlin, D. Host-derived fecal microRNAs can indicate gut microbiota healthiness and ability to induce inflammation. *Theranostics* **2019**, *9*, 4542–4557. [\[CrossRef\]](#) [\[PubMed\]](#)
84. Schächtle, M.A.; Rosshart, S.P. The Microbiota-Gut-Brain Axis in Health and Disease and Its Implications for Translational Research. *Front. Cell. Neurosci.* **2021**, *15*, 698172. [\[CrossRef\]](#) [\[PubMed\]](#)
85. Cryan, J.F.; O'Riordan, K.J.; Cowan, C.S.M.; Sandhu, K.V.; Bastiaanssen, T.F.S.; Boehme, M.; Codagnone, M.G.; Cusotto, S.; Fulling, C.; Golubeva, A.V.; et al. The Microbiota-Gut-Brain Axis. *Physiol. Rev.* **2019**, *99*, 1877–2013. [\[CrossRef\]](#) [\[PubMed\]](#)
86. Wu, K.; Taki, Y.; Sato, K.; Kinomura, S.; Goto, R.; Okada, K.; Kawashima, R.; He, Y.; Evans, A.C.; Fukuda, H. Age-related changes in topological organization of structural brain networks in healthy individuals. *Hum. Brain Mapp.* **2011**, *33*, 552–568. [\[CrossRef\]](#) [\[PubMed\]](#)
87. De la Fuente-Nunez, C.; Meneguetti, B.T.; Franco, O.L.; Lu, T.K. Neuromicrobiology: How Microbes Influence the Brain. *ACS Chem. Neurosci.* **2017**, *9*, 141–150. [\[CrossRef\]](#)
88. Ignatova, V. Influence of Gut Microbiota on Behavior and Its Disturbances. In *Behavioral Neuroscience*; Palermo, S., Morese, R., Eds.; IntechOpen: London, UK, 2019; ISBN 978-1-78984-051-3.
89. Al-Asmakh, M.; Anuar, F.; Zadjali, F.; Rafter, J.; Pettersson, S. Gut microbial communities modulating brain development and function. *Gut Microbes* **2012**, *3*, 366–373. [\[CrossRef\]](#)
90. Ogbonnaya, E.S.; Clarke, G.; Shanahan, F.; Dinan, T.G.; Cryan, J.F.; O'Leary, O.F. Adult Hippocampal Neurogenesis Is Regulated by the Microbiome. *Biol. Psychiatry* **2015**, *78*, e7–e9. [\[CrossRef\]](#)
91. Erny, D.; Hrabě de Angelis, A.L.; Jaitin, D.; Wieghofer, P.; Staszewski, O.; David, E.; Keren-Shaul, H.; Mhlahkoiv, T.; Jakobshagen, K.; Buch, T.; et al. Host microbiota constantly control maturation and function of microglia in the CNS. *Nat. Neurosci.* **2015**, *18*, 965–977. [\[CrossRef\]](#)
92. Carabotti, M.; Scirocco, A.; Maselli, M.A.; Severi, C. The gut-brain axis: Interactions between enteric microbiota, central and enteric nervous systems. *Ann. Gastroenterol.* **2015**, *28*, 203–209.
93. Martin, C.R.; Osadchiy, V.; Kalani, A.; Mayer, E.A. The Brain-Gut-Microbiome Axis. *Cell. Mol. Gastroenterol. Hepatol.* **2018**, *6*, 133–148. [\[CrossRef\]](#)
94. Indari, O.; Chandramohanadas, R.; Jha, H.C. Epstein-Barr virus infection modulates blood-brain barrier cells and its co-infection with *Plasmodium falciparum* induces RBC adhesion. *Pathog. Dis.* **2020**, *79*, ftaa080. [\[CrossRef\]](#)
95. Kashyap, D.; Baral, B.; Jakhmola, S.; Singh, A.K.; Jha, H.C. *Helicobacter pylori* and Epstein-Barr Virus Coinfection Stimulates Aggressiveness in Gastric Cancer through the Regulation of Gankyrin. *mSphere* **2021**, *6*, e00751-21. [\[CrossRef\]](#)

96. Singh, S.; Jha, H.C. Status of Epstein-Barr Virus Coinfection with *Helicobacter pylori* in Gastric Cancer. *J. Oncol.* **2017**, *2017*, 1–17. [[CrossRef](#)]
97. Montiel-Castro, A.J.; González-Cervantes, R.M.; Ebravo-Ruiseco, G.; Epacheco-Lopez, G. The microbiota-gut-brain axis: Neurobehavioral correlates, health and sociality. *Front. Integr. Neurosci.* **2013**, *7*, 70. [[CrossRef](#)]
98. Mazzoli, R.; Pessione, E. The Neuro-endocrinological Role of Microbial Glutamate and GABA Signaling. *Front. Microbiol.* **2016**, *7*, 1934. [[CrossRef](#)]
99. Chen, Y.; Xu, J.; Chen, Y. Regulation of Neurotransmitters by the Gut Microbiota and Effects on Cognition in Neurological Disorders. *Nutrients* **2021**, *13*, 2099. [[CrossRef](#)]
100. Caspani, G.; Swann, J. Small talk: Microbial metabolites involved in the signaling from microbiota to brain. *Curr. Opin. Pharmacol.* **2019**, *48*, 99–106. [[CrossRef](#)]
101. Breit, S.; Kupferberg, A.; Rogler, G.; Hasler, G. Vagus Nerve as Modulator of the Brain–Gut Axis in Psychiatric and Inflammatory Disorders. *Front. Psychiatry* **2018**, *9*, 44. [[CrossRef](#)]
102. McCorry, L.K. Physiology of the Autonomic Nervous System. *Am. J. Pharm. Educ.* **2007**, *71*, 78. [[CrossRef](#)]
103. Cunningham, D.J.C.; Robbins, P.A.; Wolff, C.B. Integration of Respiratory Responses to Changes in Alveolar Partial Pressures of CO₂ and O₂ and in Arterial pH. In *Comprehensive Physiology*; Terjung, R., Ed.; Wiley: Hoboken, NJ, USA, 1986; pp. 475–528, ISBN 978-0-470-65071-4.
104. Costa, M.; Brookes, S.J.H.; Hennig, G.W. Anatomy and physiology of the enteric nervous system. *Gut* **2000**, *47*, 15iv–19. [[CrossRef](#)]
105. Fülling, C.; Dinan, T.G.; Cryan, J.F. Gut Microbe to Brain Signaling: What Happens in Vagus *Neuron* **2019**, *101*, 998–1002. [[CrossRef](#)] [[PubMed](#)]
106. Rangon, C.-M.; Niezgoda, A. Understanding the Pivotal Role of the Vagus Nerve in Health from Pandemics. *Bioengineering* **2022**, *9*, 352. [[CrossRef](#)] [[PubMed](#)]
107. Bonaz, B.; Bazin, T.; Pellissier, S. The Vagus Nerve at the Interface of the Microbiota-Gut-Brain Axis. *Front. Neurosci.* **2018**, *12*, 49. [[CrossRef](#)]
108. Yu, K.B.; Hsiao, E.Y. Roles for the gut microbiota in regulating neuronal feeding circuits. *J. Clin. Investig.* **2021**, *131*, e143772. [[CrossRef](#)] [[PubMed](#)]
109. Forsythe, P.; Bienenstock, J.; Kunze, W.A. Vagal Pathways for Microbiome-Brain-Gut Axis Communication. In *Microbial Endocrinology: The Microbiota-Gut-Brain Axis in Health and Disease*; Advances in Experimental Medicine and Biology; Lyte, M., Cryan, J.F., Eds.; Springer: New York, NY, USA, 2014; Volume 817, pp. 115–133, ISBN 978-1-4939-0896-7.
110. Goehler, L.E.; Gaykema, R.P.; Opitz, N.; Reddaway, R.; Badr, N.; Lyte, M. Activation in vagal afferents and central autonomic pathways: Early responses to intestinal infection with *Campylobacter jejuni*. *Brain Behav. Immun.* **2005**, *19*, 334–344. [[CrossRef](#)] [[PubMed](#)]
111. Bercik, P.; Park, A.J.; Sinclair, D.; Khoshdel, A.; Lu, J.; Huang, X.; Deng, Y.; Blennerhassett, P.A.; Fahnestock, M.; Moine, D.; et al. The anxiolytic effect of *Bifidobacterium longum* NCC3001 involves vagal pathways for gut-brain communication. *Neurogastroenterol. Motil.* **2011**, *23*, 1132–1139. [[CrossRef](#)]
112. Alim, E.; Gülekon, I.N.; Atalar, K.; Bahçelioğlu, M. Enteric nervous system, gut-brain connection and related neurodevelopmental disorders. *Anatomy* **2020**, *14*, 61–67. [[CrossRef](#)]
113. Goehler, L.E.; Park, S.M.; Opitz, N.; Lyte, M.; Gaykema, R.P. *Campylobacter jejuni* infection increases anxiety-like behavior in the holeboard: Possible anatomical substrates for viscerosensory modulation of exploratory behavior. *Brain Behav. Immun.* **2008**, *22*, 354–366. [[CrossRef](#)]
114. Browning, K.N.; Verheijden, S.; Boeckstaens, G.E. The Vagus Nerve in Appetite Regulation, Mood, and Intestinal Inflammation. *Gastroenterology* **2017**, *152*, 730–744. [[CrossRef](#)]
115. Silva, Y.P.; Bernardi, A.; Frozza, R.L. The Role of Short-Chain Fatty Acids From Gut Microbiota in Gut-Brain Communication. *Front. Endocrinol. (Lausanne)* **2020**, *11*, 25. [[CrossRef](#)]
116. Ma, Q.; Xing, C.; Long, W.; Wang, H.Y.; Liu, Q.; Wang, R.-F. Impact of microbiota on central nervous system and neurological diseases: The gut-brain axis. *J. Neuroinflamm.* **2019**, *16*, 53. [[CrossRef](#)]
117. Bravo, J.A.; Forsythe, P.; Chew, M.V.; Escaravage, E.; Savignac, H.M.; Dinan, T.G.; Bienenstock, J.; Cryan, J.F. Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 16050–16055. [[CrossRef](#)]
118. Sarubbo, F.; Cavallucci, V.; Pani, G. The Influence of Gut Microbiota on Neurogenesis: Evidence and Hopes. *Cells* **2022**, *11*, 382. [[CrossRef](#)]
119. Ahmed, H.; Leyrolle, Q.; Koistinen, V.; Kärkkäinen, O.; Layé, S.; Delzenne, N.; Hanhineva, K. Microbiota-derived metabolites as drivers of gut–brain communication. *Gut Microbes* **2022**, *14*, 2102878. [[CrossRef](#)]
120. Yarandi, S.S.; Peterson, D.A.; Treisman, G.J.; Moran, T.H.; Pasricha, P.J. Modulatory Effects of Gut Microbiota on the Central Nervous System: How Gut Could Play a Role in Neuropsychiatric Health and Diseases. *J. Neurogastroenterol. Motil.* **2016**, *22*, 201–212. [[CrossRef](#)]
121. Rhee, S.H.; Pothoulakis, C.; Mayer, E.A. Principles and clinical implications of the brain–gut–enteric microbiota axis. *Nat. Rev. Gastroenterol. Hepatol.* **2009**, *6*, 306–314. [[CrossRef](#)]
122. Talham, G.L.; Jiang, H.-Q.; Bos, N.A.; Cebra, J.J. Segmented Filamentous Bacteria Are Potent Stimuli of a Physiologically Normal State of the Murine Gut Mucosal Immune System. *Infect. Immun.* **1999**, *67*, 1992–2000. [[CrossRef](#)]

123. McKernan, D.P. Toll-like receptors and immune cell crosstalk in the intestinal epithelium. *AIMS Allergy Immunol.* **2019**, *3*, 13–31. [[CrossRef](#)]
124. Vijay, K. Toll-like receptors in immunity and inflammatory diseases: Past, present, and future. *Int. Immunopharmacol.* **2018**, *59*, 391–412. [[CrossRef](#)]
125. Frosali, S.; Pagliari, D.; Gambassi, G.; Landolfi, R.; Pandolfi, F.; Cianci, R. How the Intricate Interaction among Toll-Like Receptors, Microbiota, and Intestinal Immunity Can Influence Gastrointestinal Pathology. *J. Immunol. Res.* **2015**, *2015*, 1–12. [[CrossRef](#)]
126. Alhasson, F.; Das, S.; Seth, R.; Dattaroy, D.; Chandrashekar, V.; Ryan, C.N.; Chan, L.S.; Testerman, T.; Burch, J.; Hofseth, L.J.; et al. Altered gut microbiome in a mouse model of Gulf War Illness causes neuroinflammation and intestinal injury via leaky gut and TLR4 activation. *PLoS ONE* **2017**, *12*, e0172914. [[CrossRef](#)]
127. Holzer, P.; Farzi, A. Neuropeptides and the Microbiota-Gut-Brain Axis. In *Microbial Endocrinology: The Microbiota-Gut-Brain Axis in Health and Disease*; Advances in Experimental Medicine and Biology; Lyte, M., Cryan, J.F., Eds.; Springer: New York, NY, USA, 2014; Volume 817, pp. 195–219, ISBN 978-1-4939-0896-7.
128. Guinane, C.M.; Cotter, P.D. Role of the gut microbiota in health and chronic gastrointestinal disease: Understanding a hidden metabolic organ. *Ther. Adv. Gastroenterol.* **2013**, *6*, 295–308. [[CrossRef](#)] [[PubMed](#)]
129. Zhang, Y.-J.; Li, S.; Gan, R.-Y.; Zhou, T.; Xu, D.-P.; Li, H.-B. Impacts of Gut Bacteria on Human Health and Diseases. *Int. J. Mol. Sci.* **2015**, *16*, 7493–7519. [[CrossRef](#)] [[PubMed](#)]
130. Sharon, G.; Sampson, T.R.; Geschwind, D.H.; Mazmanian, S.K. The Central Nervous System and the Gut Microbiome. *Cell* **2016**, *167*, 915–932. [[CrossRef](#)] [[PubMed](#)]
131. Faruqui, N.A.; Prium, D.H.; Mowna, S.A.; Ullah, A.; Araf, Y.; Sarkar, B.; Zohora, U.S.; Rahman, M.S. Gut microorganisms and neurological disease perspectives. *Futur. Neurol.* **2021**, *16*, FNL53. [[CrossRef](#)]
132. Lehner, C.; Gehwolf, R.; Tempfer, H.; Krizbai, I.; Hennig, B.; Bauer, H.-C.; Bauer, H. Oxidative Stress and Blood-Brain Barrier Dysfunction Under Particular Consideration of Matrix Metalloproteinases. *Antioxid. Redox Signal.* **2011**, *15*, 1305–1323. [[CrossRef](#)]
133. Liebner, S.; Dijkhuizen, R.M.; Reiss, Y.; Plate, K.H.; Agalliu, D.; Constantin, G. Functional morphology of the blood-brain barrier in health and disease. *Acta Neuropathol.* **2018**, *135*, 311–336. [[CrossRef](#)]
134. Mu, C.; Yang, Y.; Zhu, W. Gut Microbiota: The Brain Peacekeeper. *Front. Microbiol.* **2016**, *7*, 345. [[CrossRef](#)]
135. Jena, A.; Montoya, C.A.; Mullaney, J.A.; Dilger, R.N.; Young, W.; McNabb, W.C.; Roy, N.C. Gut-Brain Axis in the Early Postnatal Years of Life: A Developmental Perspective. *Front. Integr. Neurosci.* **2020**, *14*, 44. [[CrossRef](#)]
136. Braniste, V.; Al-Asmakh, M.; Kowal, C.; Anuar, F.; Abbaspour, A.; Tóth, M.; Korecka, A.; Bakocevic, N.; Ng, L.G.; Kundu, P.; et al. The gut microbiota influences blood-brain barrier permeability in mice. *Sci. Transl. Med.* **2014**, *6*, 263ra158. [[CrossRef](#)]
137. Parker, A.; Fonseca, S.; Carding, S.R. Gut microbes and metabolites as modulators of blood-brain barrier integrity and brain health. *Gut Microbes* **2019**, *11*, 135–157. [[CrossRef](#)]
138. Liu, W.-Y.; Wang, Z.-B.; Zhang, L.-C.; Wei, X.; Li, L. Tight Junction in Blood-Brain Barrier: An Overview of Structure, Regulation, and Regulator Substances. *CNS Neurosci. Ther.* **2012**, *18*, 609–615. [[CrossRef](#)]
139. Tang, W.; Zhu, H.; Feng, Y.; Guo, R.; Wan, D. The Impact of Gut Microbiota Disorders on the Blood-Brain Barrier. *Infect. Drug Resist.* **2020**, *13*, 3351–3363. [[CrossRef](#)]
140. Kabouridis, P.S.; Lasrado, R.; McCallum, S.; Chng, S.H.; Snippert, H.J.; Clevers, H.; Pettersson, S.; Pachnis, V. The gut microbiota keeps enteric glial cells on the move; prospective roles of the gut epithelium and immune system. *Gut Microbes* **2015**, *6*, 398–403. [[CrossRef](#)]
141. Kabouridis, P.S.; Lasrado, R.; McCallum, S.; Chng, S.H.; Snippert, H.J.; Clevers, H.; Pettersson, S.; Pachnis, V. Microbiota Controls the Homeostasis of Glial Cells in the Gut Lamina Propria. *Neuron* **2015**, *85*, 289–295. [[CrossRef](#)]
142. De Vadder, F.; Gasset, E.; Holm, L.M.; Karsenty, G.; MacPherson, A.J.; Olofsson, L.E.; Bäckhed, F. Gut microbiota regulates maturation of the adult enteric nervous system via enteric serotonin networks. *Proc. Natl. Acad. Sci. USA* **2018**, *115*, 6458–6463. [[CrossRef](#)]
143. Grice, E.A.; Segre, J.A. The Human Microbiome: Our Second Genome. *Annu. Rev. Genom. Hum. Genet.* **2012**, *13*, 151–170. [[CrossRef](#)]
144. Clarke, S.; Murphy, E.F.; Nilaweera, K.; Ross, R.; Shanahan, F.; O’Toole, P.W.; Cotter, P.D. The gut microbiota and its relationship to diet and obesity. *Gut Microbes* **2012**, *3*, 186–202. [[CrossRef](#)]
145. Dhikav, V.; Anand, K.S. Hippocampus in health and disease: An overview. *Ann. Indian Acad. Neurol.* **2012**, *15*, 239–246. [[CrossRef](#)]
146. Luczynski, P.; Whelan, S.O.; O’Sullivan, C.; Clarke, G.; Shanahan, F.; Dinan, T.G.; Cryan, J.F. Adult microbiota-deficient mice have distinct dendritic morphological changes: Differential effects in the amygdala and hippocampus. *Eur. J. Neurosci.* **2016**, *44*, 2654–2666. [[CrossRef](#)]
147. Kawase, T.; Nagasawa, M.; Ikeda, H.; Yasuo, S.; Koga, Y.; Furuse, M. Gut microbiota of mice putatively modifies amino acid metabolism in the host brain. *Br. J. Nutr.* **2017**, *117*, 775–783. [[CrossRef](#)] [[PubMed](#)]
148. Swann, J.; Garcia-Perez, I.; Braniste, V.; Wilson, I.; Sidaway, J.; Nicholson, J.; Pettersson, S.; Holmes, E. Application of 1 H NMR spectroscopy to the metabolic phenotyping of rodent brain extracts: A metabolomic study of gut microbial influence on host brain metabolism. *J. Pharm. Biomed. Anal.* **2017**, *143*, 141–146. [[CrossRef](#)] [[PubMed](#)]
149. Tang, W.; Meng, Z.; Li, N.; Liu, Y.; Li, L.; Chen, D.; Yang, Y. Roles of Gut Microbiota in the Regulation of Hippocampal Plasticity, Inflammation, and Hippocampus-Dependent Behaviors. *Front. Cell. Infect. Microbiol.* **2021**, *10*, 611014. [[CrossRef](#)] [[PubMed](#)]

150. Luck, B.; Engevik, M.A.; Ganesh, B.; Lackey, E.P.; Lin, T.; Balderas, M.; Major, A.; Runge, J.; Luna, R.A.; Sillitoe, R.V.; et al. Bifidobacteria shape host neural circuits during postnatal development by promoting synapse formation and microglial function. *Sci. Rep.* **2020**, *10*, 7737. [[CrossRef](#)] [[PubMed](#)]
151. Björkholm, C.; Monteggia, L.M. BDNF—A Key Transducer of Antidepressant Effects. *Neuropharmacology* **2015**, *102*, 72–79. [[CrossRef](#)]
152. Sharma, P.; Kumar, A.; Singh, D. Dietary Flavonoids Interaction with CREB-BDNF Pathway: An Unconventional Approach for Comprehensive Management of Epilepsy. *Curr. Neuropharmacol.* **2019**, *17*, 1158–1175. [[CrossRef](#)]
153. Lv, M.; Yang, S.; Cai, L.; Qin, L.-Q.; Li, B.-Y.; Wan, Z. Effects of Quercetin Intervention on Cognition Function in APP/PS1 Mice was Affected by Vitamin D Status. *Mol. Nutr. Food Res.* **2018**, *62*, e1800621. [[CrossRef](#)]
154. Carding, S.; Verbeke, K.; Vipond, D.T.; Corfe, B.M.; Owen, L.J. Dysbiosis of the gut microbiota in disease. *Microb. Ecol. Health Dis.* **2015**, *26*, 26191. [[CrossRef](#)]
155. DeGruttola, A.K.; Low, D.; Mizoguchi, A.; Mizoguchi, E. Current Understanding of Dysbiosis in Disease in Human and Animal Models. *Inflamm. Bowel Dis.* **2016**, *22*, 1137–1150. [[CrossRef](#)]
156. Vogt, N.M.; Kerby, R.L.; Dill-McFarland, K.A.; Harding, S.J.; Merluzzi, A.P.; Johnson, S.C.; Carlsson, C.M.; Asthana, S.; Zetterberg, H.; Blennow, K.; et al. Gut microbiome alterations in Alzheimer’s disease. *Sci. Rep.* **2017**, *7*, 13537. [[CrossRef](#)]
157. Vascellari, S.; Palmas, V.; Melis, M.; Pisanu, S.; Cusano, R.; Uva, P.; Perra, D.; Madau, V.; Sarchioto, M.; Oppo, V.; et al. Gut Microbiota and Metabolome Alterations Associated with Parkinson’s Disease. *mSystems* **2020**, *5*, e00561-20. [[CrossRef](#)] [[PubMed](#)]
158. Tremlett, H.; Fadrosh, D.W.; Faruqi, A.A.; Zhu, F.; Hart, J.; Roalstad, S.; Graves, J.; Lynch, S.; Waubant, E.; Centers, T.U.N.O.P.M. Gut microbiota in early pediatric multiple sclerosis: A case–control study. *Eur. J. Neurol.* **2016**, *23*, 1308–1321. [[CrossRef](#)] [[PubMed](#)]
159. Sampson, T.R.; Mazmanian, S.K. Control of Brain Development, Function, and Behavior by the Microbiome. *Cell Host Microbe* **2015**, *17*, 565–576. [[CrossRef](#)] [[PubMed](#)]
160. Valdes, A.; Walter, J.; Segal, E.; Spector, T.D. Role of the gut microbiota in nutrition and health. *BMJ* **2018**, *361*, k2179. [[CrossRef](#)] [[PubMed](#)]
161. Yang, D.; Zhao, D.; Shah, S.Z.A.; Wu, W.; Lai, M.; Zhang, X.; Li, J.; Guan, Z.; Zhao, H.; Li, W.; et al. The Role of the Gut Microbiota in the Pathogenesis of Parkinson’s Disease. *Front. Neurol.* **2019**, *10*, 1155. [[CrossRef](#)]
162. Park, J.C.; Im, S.-H. The gut-immune-brain axis in neurodevelopment and neurological disorders. *Microbiome Res. Rep.* **2022**, *1*, 23. [[CrossRef](#)]
163. Sarkar, S.R.; Banerjee, S. Gut microbiota in neurodegenerative disorders. *J. Neuroimmunol.* **2019**, *328*, 98–104. [[CrossRef](#)]
164. Nogal, A.; Valdes, A.M.; Menni, C. The role of short-chain fatty acids in the interplay between gut microbiota and diet in cardio-metabolic health. *Gut Microbes* **2021**, *13*, 1897212. [[CrossRef](#)]
165. Mirzaei, R.; Bouzari, B.; Hosseini-Fard, S.R.; Mazaheri, M.; Ahmadyousefi, Y.; Abdi, M.; Jalalifar, S.; Karimitabar, Z.; Teimoori, A.; Keyvani, H.; et al. Role of microbiota-derived short-chain fatty acids in nervous system disorders. *Biomed. Pharmacother.* **2021**, *139*, 111661. [[CrossRef](#)]
166. Tran, S.M.-S.; Mohajeri, M.H. The Role of Gut Bacterial Metabolites in Brain Development, Aging and Disease. *Nutrients* **2021**, *13*, 732. [[CrossRef](#)]
167. Vogt, N.M.; Romano, K.A.; Darst, B.F.; Engelman, C.D.; Johnson, S.C.; Carlsson, C.M.; Asthana, S.; Blennow, K.; Zetterberg, H.; Bendlin, B.B.; et al. The gut microbiota-derived metabolite trimethylamine N-oxide is elevated in Alzheimer’s disease. *Alzheimer’s Res. Ther.* **2018**, *10*, 124. [[CrossRef](#)]
168. Koszewicz, M.; Jaroch, J.; Brzecka, A.; Ejma, M.; Budrewicz, S.; Mikhaleva, L.M.; Muresanu, C.; Schield, P.; Somasundaram, S.G.; Kirkland, C.E.; et al. Dysbiosis is one of the risk factor for stroke and cognitive impairment and potential target for treatment. *Pharmacol. Res.* **2020**, *164*, 105277. [[CrossRef](#)]
169. Marchesi, J.R.; Adams, D.H.; Fava, F.; Hermes, G.D.A.; Hirschfield, G.M.; Hold, G.; Quraishi, M.N.; Kinross, J.; Smidt, H.; Tuohy, K.M.; et al. The gut microbiota and host health: A new clinical frontier. *Gut* **2016**, *65*, 330–339. [[CrossRef](#)]
170. Chen, C.; Ahn, E.H.; Kang, S.S.; Liu, X.; Alam, A.; Ye, K. Gut dysbiosis contributes to amyloid pathology, associated with C/EBP β /AEP signaling activation in Alzheimer’s disease mouse model. *Sci. Adv.* **2020**, *6*, eaba0466. [[CrossRef](#)]
171. Gagliardi, A.; Totino, V.; Cacciotti, F.; Iebba, V.; Neroni, B.; Bonfiglio, G.; Trancassini, M.; Passariello, C.; Pantanella, F.; Schippa, S. Rebuilding the Gut Microbiota Ecosystem. *Int. J. Environ. Res. Public Health* **2018**, *15*, 1679. [[CrossRef](#)]
172. Markowiak, P.; Śliżewska, K. Effects of Probiotics, Prebiotics, and Synbiotics on Human Health. *Nutrients* **2017**, *9*, 1021. [[CrossRef](#)]
173. Hill, C.; Guarner, F.; Reid, G.; Gibson, G.R.; Merenstein, D.J.; Pot, B.; Morelli, L.; Canani, R.B.; Flint, H.J.; Salminen, S.; et al. Expert consensus document: The International Scientific Association for Probiotics and Prebiotics consensus statement on the scope and appropriate use of the term probiotic. *Nat. Rev. Gastroenterol. Hepatol.* **2014**, *11*, 506–514. [[CrossRef](#)]
174. Ochoa-Repáraz, J.; Mielcarz, D.W.; Ditrío, L.E.; Burroughs, A.R.; Begum-Haque, S.; Dasgupta, S.; Kasper, D.L.; Kasper, L.H. Central Nervous System Demyelinating Disease Protection by the Human Commensal *Bacteroides fragilis* Depends on Polysaccharide A Expression. *J. Immunol.* **2010**, *185*, 4101–4108. [[CrossRef](#)]
175. Huang, H.; Xu, H.; Luo, Q.; He, J.; Li, M.; Chen, H.; Tang, W.; Nie, Y.; Zhou, Y. Fecal microbiota transplantation to treat Parkinson’s disease with constipation. *Medicine* **2019**, *98*, e16163. [[CrossRef](#)]
176. König, R.S.; Albrich, W.C.; Kahlert, C.R.; Bahr, L.S.; Löber, U.; Vernazza, P.; Scheibenbogen, C.; Forslund, S.K. The Gut Microbiome in Myalgic Encephalomyelitis (ME)/Chronic Fatigue Syndrome (CFS). *Front. Immunol.* **2022**, *12*, 628741. [[CrossRef](#)]

177. Hsiao, E.Y.; McBride, S.W.; Hsien, S.; Sharon, G.; Hyde, E.R.; McCue, T.; Codelli, J.A.; Chow, J.; Reisman, S.E.; Petrosino, J.F.; et al. Microbiota Modulate Behavioral and Physiological Abnormalities Associated with Neurodevelopmental Disorders. *Cell* **2013**, *155*, 1451–1463. [[CrossRef](#)] [[PubMed](#)]
178. Azm, S.A.N.; Djazayeri, A.; Safa, M.; Azami, K.; Ahmadvand, B.; Sabbaghziarani, F.; Sharifzadeh, M.; Vafa, M. Lactobacilli and bifidobacteria ameliorate memory and learning deficits and oxidative stress in β -amyloid (1–42) injected rats. *Appl. Physiol. Nutr. Metab.* **2018**, *43*, 718–726. [[CrossRef](#)]
179. Jing, Y.; Yu, Y.; Bai, F.; Wang, L.; Yang, D.; Zhang, C.; Qin, C.; Yang, M.; Zhang, D.; Zhu, Y.; et al. Effect of fecal microbiota transplantation on neurological restoration in a spinal cord injury mouse model: Involvement of brain-gut axis. *Microbiome* **2021**, *9*, 1–21. [[CrossRef](#)] [[PubMed](#)]
180. Kim, C.-S.; Cha, L.; Sim, M.; Jung, S.; Chun, W.Y.; Baik, H.W.; Shin, D.-M. Probiotic Supplementation Improves Cognitive Function and Mood with Changes in Gut Microbiota in Community-Dwelling Older Adults: A Randomized, Double-Blind, Placebo-Controlled, Multicenter Trial. *J. Gerontol. Ser. A* **2020**, *76*, 32–40. [[CrossRef](#)] [[PubMed](#)]
181. Desbonnet, L.; Garrett, L.; Clarke, G.; Kiely, B.; Cryan, J.F.; Dinan, T.G. Effects of the probiotic *Bifidobacterium infantis* in the maternal separation model of depression. *Neuroscience* **2010**, *170*, 1179–1188. [[CrossRef](#)]
182. Du, D.; Tang, W.; Zhou, C.; Sun, X.; Wei, Z.; Zhong, J.; Huang, Z. Fecal Microbiota Transplantation Is a Promising Method to Restore Gut Microbiota Dysbiosis and Relieve Neurological Deficits after Traumatic Brain Injury. *Oxidative Med. Cell. Longev.* **2021**, *2021*, 1–21. [[CrossRef](#)]
183. Bonfili, L.; Cecarini, V.; Gogoi, O.; Berardi, S.; Scarpona, S.; Angeletti, M.; Rossi, G.; Eleuteri, A.M. Gut microbiota manipulation through probiotics oral administration restores glucose homeostasis in a mouse model of Alzheimer’s disease. *Neurobiol. Aging* **2019**, *87*, 35–43. [[CrossRef](#)]
184. Kechagia, M.; Basoulis, D.; Konstantopoulou, S.; Dimitriadi, D.; Gyftopoulou, K.; Skarmoutsou, N.; Fakiri, E.M. Health Benefits of Probiotics: A Review. *Int. Sch. Res. Not.* **2013**, *2013*, 1–7. [[CrossRef](#)]
185. Aponte, M.; Murru, N.; Shoukat, M. Therapeutic, Prophylactic, and Functional Use of Probiotics: A Current Perspective. *Front. Microbiol.* **2020**, *11*, 562048. [[CrossRef](#)]
186. Berg, R.D. The indigenous gastrointestinal microflora. *Trends Microbiol.* **1996**, *4*, 430–435. [[CrossRef](#)]
187. Liao, S.F.; Nyachoti, M. Using Probiotics to Improve Swine Gut Health and Nutrient Utilization. *Anim. Nutr.* **2017**, *3*, 331–343. [[CrossRef](#)]
188. Maldonado Galdeano, C.; Cazorla, S.I.; Lemme Dumit, J.M.; Vélez, E.; Perdigon, G. Beneficial Effects of Probiotic Consumption on the Immune System. *Ann. Nutr. Metab.* **2019**, *74*, 115–124. [[CrossRef](#)]
189. Fijan, S. Microorganisms with Claimed Probiotic Properties: An Overview of Recent Literature. *Int. J. Environ. Res. Public Health* **2014**, *11*, 4745–4767. [[CrossRef](#)]
190. Mani-López, E.; Palou, E.; López-Malo, A. Probiotic viability and storage stability of yogurts and fermented milks prepared with several mixtures of lactic acid bacteria. *J. Dairy Sci.* **2014**, *97*, 2578–2590. [[CrossRef](#)]
191. Amara, A.; Shibl, A. Role of Probiotics in health improvement, infection control and disease treatment and management. *Saudi Pharm. J.* **2013**, *23*, 107–114. [[CrossRef](#)]
192. Stavropoulou, E.; Bezirtzoglou, E. Probiotics in Medicine: A Long Debate. *Front. Immunol.* **2020**, *11*, 2192. [[CrossRef](#)]
193. Hemarajata, P.; Versalovic, J. Effects of probiotics on gut microbiota: Mechanisms of intestinal immunomodulation and neuro-modulation. *Ther. Adv. Gastroenterol.* **2012**, *6*, 39–51. [[CrossRef](#)]
194. Neef, A.; Sanz, Y. Future for probiotic science in functional food and dietary supplement development. *Curr. Opin. Clin. Nutr. Metab. Care* **2013**, *16*, 679–687. [[CrossRef](#)]
195. Khangwal, I.; Shukla, P. Prospecting prebiotics, innovative evaluation methods, and their health applications: A review. *3 Biotech* **2019**, *9*, 187. [[CrossRef](#)]
196. Davani-Davari, D.; Negahdaripour, M.; Karimzadeh, I.; Seifan, M.; Mohkam, M.; Masoumi, S.; Berenjian, A.; Ghasemi, Y. Prebiotics: Definition, Types, Sources, Mechanisms, and Clinical Applications. *Foods* **2019**, *8*, 92. [[CrossRef](#)]
197. Holscher, H.D. Dietary fiber and prebiotics and the gastrointestinal microbiota. *Gut Microbes* **2017**, *8*, 172–184. [[CrossRef](#)] [[PubMed](#)]
198. Liu, F.; Li, P.; Chen, M.; Luo, Y.; Prabhakar, M.; Zheng, H.; He, Y.; Qi, Q.; Long, H.; Zhang, Y.; et al. Fructooligosaccharide (FOS) and Galactooligosaccharide (GOS) Increase *Bifidobacterium* but Reduce Butyrate Producing Bacteria with Adverse Glycemic Metabolism in healthy young population. *Sci. Rep.* **2017**, *7*, 11789. [[CrossRef](#)] [[PubMed](#)]
199. Michel, M.R.; Rodríguez-Jasso, R.M.; Aguilar, C.N.; Gonzalez-Herrera, S.M.; Flores-Gallegos, A.C.; Rodríguez-Herrera, R. Fructosyltransferase Sources, Production, and Applications for Prebiotics Production. In *Probiotics and Prebiotics in Human Nutrition and Health*; Rao, V., Rao, L.G., Eds.; IntechOpen: London, UK, 2016; ISBN 978-953-51-2475-7.
200. Savaignac, H.M.; Corona, G.; Mills, H.; Chen, L.; Spencer, J.P.; Tzortzis, G.; Burnet, P.W. Prebiotic feeding elevates central brain derived neurotrophic factor, N-methyl-d-aspartate receptor subunits and d-serine. *Neurochem. Int.* **2013**, *63*, 756–764. [[CrossRef](#)] [[PubMed](#)]
201. Rossen, N.G.; Fuentes, S.; van der Spek, M.J.; Tijssen, J.G.; Hartman, J.H.A.; Duflou, A.; Löwenberg, M.; Van Den Brink, G.R.; Mathus-Vliegen, E.M.H.; de Vos, W.M.; et al. Findings From a Randomized Controlled Trial of Fecal Transplantation for Patients With Ulcerative Colitis. *Gastroenterology* **2015**, *149*, 110–118.e4. [[CrossRef](#)]

202. Chen, H.; Athar, R.; Zheng, G.; Williams, H.N. Prey bacteria shape the community structure of their predators. *ISME J.* **2011**, *5*, 1314–1322. [[CrossRef](#)]
203. Atterbury, R.J.; Hobbey, L.; Till, R.; Lambert, C.; Capeness, M.J.; Lerner, T.R.; Fenton, A.K.; Barrow, P.; Sockett, R.E. Effects of Orally Administered *Bdellovibrio bacteriovorus* on the Well-Being and *Salmonella* Colonization of Young Chicks. *Appl. Environ. Microbiol.* **2011**, *77*, 5794–5803. [[CrossRef](#)]
204. Hajjo, R.; Sabbah, D.A.; Al Bawab, A.Q. Unlocking the Potential of the Human Microbiome for Identifying Disease Diagnostic Biomarkers. *Diagnostics* **2022**, *12*, 1742. [[CrossRef](#)]
205. Zhou, Z.; Ge, S.; Li, Y.; Ma, W.; Liu, Y.; Hu, S.; Zhang, R.; Ma, Y.; Du, K.; Syed, A.; et al. Human Gut Microbiome-Based Knowledgebase as a Biomarker Screening Tool to Improve the Predicted Probability for Colorectal Cancer. *Front. Microbiol.* **2020**, *11*, 596027. [[CrossRef](#)]
206. Romo-Vaquero, M.; Fernández-Villalba, E.; Gil-Martinez, A.-L.; Cuenca-Bermejo, L.; Espín, J.C.; Herrero, M.T.; Selma, M.V. Urolithins: Potential biomarkers of gut dysbiosis and disease stage in Parkinson’s patients. *Food Funct.* **2022**, *13*, 6306–6316. [[CrossRef](#)]
207. Schwartz, A.; Spiegel, J.; Dillmann, U.; Grundmann, D.; Bürmann, J.; Faßbender, K.; Schäfer, K.-H.; Unger, M.M. Fecal markers of intestinal inflammation and intestinal permeability are elevated in Parkinson’s disease. *Park. Relat. Disord.* **2018**, *50*, 104–107. [[CrossRef](#)]
208. Rüb, A.M.; Tsakmaklis, A.; Gräfe, S.K.; Simon, M.-C.; Vehreschild, M.J.; Wuethrich, I. Biomarkers of human gut microbiota diversity and dysbiosis. *Biomark. Med.* **2021**, *15*, 139–150. [[CrossRef](#)]
209. Brydges, C.R.; Fiehn, O.; Mayberg, H.S.; Schreiber, H.; Dehkordi, S.M.; Bhattacharyya, S.; Cha, J.; Choi, K.S.; Craighead, W.E.; Krishnan, R.R.; et al. Indoxyl Sulfate, a Gut Microbiome-Derived Uremic Toxin, Is Associated with Psychic Anxiety and Its Functional Magnetic Resonance Imaging-Based Neurologic Signature. *Neuroscience* **2020**, *11*, 21011. [[CrossRef](#)]
210. Ling, Y.; Gong, T.; Zhang, J.; Gu, Q.; Gao, X.; Weng, X.; Liu, J.; Sun, J. Gut Microbiome Signatures Are Biomarkers for Cognitive Impairment in Patients With Ischemic Stroke. *Front. Aging Neurosci.* **2020**, *12*, 511562. [[CrossRef](#)]
211. De Ture, M.A.; Dickson, D.W. The neuropathological diagnosis of Alzheimer’s disease. *Mol. Neurodegener.* **2019**, *14*, 32. [[CrossRef](#)]
212. Serrano-Pozo, A.; Frosch, M.P.; Masliah, E.; Hyman, B.T. Neuropathological Alterations in Alzheimer Disease. *Cold Spring Harb. Perspect. Med.* **2011**, *1*, a006189. [[CrossRef](#)]
213. Murphy, M.P.; LeVine, H., 3rd. Alzheimer’s Disease and the Amyloid- β Peptide. *J. Alzheimer’s Dis.* **2010**, *19*, 311–323. [[CrossRef](#)]
214. Kim, H.S.; Kim, S.; Shin, S.J.; Park, Y.H.; Nam, Y.; Kim, C.W.; Lee, K.W.; Kim, S.-M.; Jung, I.D.; Yang, H.D.; et al. Gram-negative bacteria and their lipopolysaccharides in Alzheimer’s disease: Pathologic roles and therapeutic implications. *Transl. Neurodegener.* **2021**, *10*, 49. [[CrossRef](#)]
215. Zhan, X.; Stamova, B.; Sharp, F.R. Lipopolysaccharide Associates with Amyloid Plaques, Neurons and Oligodendrocytes in Alzheimer’s Disease Brain: A Review. *Front. Aging Neurosci.* **2018**, *10*, 42. [[CrossRef](#)]
216. Kowalski, K.; Mulak, A. Brain-Gut-Microbiota Axis in Alzheimer’s Disease. *J. Neurogastroenterol. Motil.* **2019**, *25*, 48–60. [[CrossRef](#)]
217. Chapman, M.; Robinson, L.S.; Pinkner, J.S.; Roth, R.; Heuser, J.; Hammar, M.; Normark, S.; Hultgren, S.J. Role of *Escherichia coli* Curli Operons in Directing Amyloid Fiber Formation. *Science* **2002**, *295*, 851–855. [[CrossRef](#)]
218. Friedland, R.P.; Chapman, M.R. The role of microbial amyloid in neurodegeneration. *PLoS Pathog.* **2017**, *13*, e1006654. [[CrossRef](#)] [[PubMed](#)]
219. Tiwari, D.; Singh, V.K.; Baral, B.; Pathak, D.K.; Jayabalan, J.; Kumar, R.; Tapryal, S.; Jha, H.C. Indication of Neurodegenerative Cascade Initiation by Amyloid-like Aggregate-Forming EBV Proteins and Peptide in Alzheimer’s Disease. *ACS Chem. Neurosci.* **2021**, *12*, 3957–3967. [[CrossRef](#)] [[PubMed](#)]
220. Alexander, G.E. Biology of Parkinson’s disease: Pathogenesis and pathophysiology of a multisystem neurodegenerative disorder. *Dialog- Clin. Neurosci.* **2004**, *6*, 259–280. [[CrossRef](#)]
221. Romano, S.; Savva, G.M.; Bedarf, J.R.; Charles, I.G.; Hildebrand, F.; Narbad, A. Meta-analysis of the Parkinson’s disease gut microbiome suggests alterations linked to intestinal inflammation. *npj Park. Dis.* **2021**, *7*, 27. [[CrossRef](#)]
222. Zapala, B.; Stefura, T.; Wójcik-Pędziwiatr, M.; Kabut, R.; Bałajewicz-Nowak, M.; Milewicz, T.; Dudek, A.; Stój, A.; Rudzińska-Bar, M. Differences in the Composition of Gut Microbiota between Patients with Parkinson’s Disease and Healthy Controls: A Cohort Study. *J. Clin. Med.* **2021**, *10*, 5698. [[CrossRef](#)]
223. Shen, T.; Yue, Y.; He, T.; Huang, C.; Qu, B.; Lv, W.; Lai, H.-Y. The Association Between the Gut Microbiota and Parkinson’s Disease, a Meta-Analysis. *Front. Aging Neurosci.* **2021**, *13*, 636545. [[CrossRef](#)]
224. Huang, Y.; Liao, J.; Liu, X.; Zhong, Y.; Cai, X.; Long, L. Review: The Role of Intestinal Dysbiosis in Parkinson’s Disease. *Front. Cell. Infect. Microbiol.* **2021**, *11*, 615075. [[CrossRef](#)]
225. Caspani, G.; Kennedy, S.; Foster, J.A.; Swann, J. Gut microbial metabolites in depression: Understanding the biochemical mechanisms. *Microb. Cell* **2019**, *6*, 454–481. [[CrossRef](#)]
226. Van Ijzendoorn, S.C.D.; Derkinderen, P. The Intestinal Barrier in Parkinson’s Disease: Current State of Knowledge. *J. Park. Dis.* **2019**, *9*, S323–S329. [[CrossRef](#)]
227. Mandel, I.; Paperna, T.; Glass-Marmor, L.; Volkowich, A.; Badarny, S.; Schwartz, I.; Vardi, P.; Koren, I.; Miller, A. Tight junction proteins expression and modulation in immune cells and multiple sclerosis. *J. Cell. Mol. Med.* **2011**, *16*, 765–775. [[CrossRef](#)]
228. Gorré, N.; Bauwens, E.; Haesebrouck, F.; Smet, A.; Vandenbroucke, R.E. *Helicobacter* and the Potential Role in Neurological Disorders: There Is More Than *Helicobacter pylori*. *Front. Immunol.* **2021**, *11*, 584165. [[CrossRef](#)]

229. Goldenberg, M.M. Medical management of Parkinson's disease. *Pharm. Ther.* **2008**, *33*, 590–606.
230. Nyholm, D.; Hellström, P.M. Effects of *Helicobacter pylori* on Levodopa Pharmacokinetics. *J. Park. Dis.* **2021**, *11*, 61–69. [[CrossRef](#)]
231. Lolekha, P.; Sriphanom, T.; Vilaichone, R.-K. *Helicobacter pylori* eradication improves motor fluctuations in advanced Parkinson's disease patients: A prospective cohort study (HP-PD trial). *PLoS ONE* **2021**, *16*, e0251042. [[CrossRef](#)]
232. Niehues, M.; Hensel, A. In vitro interaction of L-Dopa with bacterial adhesins of *Helicobacter pylori*: An explanation for clinical differences in bioavailability? *Planta Med.* **2009**, *75*, s-0029-1234811. [[CrossRef](#)]
233. Abdollahi, H.; Tadjrobehkar, O. The Role of Different Sugars, Amino Acids and Few Other Substances in Chemotaxis Directed Motility of *Helicobacter Pylori*. *Iran. J. Basic Med. Sci.* **2012**, *15*, 787–794. [[CrossRef](#)]
234. Rekdal, V.M.; Bess, E.N.; Bisanz, J.E.; Turnbaugh, P.J.; Balskus, E.P. Discovery and inhibition of an interspecies gut bacterial pathway for Levodopa metabolism. *Science* **2019**, *364*, eaau6323. [[CrossRef](#)]
235. Zuccato, C.; Valenza, M.; Cattaneo, E. Molecular Mechanisms and Potential Therapeutical Targets in Huntington's Disease. *Physiol. Rev.* **2010**, *90*, 905–981. [[CrossRef](#)]
236. Kong, G.; Cao, K.-A.L.; Judd, L.M.; Li, S.; Renoir, T.; Hannan, A.J. Microbiome profiling reveals gut dysbiosis in a transgenic mouse model of Huntington's disease. *Neurobiol. Dis.* **2020**, *135*, 104268. [[CrossRef](#)]
237. Du, G.; Dong, W.; Yang, Q.; Yu, X.; Ma, J.; Gu, W.; Huang, Y. Altered Gut Microbiota Related to Inflammatory Responses in Patients With Huntington's Disease. *Front. Immunol.* **2021**, *11*, 603594. [[CrossRef](#)]
238. Gerhardt, S.; Mohajeri, M.H. Changes of Colonic Bacterial Composition in Parkinson's Disease and Other Neurodegenerative Diseases. *Nutrients* **2018**, *10*, 708. [[CrossRef](#)] [[PubMed](#)]
239. Engels, C.; Ruscheweyh, H.-J.; Beerenwinkel, N.; Lacroix, C.; Schwab, C. The Common Gut Microbe *Eubacterium hallii* also Contributes to Intestinal Propionate Formation. *Front. Microbiol.* **2016**, *7*, 713. [[CrossRef](#)] [[PubMed](#)]
240. Pochard, P.; Gosset, P.; Grangette, C.; Andre, C.; Tonnel, A.-B.; Pestel, J.; Mercenier, A. Lactic acid bacteria inhibit TH2 cytokine production by mononuclear cells from allergic patients. *J. Allergy Clin. Immunol.* **2002**, *110*, 617–623. [[CrossRef](#)] [[PubMed](#)]
241. Bjorkqvist, M.; Wild, E.; Thiele, J.; Silvestroni, A.; Andre, R.; Lahiri, N.; Raibon, E.; Lee, R.V.; Benn, C.L.; Soulet, D.; et al. A novel pathogenic pathway of immune activation detectable before clinical onset in Huntington's disease. *J. Exp. Med.* **2008**, *205*, 1869–1877. [[CrossRef](#)] [[PubMed](#)]
242. Gadani, S.P.; Cronk, J.C.; Norris, G.T.; Kipnis, J. IL-4 in the Brain: A Cytokine To Remember. *J. Immunol.* **2012**, *189*, 4213–4219. [[CrossRef](#)]
243. Pirko, I.; Noseworthy, J.H. Demyelinating Disorders of the Central Nervous System. In *Textbook of Clinical Neurology*; Elsevier: Amsterdam, The Netherlands, 2007; pp. 1103–1133, ISBN 978-1-4160-3618-0.
244. Wootla, B.; Eriguchi, M.; Rodriguez, M. Is Multiple Sclerosis an Autoimmune Disease? *Autoimmune Dis.* **2012**, *2012*, 969657. [[CrossRef](#)]
245. Errede, M.; Girolamo, F.; Ferrara, G.; Strippoli, M.; Morando, S.; Boldrin, V.; Rizzi, M.; Uccelli, A.; Perris, R.; Bendotti, C.; et al. Blood-Brain Barrier Alterations in the Cerebral Cortex in Experimental Autoimmune Encephalomyelitis. *J. Neuropathol. Exp. Neurol.* **2012**, *71*, 840–854. [[CrossRef](#)]
246. Mu, Q.; Kirby, J.; Reilly, C.M.; Luo, X.M. Leaky Gut As a Danger Signal for Autoimmune Diseases. *Front. Immunol.* **2017**, *8*, 598. [[CrossRef](#)]
247. Cox, L.M.; Maghzi, A.H.; Liu, S.; Tankou, S.K.; Dhang, F.H.; Willocq, V.; Song, A.; Wasén, C.; Tauhid, S.; Chu, R.; et al. Gut Microbiome in Progressive Multiple Sclerosis. *Ann. Neurol.* **2021**, *89*, 1195–1211. [[CrossRef](#)]
248. Melbye, P.; Olsson, A.; Hansen, T.H.; Søndergaard, H.B.; Oturai, A.B. Short-chain fatty acids and gut microbiota in multiple sclerosis. *Acta Neurol. Scand.* **2018**, *139*, 208–219. [[CrossRef](#)]
249. Liu, J.; Jin, Y.; Ye, Y.; Tang, Y.; Dai, S.; Li, M.; Zhao, G.; Hong, G.; Lu, Z.-Q. The Neuroprotective Effect of Short Chain Fatty Acids Against Sepsis-Associated Encephalopathy in Mice. *Front. Immunol.* **2021**, *12*, 626894. [[CrossRef](#)]
250. Arpaia, N.; Campbell, C.; Fan, X.; Dikiy, S.; Van Der Veecken, J.; DeRoos, P.; Liu, H.; Cross, J.R.; Pfeffer, K.; Coffey, P.J.; et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* **2013**, *504*, 451–455. [[CrossRef](#)]
251. Rumah, K.R.; Linden, J.; Fischetti, V.A.; Vartanian, T. Isolation of *Clostridium perfringens* Type B in an Individual at First Clinical Presentation of Multiple Sclerosis Provides Clues for Environmental Triggers of the Disease. *PLoS ONE* **2013**, *8*, e76359. [[CrossRef](#)]
252. Tremlett, H.; Waubant, E. The multiple sclerosis microbiome? *Ann. Transl. Med.* **2017**, *5*, 53. [[CrossRef](#)]
253. Cekanaviciute, E.; Yoo, B.B.; Runia, T.F.; Debelius, J.W.; Singh, S.; Nelson, C.A.; Kanner, R.; Bencosme, Y.; Lee, Y.K.; Hauser, S.L.; et al. Gut bacteria from multiple sclerosis patients modulate human T cells and exacerbate symptoms in mouse models. *Proc. Natl. Acad. Sci. USA* **2017**, *114*, 10713–10718. [[CrossRef](#)]
254. Grant, S.M.; DeMorrow, S. Bile Acid Signaling in Neurodegenerative and Neurological Disorders. *Int. J. Mol. Sci.* **2020**, *21*, 75982. [[CrossRef](#)]
255. Marksteiner, J.; Blasko, I.; Kemmler, G.; Koal, T.; Humpel, C. Bile acid quantification of 20 plasma metabolites identifies lithocholic acid as a putative biomarker in Alzheimer's disease. *Metabolomics* **2017**, *14*, 1. [[CrossRef](#)]
256. Graham, S.F.; Rey, N.L.; Ugur, Z.; Yilmaz, A.; Sherman, E.; Maddens, M.; Bahado-Singh, R.O.; Becker, K.; Schulz, E.; Meyerdirk, L.K.; et al. Metabolomic Profiling of Bile Acids in an Experimental Model of Prodromal Parkinson's Disease. *Metabolites* **2018**, *8*, 71. [[CrossRef](#)]
257. Ackerman, H.D.; Gerhard, G.S. Bile Acids in Neurodegenerative Disorders. *Front. Aging Neurosci.* **2016**, *8*, 263. [[CrossRef](#)]

258. Nunes, A.F.; Amaral, J.D.; Lo, A.C.; Fonseca, M.B.; Viana, R.J.; Callaerts-Vegh, Z.; D'Hooge, R.; Rodrigues, C.M. TUDCA, a bile acid, attenuates amyloid precursor protein processing and amyloid- β deposition in APP/PS1 mice. *Mol. Neurobiol.* **2012**, *45*, 440–454. [[CrossRef](#)]
259. Castro-Caldas, M.; Carvalho, A.N.; Rodrigues, E.; Henderson, C.; Wolf, C.R.; Rodrigues, C.; Gama, M.J. Tauroursodeoxycholic Acid Prevents MPTP-Induced Dopaminergic Cell Death in a Mouse Model of Parkinson's Disease. *Mol. Neurobiol.* **2012**, *46*, 475–486. [[CrossRef](#)] [[PubMed](#)]
260. Nagpal, R.; Mainali, R.; Ahmadi, S.; Wang, S.; Singh, R.; Kavanagh, K.; Kitzman, D.W.; Kushugulova, A.; Marotta, F.; Yadav, H. Gut microbiome and aging: Physiological and mechanistic insights. *Nutr. Health Aging* **2018**, *4*, 267–285. [[CrossRef](#)] [[PubMed](#)]
261. Leite, G.; Pimentel, M.; Barlow, G.M.; Chang, C.; Hosseini, A.; Wang, J.; Parodi, G.; Sedighi, R.; Rezaie, A.; Mathur, R. Age and the aging process significantly alter the small bowel microbiome. *Cell Rep.* **2021**, *36*, 109765. [[CrossRef](#)] [[PubMed](#)]
262. Sharma, R. Emerging Interrelationship Between the Gut Microbiome and Cellular Senescence in the Context of Aging and Disease: Perspectives and Therapeutic Opportunities. *Probiotics Antimicrob. Proteins* **2022**, *14*, 648–663. [[CrossRef](#)] [[PubMed](#)]
263. Wilmanski, T.; Diener, C.; Rappaport, N.; Patwardhan, S.; Wiedrick, J.; Lapidus, J.; Earls, J.C.; Zimmer, A.; Glusman, G.; Robinson, M.; et al. Gut microbiome pattern reflects healthy ageing and predicts survival in humans. *Nat. Metab.* **2021**, *3*, 274–286. [[CrossRef](#)]
264. Lee, H.-Y.; Lee, S.-H.; Lee, J.-H.; Lee, W.-J.; Min, K.-J. The role of commensal microbes in the lifespan of *Drosophila melanogaster*. *Aging* **2019**, *11*, 4611–4640. [[CrossRef](#)]
265. Smith, P.; Willemsen, D.; Popkes, M.; Metge, F.; Gandiwa, E.; Reichard, M.; Valenzano, D.R. Regulation of life span by the gut microbiota in the short-lived *African turquoise killifish*. *eLife* **2017**, *6*, e27014. [[CrossRef](#)]
266. Hou, Y.; Dan, X.; Babbar, M.; Wei, Y.; Hasselbalch, S.G.; Croteau, D.L.; Bohr, V.A. Ageing as a risk factor for neurodegenerative disease. *Nat. Rev. Neurol.* **2019**, *15*, 565–581. [[CrossRef](#)]
267. Shah, Z.; Asejani, A. The influence of gut microbiota alteration on age-related neuroinflammation and cognitive decline. *Neural Regen. Res.* **2022**, *17*, 2407. [[CrossRef](#)]
268. Zhan, X.; Hakoupan, M.; Jin, L.-W.; Sharp, F.R. Lipopolysaccharide, Identified Using an Antibody and by PAS Staining, Is Associated With Corpora amylacea and White Matter Injury in Alzheimer's Disease and Aging Brain. *Front. Aging Neurosci.* **2021**, *13*, 705594. [[CrossRef](#)]
269. Wu, M.-L.; Yang, X.-Q.; Xue, L.; Duan, W.; Du, J.-R. Age-related cognitive decline is associated with microbiota-gut-brain axis disorders and neuroinflammation in mice. *Behav. Brain Res.* **2021**, *402*, 113125. [[CrossRef](#)]
270. Villarán, R.F.; Espinosa-Oliva, A.M.; Sarmiento, M.; De Pablos, R.M.; Argüelles, S.; Delgado-Cortés, M.J.; Sobrino, V.; Van Rooijen, N.; Venero, J.L.; Herrera, A.J.; et al. Ulcerative colitis exacerbates lipopolysaccharide-induced damage to the nigral dopaminergic system: Potential risk factor in Parkinson's disease. *J. Neurochem.* **2010**, *114*, 1687–1700. [[CrossRef](#)]
271. Devi, P.; Khan, A.; Chattopadhyay, P.; Mehta, P.; Sahni, S.; Sharma, S.; Pandey, R. Co-infections as Modulators of Disease Outcome: Minor Players or Major Players? *Front. Microbiol.* **2021**, *12*, 664386. [[CrossRef](#)]
272. Traylen, C.M.; Patel, H.R.; Fondaw, W.; Mahatme, S.; Williams, J.F.; Walker, L.R.; Dyson, O.F.; Arce, S.; Akula, S.M. Virus reactivation: A panoramic view in human infections. *Future Virol.* **2011**, *6*, 451–463. [[CrossRef](#)]
273. Li, N.; Ma, W.-T.; Pang, M.; Fan, Q.-L.; Hua, J.-L. The Commensal Microbiota and Viral Infection: A Comprehensive Review. *Front. Immunol.* **2019**, *10*, 1551. [[CrossRef](#)]
274. Pérez-Losada, M.; Arenas, M.; Galán, J.C.; Palero, F.; González-Candelas, F. Recombination in viruses: Mechanisms, methods of study, and evolutionary consequences. *Infect. Genet. Evol.* **2015**, *30*, 296–307. [[CrossRef](#)]
275. Diaz, C.D.; García-Orozco, A.; Riera-Leal, A.; Padilla-Arellano, J.R.; Fafutis-Morris, M. Microbiota and Its Role on Viral Evasion: Is It With Us or Against Us? *Front. Cell. Infect. Microbiol.* **2019**, *9*, 256. [[CrossRef](#)]
276. Follmer, C. Viral Infection-Induced Gut Dysbiosis, Neuroinflammation, and α -Synuclein Aggregation: Updates and Perspectives on COVID-19 and Neurodegenerative Disorders. *ACS Chem. Neurosci.* **2020**, *11*, 4012–4016. [[CrossRef](#)]
277. Dutta, S.K.; Verma, S.; Jain, V.; Surapaneni, B.; Vinayek, R.; Phillips, L.; Nair, P.P. Parkinson's Disease: The Emerging Role of Gut Dysbiosis, Antibiotics, Probiotics, and Fecal Microbiota Transplantation. *J. Neurogastroenterol. Motil.* **2019**, *25*, 363–376. [[CrossRef](#)]
278. Nandwana, V.; Debbarma, S. Fecal Microbiota Transplantation: A Microbiome Modulation Technique for Alzheimer's Disease. *Cureus* **2021**, *13*, e16503. [[CrossRef](#)]
279. Gupta, S.; Allen-Vercoe, E.; Petrof, E.O. Fecal microbiota transplantation: In perspective. *Ther. Adv. Gastroenterol.* **2015**, *9*, 229–239. [[CrossRef](#)] [[PubMed](#)]
280. Borody, T.; Brandt, L.J.; Paramsothy, S. Therapeutic faecal microbiota transplantation. *Curr. Opin. Gastroenterol.* **2014**, *30*, 97–105. [[CrossRef](#)] [[PubMed](#)]
281. Strandwitz, P. Neurotransmitter modulation by the gut microbiota. *Brain Res.* **2018**, *1693*, 128–133. [[CrossRef](#)] [[PubMed](#)]
282. Vendrik, K.E.W.; Ooijevaar, R.E.; De Jong, P.R.C.; Laman, J.D.; van Oosten, B.W.; Van Hilten, J.J.; Ducarmon, Q.R.; Keller, J.J.; Kuijper, E.J.; Contarino, M.F. Fecal Microbiota Transplantation in Neurological Disorders. *Front. Cell. Infect. Microbiol.* **2020**, *10*, 98. [[CrossRef](#)] [[PubMed](#)]
283. Van Kessel, S.P.; Frye, A.K.; El-Gendy, A.O.; Castejon, M.; Keshavarzian, A.; van Dijk, G.; El Aidy, S. Gut bacterial tyrosine decarboxylases restrict levels of levodopa in the treatment of Parkinson's disease. *Nat. Commun.* **2019**, *10*, 310–311. [[CrossRef](#)]
284. Holmes, A.; Finger, C.; Morales-Scheihing, D.; Lee, J.; McCullough, L.D. Gut dysbiosis and age-related neurological diseases; an innovative approach for therapeutic interventions. *Transl. Res.* **2020**, *226*, 39–56. [[CrossRef](#)]

285. Zhang, T.; Li, Q.; Cheng, L.; Buch, H.; Zhang, F. *Akkermansia muciniphila* is a promising probiotic. *Microb. Biotechnol.* **2019**, *12*, 1109–1125. [[CrossRef](#)]
286. Han, B.; Sivaramakrishnan, P.; Lin, C.-C.J.; Neve, I.A.; He, J.; Tay, L.W.R.; Sowa, J.N.; Sizovs, A.; Du, G.; Wang, J.; et al. Microbial Genetic Composition Tunes Host Longevity. *Cell* **2017**, *169*, 1249–1262.e13. [[CrossRef](#)]
287. Shin, J.; Noh, J.-R.; Choe, D.; Lee, N.; Song, Y.; Cho, S.; Kang, E.-J.; Go, M.-J.; Ha, S.K.; Chang, D.-H.; et al. Ageing and rejuvenation models reveal changes in key microbial communities associated with healthy ageing. *Microbiome* **2021**, *9*, 240. [[CrossRef](#)]
288. Wu, T.; Lyu, Y.; Li, X.; Wu, M.; Yu, K.; Li, S.; Ji, C.; Zhang, Q.; Zhang, Y.; Zhao, D.; et al. Impact of N-Acetylcysteine on the Gut Microbiota in the Piglets Infected With Porcine Epidemic Diarrhea Virus. *Front. Veter.-Sci.* **2021**, *7*, 582338. [[CrossRef](#)]
289. Tenório, M.C.d.S.; Graciliano, N.G.; Moura, F.; de Oliveira, A.C.M.; Goulart, M.O.F. N-Acetylcysteine (NAC): Impacts on Human Health. *Antioxidants* **2021**, *10*, 967. [[CrossRef](#)] [[PubMed](#)]
290. Zheng, J.; Yuan, X.; Zhang, C.; Jia, P.; Jiao, S.; Zhao, X.; Yin, H.; Du, Y.; Liu, H. N -Acetylcysteine Alleviates Gut Dysbiosis and Glucose Metabolic Disorder in High-Fat Diet-Fed Mice. *J. Diabetes* **2018**, *11*, 32–45. [[CrossRef](#)] [[PubMed](#)]
291. Lee, S.I.; Kang, K.S. N-acetylcysteine modulates lipopolysaccharide-induced intestinal dysfunction. *Sci. Rep.* **2019**, *9*, 1004. [[CrossRef](#)] [[PubMed](#)]