

Supplementary Materials

Isomadecassoside, a New Ursane-Type Triterpene Glycoside from Centella asiatica Leaves, Reduces Nitrite Levels in LPS-Stimulated Macrophages

Giuseppina Chianese¹, Francesca Masi¹, Donatella Ciccia¹, Daniele Ciceri², Sabrina Arpini², Mario Falzoni², Ester Pagano^{1,3}, Orazio Taglialatela-Scafati^{1,*}

Supporting data	Page
S1: ^1H NMR spectrum of isomadecassoside	2
S2: 2D NMR HSQC spectrum of isomadecassoside	2
S3: 2D NMR HMBC spectrum of isomadecassoside	3
S4: 2D NMR COSY spectrum of isomadecassoside	3
S5: 2D NMR NOESY spectrum of isomadecassoside	4

Figure S1: ^1H NMR spectrum of isomadecassoside (CD_3OD , 700MHz)




Figure S2: 2D NMR HSQC spectrum of isomadecassoside (CD_3OD)




Figure S3: 2D NMR HMBC spectrum of isomadecassoside (CD_3OD)




Figure S4: 2D NMR COSY spectrum of isomadecassoside (CD_3OD)




Figure S5: 2D NMR NOESY spectrum of isomadecassoside (CD_3OD)

