

Article

GC-MS Based Identification of the Volatile Components of Six *Astragalus* Species from Uzbekistan and Their Biological Activity

Haidy A. Gad ^{1,2}, Nilufar Z. Mamadalieva ^{3,4}, Stefan Böhmendorfer ⁵, Thomas Rosenau ⁵, Gokhan Zengin ⁶, Rano Z. Mamadalieva ⁷, Nawal M. Al Musayeib ^{8,*} and Mohamed L. Ashour ^{1,*}

- ¹ Department of Pharmacognosy, Faculty of Pharmacy, Ain Shams University, 11566 Cairo, Egypt; haidygad@pharma.asu.edu.eg
 - ² Department of Pharmacognosy, Faculty of Pharmacy, King Salman International University, South Sinai, 46612 Ras Sidr, Egypt
 - ³ Institute of the Chemistry of Plant Substances, Academy of Sciences of RUz, Mirzo Ulugbek str. 77, Tashkent 100170, Uzbekistan; nmamadalieva@yahoo.com
 - ⁴ Department of Bioorganic Chemistry, Leibniz Institute of Plant Biochemistry, Weinberg 3, D-06120 Halle (Saale), Germany
 - ⁵ Department of Chemistry, Institute of Chemistry of Renewable Resources, University of Natural Resources and Life Sciences, Vienna (BOKU University), Konrad-Lorenz-Straße 24, 3430 Tulln, Vienna, Austria; stefan.boehmdorfer@boku.ac.at (S.B.); thomas.rosenau@boku.ac.at (T.R.)
 - ⁶ Biochemistry and Physiology Research Laboratory, Department of Biology, Selcuk University, Science Faculty, 42130 Konya, Turkey; gokhanzengin@selcuk.edu.tr
 - ⁷ Kokand State Pedagogical Institute, Turon str. 23, 713000 Kokand, Uzbekistan; rmamadalieva@yahoo.com
 - ⁸ Department of Pharmacognosy, College of Pharmacy, King Saud University, 11495 Riyadh, Saudi Arabia
- * Correspondence: nalmusayeib@ksu.edu.sa (N.M.A.); ashour@pharma.asu.edu.eg (M.L.A.); Tel.: +20-10-68-222-354 (M.L.A.); Fax: +20-22-405-1107 (M.L.A.)

2. Results and Discussion

2.2. Chemometric analysis based on GC-MS




Figure 1S. Enlarged correlation-loading plot of GC-MS analysis of essential oils of different *Astragalus* species based on the identification of compounds shown in Table (1).

3. Materials and methods

3.5. Antioxidant and enzyme inhibitory assays

Determination of Antioxidant and Enzyme Inhibitory Effects

Antioxidant (DPPH and ABTS radical scavenging, reducing power (CUPRAC and FRAP), phosphomolybdenum and metal chelating (ferrozine method)) and enzyme inhibitory activities (cholinesterase (Eldmann's method), tyrosinase (dopachrome method), α -amylase (iodine/potassium iodide method), and α -glucosidase (chromogenic PNPG method)) were determined using the methods previously described by Uysal et al. [1] and Grochowski et al. [2]

For the DPPH (1,1-diphenyl-2-picrylhydrazyl) radical scavenging assay: Sample solution was added to 4 mL of a 0.004% methanol solution of DPPH. The sample absorbance was read at 517 nm after a 30 min incubation at room temperature in the dark. DPPH radical scavenging activity was expressed as milligrams of trolox equivalents (mg TE/g oil).

For ABTS (2,2'-azino-bis(3-ethylbenzothiazoline) 6-sulfonic acid) radical scavenging assay: Briefly, ABTS⁺ was produced directly by reacting 7 mM ABTS solution with 2.45 mM potassium persulfate and allowing the mixture to stand for 12–16 h in the dark at room temperature. Prior to beginning the assay, ABTS solution was diluted with methanol to an absorbance of 0.700 ± 0.02 at 734 nm. Sample solution was added to ABTS solution (2 mL) and mixed. The sample absorbance was read at 734 nm after a 30 min incubation at room temperature. The ABTS radical scavenging activity was expressed as milligrams of trolox equivalents (mg TE/g oil).

For CUPRAC (cupric ion reducing activity) activity assay: Sample solution was added to premixed reaction mixture containing CuCl₂ (1 mL, 10 mM), neocuproine (1 mL, 7.5 mM) and NH₄Ac buffer (1 mL, 1 M, pH 7.0). Similarly, a blank was prepared by adding sample solution (0.5 mL) to premixed reaction mixture (3 mL) without CuCl₂. Then, the sample and blank absorbances were read at 450 nm after a 30 min incubation at room temperature. The absorbance of the blank was subtracted from that of the sample. CUPRAC activity was expressed as milligrams of trolox equivalents (mg TE/g oil).

For FRAP (ferric reducing antioxidant power) activity assay: Sample solution was added to premixed FRAP reagent (2 mL) containing acetate buffer (0.3 M, pH 3.6), 2,4,6-tris(2-pyridyl)-S-triazine (TPTZ) (10 mM) in 40 mM HCl and ferric chloride (20 mM) in a ratio of 10:1:1 (v/v/v). Then, the sample absorbance was read at 593 nm after a 30 min incubation at room temperature. FRAP activity was expressed as milligrams of trolox equivalents (mg TE/g oil).

For phosphomolybdenum method: Sample solution was combined with 3 mL of reagent solution (0.6 M sulfuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The sample absorbance was read at 695 nm after a 90 min incubation at 95 °C. The total antioxidant capacity was expressed as millimoles of trolox equivalents (mmol TE/g oil).

For metal chelating activity assay: Briefly, sample solution was added to FeCl₂ solution (0.05 mL, 2 mM). The reaction was initiated by the addition of 5 mM ferrozine (0.2 mL). Similarly, a blank was prepared by adding sample solution (2 mL) to FeCl₂ solution (0.05 mL, 2 mM) and water (0.2 mL) without ferrozine. Then, the sample and blank absorbances were read at 562 nm after 10 min incubation at room temperature. The absorbance of the blank was subtracted from that of the sample. The metal chelating activity was expressed as milligrams of EDTA (disodium edetate) equivalents (mg EDTAE/g oil).

For Cholinesterase (ChE) inhibitory activity assay: Sample solution (was mixed with DTNB (5,5-dithio-bis(2-nitrobenzoic) acid, Sigma, St. Louis, MO, USA) (125 μ L) and AChE (acetylcholinesterase (Electric eel acetylcholinesterase, Type-VI-S, EC 3.1.1.7, Sigma)), or BChE (butyrylcholinesterase (horse

serum butyrylcholinesterase, EC 3.1.1.8, Sigma)) solution (25 μ L) in Tris-HCl buffer (pH 8.0) in a 96-well microplate and incubated for 15 min at 25 °C. The reaction was then initiated with the addition of acetylthiocholine iodide (ATCI, Sigma) or butyrylthiocholine chloride (BTCl, Sigma) (25 μ L). Similarly, a blank was prepared by adding sample solution to all reaction reagents without enzyme (AChE or BChE) solution. The sample and blank absorbances were read at 405 nm after 10 min incubation at 25 °C. The absorbance of the blank was subtracted from that of the sample and the cholinesterase inhibitory activity was expressed as galanthamine equivalents (mg GALAE/g oil).

For Tyrosinase inhibitory activity assay: Sample solution was mixed with tyrosinase solution (40 μ L, Sigma) and phosphate buffer (100 μ L, pH 6.8) in a 96-well microplate and incubated for 15 min at 25 °C. The reaction was then initiated with the addition of L-DOPA (40 μ L, Sigma). Similarly, a blank was prepared by adding sample solution to all reaction reagents without enzyme (tyrosinase) solution. The sample and blank absorbances were read at 492 nm after a 10 min incubation at 25 °C. The absorbance of the blank was subtracted from that of the sample and the tyrosinase inhibitory activity was expressed as kojic acid equivalents (mg KAE/g oil).

For α -amylase inhibitory activity assay: Sample solution was mixed with α -amylase solution (ex-porcine pancreas, EC 3.2.1.1, Sigma) (50 μ L) in phosphate buffer (pH 6.9 with 6 mM sodium chloride) in a 96-well microplate and incubated for 10 min at 37 °C. After pre-incubation, the reaction was initiated with the addition of starch solution (50 μ L, 0.05%). Similarly, a blank was prepared by adding sample solution to all reaction reagents without enzyme (α -amylase) solution. The reaction mixture was incubated 10 min at 37 °C. The reaction was then stopped with the addition of HCl (25 μ L, 1 M). This was followed by addition of the iodine-potassium iodide solution (100 μ L). The sample and blank absorbances were read at 630 nm. The absorbance of the blank was subtracted from that of the sample and the α -amylase inhibitory activity was expressed as acarbose equivalents (mmol ACE/g oil).

1. Uysal, S.; Zengin, G.; Locatelli, M.; Bahadori, M. B.; Mocan, A.; Bellagamba, G.; De Luca, E.; Mollica, A.; Aktumsek, A., Cytotoxic and enzyme inhibitory potential of two *Potentilla* species (*P. speciosa* L. and *P. reptans* Willd.) and their chemical composition. *Frontiers in pharmacology* **2017**, *8*, 290.
2. Grochowski, D. M.; Uysal, S.; Aktumsek, A.; Granica, S.; Zengin, G.; Ceylan, R.; Locatelli, M.; Tomczyk, M., In vitro enzyme inhibitory properties, antioxidant activities, and phytochemical profile of *Potentilla thuringiaca*. *Phytochemistry Letters* **2017**, *20*, 365-372.