



Supplementary Figure S1. Illustrative picture showing the experimental design used in this study with the representative phenotype of sweet pepper (*Capsicum annuum* L.) fruits at different stages and treatments: immature green, breaking point 1 (BP1), breaking point 2 without NO treatment (BP2 – NO), breaking point 2 with NO treatment (BP2 + NO), and ripe red. Fruits were subjected to a NO-enriched atmosphere (5 ppm) in a hermetic box for 1 h and were then stored at room temperature (RT) for 3 days. Reproduced with permission from González-Gordo et al. (2022b) [46].

Supplementary Table S1. List of the 53 different isozymes of G6PDH belonging to 15 different plant species used in this work for phylogenetic analysis.

Plant species	G6PDH isozyme	Protein ID	Reference
Pepper (<i>Capsicum annuum</i>)	CaG6PDH1 CaG6PDH2 CaG6PDH3 CaG6PDH4	XP_016561682.1 XP_016570698.1 XP_016580465.1 XP_016573493.1	The present study
<i>Arabidopsis thaliana</i>	AtG6PDH1 AtG6PDH2 AtG6PDH3 AtG6PDH4 AtG6PDH5 AtG6PDH6	Q43727.2 Q9FY99.2 Q8L743.2 Q93ZW0.1 Q9LK23.1 Q9FJI5.1	Wendt et al., 1999; Wakao and Benning, 2004)
Alfalfa (<i>Medicago sativa</i>)	MsG6PDH1	AAB41552.1	(Wendt et al., 1999)
Tobacco (<i>Nicotiana tabacum</i>)	NtG6PDH1 NtG6PDH2 NtG6PDH3	CAA04994.1 AAF87216.1 CAA04992.1	(Wendt et al., 1999)
Rice (<i>Oryza sativa</i>)	OsG6PDH1 OsG6PDH2 OsG6PDH3	AAS07054.1 BAC84352.1 CAC09489.2	(Wendt et al., 1999)
Parsley (<i>Petroselinum crispum</i>)	PcG6PDH1 PcG6PDH2 PcG6PDH3	AAB69318.1 AAB69319.1 AAB69317.1	(Wendt et al., 1999)
Potato (<i>Solanum tuberosum</i>)	StG6PDH1 StG6PDH2 StG6PDH3	CAA52442.1 CAA58775.1 CAB52708.1	(Wendt et al., 1999)
Spinach (<i>Spinacia oleracea</i>)	SoG6PDH1	CAA03939.1	(Wendt et al., 1999)
Wheat (<i>Triticum aestivum</i>)	TaG6PDH1	BAA97663.1	(Wendt et al., 1999)
Rubber tree (<i>Hevea brasiliensis</i>)	HbG6PDH1 HbG6PDH2 HbG6PDH3 HbG6PDH4	AIE47266.1 AIE47267.1 AIE47268.1 AIE47269.1	(Long et al., 2016)
Black cottonwood (<i>Populus trichocarpa</i>)	PtG6PDH1 PtG6PDH2 PtG6PDH3 PtG6PDH4	EEE79649.2 ERP53365.1 EEE94856.1 EEF03929.1	(Long et al., 2016)
Castor oil (<i>Ricinus communis</i>)	RcG6PDH1 RcG6PDH2 RcG6PDH3	EEF47431.1 EEF32168.1 EEF50009.1	(Long et al., 2016)
Maize (<i>Zea mays</i>)	ZmG6PDH1 ZmG6PDH2 ZmG6PDH3 ZmG6PDH4	AFW57831.1 DAA45780.1 NP_001348206.1 ACG39996.1	(Long et al., 2016)
Tomato (<i>Solanum lycopersicum</i>)	SlG6PDH1 SlG6PDH2 SlG6PDH3 SlG6PDH4	XP_004230338.1 XP_004231802.1 XP_004239245.1 XP_004243567.1	Tomato proteome

Soybean (<i>Glycine max</i>)	GmG6PDH1	XP_003520786.1	(Zhao et al., 2020)
	GmG6PDH2	NP_001241264.1	
	GmG6PDH3	XP_003533032.1	
	GmG6PDH4	XP_003547679.1	
	GmG6PDH5	XP_003518668.1	
	GmG6PDH6	XP_003553875.1	
	GmG6PDH7	XP_003552660.1	
	GmG6PDH8	XP_014633122.1	
	GmG6PDH9	XP_003554604.1	

References

- Long, X., He, B., Fang, Y., & Tang, C. (2016). Identification and characterization of the glucose-6-phosphate dehydrogenase gene family in the para rubber tree, *Hevea brasiliensis*. *Frontiers in Plant Science*, 7, 215. <https://doi.org/10.3389/fpls.2016.00215>
- Wakao, S., & Benning, C. (2004). Genome-wide analysis of glucose-6-phosphate dehydrogenases in *Arabidopsis*. *The Plant Journal*, 41, 243–256. <https://doi.org/10.1111/j.1365-313x.2004.02293.x>
- Wendt, U. K., Hauschild, R., Lange, C., Pietersma, M., Wenderoth, I., & von Schaewen, A. (1999). *Plant Molecular Biology*, 40, 487–494. <https://doi.org/10.1023/a:1006257230779>
- Zhao, Y., Cui, Y., Huang, S., Yu, J., Wang, X., Xin, D., Li, X., Liu, Y., Dai, Y., Qi, Z., & Chen, Q. (2020). Genome-wide analysis of the glucose-6-phosphate dehydrogenase family in soybean and functional identification of GMG6PDH2 involvement in salt stress. *Frontiers in Plant Science*, 11, 214. <https://doi.org/10.3389/fpls.2020.00214>