

Survey Instrument

Public Perceptions of Community Pharmacy-Based Naloxone Services: A National Cross-Sectional Survey

I. INFORMATION ABOUT YOU

1. Enter your age in years: _____
2. Please indicate your sex
 Male Female Non-binary Other Prefer not to answer
3. With which race do you identify most closely?
 Black/African American Asian or Pacific Islander
 Native American or Alaska Native White/Caucasian Other
 Prefer not to answer
4. Are you of Hispanic, Latino(a), or Spanish origin?
 Yes No Prefer not to answer
5. What is the highest level of education you have completed?
 Some high school, no diploma
 High school graduate, diploma or the equivalent (for example: GED)
 Some college credit, no degree Trade/technical/vocational training Associate degree
 Bachelor's degree Master's degree Professional degree
 Doctorate degree
6. Please indicate your annual household income:
 Less than \$20,000 \$20,000-\$49,999 \$50,000-\$99,999 \$100,000-\$149,999
 \$150,000 or more
7. Are you employed in a healthcare profession or non-healthcare profession?
 Healthcare profession Non-healthcare profession Not employed
8. Please indicate the State in which you reside:
 Select from dropdown list of states, territories, and DC Prefer not to answer
9. Please indicate the Zip Code in which you reside: _____
10. Have you or someone you have known been personally or professionally affected by opioid use disorder (OUD) or substance misuse?
 Yes No
11. Are you currently using prescription opioid medications or have you been prescribed opioid medications in the past for the treatment of pain or for post-surgery management? Prescription opioid medications may include morphine, hydrocodone, Vicodin®, Norco®, or others.
 Yes No

II. GENERAL KNOWLEDGE ABOUT OPIOID OVERDOSE AND NALOXONE

1. Before taking this survey, were you aware of naloxone, also commonly referred to as Narcan®, and what it is used for?
 Yes No
2. What is naloxone used for?
 To reverse the effects of an opioid overdose (e.g. heroin, Oxycodone, Vicodin, methadone)
 To reverse the effects of an amphetamine overdose
 To reverse the effects of a cocaine overdose To reverse the effects of any overdose

3. [Display after Q2 has been answered] Naloxone, or Narcan[®] is a medication used for the treatment of opioid overdose and can help to reverse an opioid overdose. Were you aware that in all 50 states you can obtain this medication without a prescription by speaking with your pharmacist?
Yes No
4. Which of the following factors increase the risk of an opioid overdose? Please check all that apply.
- | | |
|--|---|
| <input type="checkbox"/> Taking larger than usual doses of opioids | <input type="checkbox"/> Increase in opioid purity |
| <input type="checkbox"/> Using an opioid again after not having used for a while | <input type="checkbox"/> A long history of opioid use |
| <input type="checkbox"/> Using an opioid when no one else is present around | |
| <input type="checkbox"/> Using an opioid again after a detoxification treatment | |
5. Which of the following are indicators of an opioid overdose? Please check all that apply.
- | | | |
|---|--|--|
| <input type="checkbox"/> Having bloodshot eyes | <input type="checkbox"/> Slow or shallow breathing | <input type="checkbox"/> Unresponsive |
| <input type="checkbox"/> Lips, hands or feet turning blue | <input type="checkbox"/> Loss of consciousness | <input type="checkbox"/> Deep snoring |
| <input type="checkbox"/> Very small pupils | <input type="checkbox"/> Agitated behavior | <input type="checkbox"/> Rapid heartbeat |
6. Which of the following should be done when managing an opioid overdose? Please check all that apply.
- | | |
|---|---|
| <input type="checkbox"/> Stay with the person until an ambulance arrives | <input type="checkbox"/> Call an ambulance |
| <input type="checkbox"/> Inject the person with salt solution or milk | <input type="checkbox"/> Give mouth to mouth resuscitation |
| <input type="checkbox"/> Give stimulants (e.g. cocaine or black coffee) | <input type="checkbox"/> Check for breathing |
| <input type="checkbox"/> Give naloxone (opioid overdose antidote) | <input type="checkbox"/> Put the person in a bath of cold water |
| <input type="checkbox"/> Check for blocked airways (nose and mouth) | <input type="checkbox"/> Put the person in bed to sleep it off |
| <input type="checkbox"/> Place the person in the recovery position (on their side with mouth clear) | |
7. How can Naloxone be administered? Please check all that apply.
- | | | |
|--|--|--|
| <input type="checkbox"/> Into the nose (intranasal) | <input type="checkbox"/> Into a muscle (intramuscular) | <input type="checkbox"/> Into a vein (intravenous) |
| <input type="checkbox"/> Under the skin (subcutaneous) | <input type="checkbox"/> Swallowing a liquid | <input type="checkbox"/> Swallowing a tablet |
| <input type="checkbox"/> Don't know | | |
8. How long does naloxone take to start having an effect?
- | | | | |
|--------------------------------------|---------------------------------------|--|--|
| <input type="checkbox"/> 2-5 minutes | <input type="checkbox"/> 6-10 minutes | <input type="checkbox"/> 11-20 minutes | <input type="checkbox"/> 21-40 minutes |
| <input type="checkbox"/> Don't know | | | |
9. How long do the effects of naloxone last for?
- | | | | |
|---|---------------------------------------|---------------------------------------|--|
| <input type="checkbox"/> Less than 20 minutes | <input type="checkbox"/> About 1 hour | <input type="checkbox"/> 1 to 6 hours | <input type="checkbox"/> 6 to 12 hours |
| <input type="checkbox"/> Don't know | | | |
10. Which statements are CORRECT about naloxone and opioid overdose? Please check all that apply.
- If the first dose of naloxone has no effect a second dose can be given
 - There is no need to call for an ambulance if I know how to manage an overdose
 - Someone can overdose again even after having received naloxone
 - Naloxone can provoke withdrawal symptoms
 - After recovering from an opioid overdose, the person must not take any heroin, but it is OK for them to drink alcohol or take sleeping pills

III. GENERAL OPINIONS ABOUT NALOXONE

1. On a scale of 1-5, with 1 being strongly disagree and 5 being strongly agree, please indicate your level of agreement or disagreement with the following statements.

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
Naloxone provides opioid users with a second chance at life	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Naloxone may cause MORE opioid overdoses	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Having naloxone available ENABLES people who misuse opioids	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Naloxone should be made available to anyone concerned about opioid overdose	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If my pharmacist speaks to me about naloxone, I would like to be moved to a private area in the pharmacy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If my doctor prescribed me opioids, I would NOT need naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

IV. HOW TO RECEIVE NALOXONE FROM THE PHARMACY

1. On a scale of 1-5, with 1 being strongly disagree and 5 being strongly agree, please indicate your level of agreement or disagreement with the following statements.

What things make it DIFFICULT to receive naloxone from the pharmacy?

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
Cost of naloxone is too high	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is uncomfortable asking the pharmacist for naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It is uncomfortable asking the pharmacist for naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Not wanting my family/friends to know that I was getting naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Not wanting people I know from my community/neighborhood to see me getting naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fear of the pharmacist having a negative view of me if I ask for naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
It will take too long to receive naloxone from the pharmacy, e.g. long wait times	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

2. What do you think would be the BEST way for a pharmacist to ask you if you wanted naloxone when you pick up a prescription for opioid pain medications?
- Universal offer – pharmacist tells all customers at the pharmacy that naloxone is available
 - Targeted offer – pharmacist tells only those customers at higher risk of overdose about naloxone
 - General advertisement – pharmacist posts a flyer so that all customers in the pharmacy are aware naloxone is available

V. OPINIONS ABOUT PHARMACISTS AND NALOXONE

1. On a scale of 1-5, with 1 being strongly disagree and 5 being strongly agree, please indicate your level of agreement or disagreement with the following statements.

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
I am confident that pharmacists have sufficient KNOWLEDGE to dispense naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I am confident that pharmacists have sufficient TRAINING to dispense naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I support community pharmacists dispensing naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I prefer receiving naloxone from my pharmacist rather than my doctor	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would only get naloxone from my pharmacy if my doctor recommended it first	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If I received naloxone from my pharmacy, I would NOT want them to tell my doctor	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I prefer to receive naloxone from a more clinical setting, like a doctor's office	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
The pharmacy is an appropriate place to provide naloxone	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I worry how the pharmacy will handle my naloxone records	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
There is not enough privacy in the pharmacy	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would prefer to receive naloxone from an anonymous source, like the public health department	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

VI. YOUR ABILITY TO MANAGE AN OPIOID OVERDOSE

On a scale of 1-5, with 1 being strongly disagree and 5 being strongly agree, please indicate your level of agreement or disagreement with the following statements.

1. Competencies to Manage an Opioid Overdose

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
I already have enough information about how to manage an overdose	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I am already able to inject naloxone into someone who has overdosed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be able to check that someone who has overdosed was breathing properly	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I am going to need more training before I would feel confident to help someone who has overdosed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be able to perform mouth to mouth resuscitation to someone who has overdosed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be able to perform chest compressions to someone who has overdosed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If someone overdoses, I would know what to do to help them	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be able to place someone who has overdosed in the recovery position	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I know very little about how to help someone who has overdosed	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be able to deal effectively with an overdose	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

2. Concerns About Managing an Opioid Overdose

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
I would be afraid of giving naloxone in case the person becomes aggressive afterwards	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would be afraid of doing something wrong in an overdose situation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

I would be reluctant to use naloxone for fear of causing withdrawal symptoms	<input type="radio"/>				
I would be concerned about calling emergency services in case the police come around	<input type="radio"/>				
If I tried to help someone who has overdosed, I might accidentally hurt them	<input type="radio"/>				
I would feel safer if I knew that naloxone was around	<input type="radio"/>				
I would be afraid of suffering a needle stick injury if I had to give someone a naloxone injection	<input type="radio"/>				
Needles frighten me and I wouldn't be able to give someone an injection of naloxone	<input type="radio"/>				

3. Readiness to Intervene in an Opioid Overdose

Questions	(1) Strongly Disagree	Disagree	Neutral	Agree	(5) Strongly Agree
Everyone at risk of witnessing an overdose should be given a naloxone supply	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I couldn't just watch someone overdose, I would have to do something to help	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If someone overdoses, I would call an ambulance but I wouldn't be willing to do anything else	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Family and friends of drug users should be prepared to deal with an overdose	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If I saw an overdose, I would panic and not be able to help	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If I witnessed an overdose, I would call an ambulance straight away	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I would stay with the overdose victim until help arrives	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
If I saw an overdose, I would feel nervous, but I would still take the necessary actions	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
I will do whatever is necessary to save someone's life in an overdose situation	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

If someone overdoses, I want to be able to help them	○	○	○	○	○
--	---	---	---	---	---

VII. CONCLUSION

Thank you so much for your time! We appreciate your thoughts and feedback regarding naloxone. If you would like more information regarding Narcan, feel free to visit <https://www.narcan.com/>