

Article

Matrix Approach for Analyzing n -Site Generalized ASIP Systems: PGF and Site Occupancy Probabilities

Uri Yechiali and Yaron Yeger * 

Department of Statistics and Operations Research, School of Mathematical Sciences, Tel Aviv University, Tel Aviv 6329302, Israel

* Correspondence: yaronyeg@mail.tau.ac.il

Abstract: The Asymmetric Simple Inclusion Process (ASIP) is an n -site tandem stochastic network with a Poisson arrival influx into the first site. Each site has an unlimited buffer with a gate in front of it. Each gate opens, independently of all other gates, following a site-dependent Exponential inter-opening time. When a site's gate opens, all particles occupying the site move simultaneously to the next site. In this paper, a Generalized ASIP network is analyzed where the influx is to all sites, while gate openings are determined by a general renewal process. A compact matrix approach—instead of the conventional (and tedious) successive substitution method—is constructed for the derivation of the multidimensional probability-generating function (PGF) of the site occupancies. It is shown that the set of $\binom{2^n}{n}$ linear equations required to obtain the PGF of an n -site network can be first cut by half into a set of $\binom{2^{n-1}}{n}$ equations, and then further reduced to a set of $2^n - (n + 1)$ equations. The latter set can be additionally split into several smaller triangular subsets. It is also shown how the PGF of an $(n + 1)$ -site network can be derived from the corresponding PGF of an n -site system. Explicit results for networks with $n = 3$ and $n = 4$ sites are obtained. The matrix approach is utilized to explicitly calculate the probability that site k ($k = 1, 2, \dots, n$) is occupied. We show that, in the case where arrivals occur to the first site only, these probabilities are functions of both the site's index and the arrival flux and not solely of the site's index. Consequently, refined formulas for the latter probabilities and for the mean conditional site occupancies are derived. We further show that in the case where the arrival process to the first site is Poisson with rate λ , the following interesting property holds: $P(\text{site } k \text{ is occupied} \mid \lambda = 1) = P(\text{site } k + 1 \text{ is occupied} \mid \lambda \rightarrow \infty)$. The case where the inter-gate opening intervals are Gamma distributed is investigated and explicit formulas are obtained. Mean site occupancy and mean total load of the first k sites are calculated. Numerical results are presented.



Citation: Yechiali, U.; Yeger, Y. Matrix Approach for Analyzing n -Site Generalized ASIP Systems: PGF and Site Occupancy Probabilities. *Mathematics* **2022**, *10*, 4624. <https://doi.org/10.3390/math10234624>

Academic Editor: Tuan Phung-Duc

Received: 24 October 2022

Accepted: 3 December 2022

Published: 6 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Keywords: Asymmetric Simple Inclusion Process (ASIP); Generalized ASIP (G-ASIP); multidimensional PGF; matrix approach; site occupancies

MSC: 60K25; 68M20; 90B22

1. Introduction and Outline

A tandem stochastic system (TSS) is an array of several sites in series, where random events cause particles (customers, messages, products, calls, jobs, molecules, etc.) to propagate unidirectionally along the one-dimensional lattice of n sites (queues, servers, stations, etc.). Particles are fed, randomly in time, into the leftmost site and propagate unidirectionally (to the right) through the system. At the rightmost site, particles exit the system randomly in time. The random inflow into the leftmost site, the random instants of movement from site to site, and the random outflow from the rightmost site are all governed by random processes. Variations of this model have been explored in various papers [1–11].

A notable example of TSS is the Tandem Jackson Network (TJN), which has been investigated thoroughly in the literature [6–11]. In this model, the buffer size of each

site is unlimited, only single particles move forward between sites and the transitions instants follow Markovian rules. The TJN is famous for its product-form solution of the multidimensional distribution function of the sites' queue sizes (occupancies).

Another well-known TSS is the Asymmetric Exclusion Process (ASEP) [3,4], where each site can hold at most a single particle, a constraint that causes blockings on particles' forward movements. The ASEP was extensively studied in [12–27]. As described in [28], “The ASEP serves as a model for a unidirectionally driven lattice gas of particles subject to exclusion interactions. The exclusion principle causes jamming throughout the system and renders the ASEP dynamics highly nontrivial. Despite its simple description and its one dimensionality, the ASEP displays a complex and intricate behavior [12]”.

An important process in the family of tandem stochastic processes is the recently introduced [28] Asymmetric Inclusion Process (ASIP) [28–32]. Similarly to the ASEP and TJN, particles propagate unidirectionally through the system. The ASIP's inclusion principle allows each site to be occupied by an arbitrary number of particles at the same time. All particles that simultaneously occupy a site form a cluster of particles that move together to the next site, where a new and larger cluster will form together with the particles already residing in the new site (if any), or out of the system, in the case of the rightmost site. Each site has an unbounded buffer capacity and an unlimited size batch service. The ASIP is a showcase of complexity [29]. To better illustrate the ASIP, one can imagine that each site has a gate in front of it and when the gate opens, all particles currently in that site move forward to the next site. The ASIP model may be viewed as a tandem array of growth–collapse processes. Furthermore, it can be applied to the analysis of road traffic along a sequence of traffic lights, where, when a traffic light turns green, all vehicles in front of this light move forward to the next light. Another application is the analysis of marine traffic along a canal with several locks (e.g., Panama Canal). An illustration of the ASIP, with arrivals to the first site only, appears in Figure 1.



Figure 1. An illustration of the Asymmetric Inclusion Process (ASIP).

In a TSS, each site is characterized by its occupancy capacity, C_{site} , and by its gate (service) capacity, C_{gate} . Consequently, the above 3 TSS variations (TJN, ASEP and ASIP) are distinguished by their values of C_{site} and C_{gate} . Table 1 below characterizes the differences and similarities between the three models.

Table 1. Characterization of Tandem Stochastic Systems (TSS).

	Site (Occupancy) Capacity $C_{site}=1$	Site (Occupancy) Capacity C_{site} Is Unlimited
Gate (service) capacity $C_{gate} = 1$	ASEP	TJN
Gate (service) capacity C_{gate} is unlimited	ASEP	ASIP

An important feature of the classical ASIP model [28] is that its multidimensional Probability-Generating Function (PGF) of the site occupancies does not admit a product-form solution (as does the TJN), which makes its analysis highly complicated and intricate. Consequently, explicit PGF expressions were obtained only for ASIPs with $n = 2$ and $n = 3$ sites.

In this work we study a generalized ASIP (abbreviated G-ASIP) model where particles arrive to all sites, and inter-gate openings follow a renewal process, as is described below.

A Generalized ASIP Model (G-ASIP)

Consider a system of n sites (queues) Q_1, \dots, Q_n in series. Each queue has one gate in front of it, which may be viewed as a server. In the classical ASIP model (Reuveni, Eliazar and Yechiali [28]) particles arrive only to the first queue according to a Poisson process with rate λ , while gate j opens independently of the other gates every Exponential time with parameter μ_j . We study a generalized ASIP where the time intervals between consecutive instants of gate opening are i.i.d random variables, all distributed like a random variable O . At an instant of gate opening, only one gate opens, and it is gate j with probability $p_j > 0$, where $\sum_{j=1}^n p_j = 1$. During intervals between successive gate openings, particles arrive according to a random process. Gates are closed almost all the time. When gate j (the gate in front Q_j) opens, all particles present in Q_j are instantaneously transferred to Q_{j+1} , $j = 1, \dots, n-1$, where they form a larger cluster with the particles already residing there. When the last gate (gate n) opens, all particles present in Q_n simultaneously leave the system. This extension of the model allows for the analysis of systems where the times between gate openings are not necessarily Exponentially distributed (as in the classical ASIP) but rather generally. For example, consider a road traffic between traffic lights, or a marine traffic through a canal with several locks, where the evolution of each system is governed by Deterministically distributed time intervals.

The classical ASIP model was further investigated and analyzed by the above authors [29–31]. Recently, Boxma, Kella and Yechiali [32] extended the investigation of ASIP models to the case where the gate opening process follows a Markov renewal process and investigated workload distributions in these queueing networks [33]. In a recent study [34], various key performance measures were analyzed, where a general process controls the gate inter-opening times, and particles arrive to the first site between gate openings.

The current work considers an ASIP-type model with a general arrival process as described above. First, we assume in Section 2 that arrivals occur to the first site only, expand the study done in [31] to a G-ASIP, and derive a formula for calculating the site occupancy probabilities in a homogeneous G-ASIP [34]. We show that these probabilities are functions of both the site index, as well as the inward flux rate (λ), thus extending and refining the result presented in [31], where those probabilities are presented as functions of the site index only. Numerical results illustrate the dependence of a site's occupancy probability on the inward flux rate λ . A refined formula for the conditional mean occupancy in each site is also derived. It is further shown that, in the case where the arrival process to the first site is Poisson with rate λ , the following interesting property holds $P(\text{site } k \text{ is occupied} \mid \lambda = 1) = P(\text{site } k + 1 \text{ is occupied} \mid \lambda \rightarrow \infty)$.

For the classical ASIP [28], a tedious successive substitution approach was constructed for the derivation of the PGF of site occupancies; however, due to the complexity of the formulas, explicit expressions were derived for networks with $n = 2$ and $n = 3$ sites only. In a generalized ASIP studied in [32], the PGF was explicitly calculated only for a network with $n = 2$ sites. In Section 3, we assume that arrivals occur to all sites and develop an innovative matrix solution approach to derive the multidimensional PGF of the site occupancies, an approach replacing the tedious successive substitution solution method developed in [28]. An explicit result is derived for the PGF of a G-ASIP with $n = 3$ sites. In addition, the case where the inter-gate opening intervals are Gamma distributed with shape parameter α is thoroughly analyzed and explicit formulas are derived, including the cases when $\alpha = 1$ (Exponential) and when $\alpha \rightarrow \infty$ (Deterministic). In Section 4, we extend the analysis to a system with $n = 4$ sites and calculate its PGF. The main contribution in Section 5 is showing that the set of $\binom{2n}{n}$ linear equations required to obtain the PGF of

an n -site G-ASIP can be first reduced by half into a set of $\binom{2n-1}{n}$ equations, and then further reduced to a set of $2^n - (n + 1)$ equations. The latter set can be additionally split

into several smaller triangular sets. Moreover, a procedure is developed showing how to derive the PGF of an $(n + 1)$ -site G-ASIP from the solution obtained for an n -site network.

Finally, in Section 6, the mean site occupancies, and the mean total load of the first k sites in an n -site system, with arrivals to all sites, are calculated. The results are compared with those of the classical ASIP model [28].

2. Occupation Probabilities

2.1. Introduction

Let X_k be the number of particles occupying site k , ($k = 1, 2, \dots, n$). The probability that site k is occupied had been first studied in [29] for the classical ASIP. It was demonstrated, via Monte Carlo simulations, that the asymptotic occupation probabilities for a homogeneous ASIP where particles arrive at Poisson rate λ to the first site only and where gate k opens, independently of other gates, every Exponential (μ_k) time. When $\lambda = \mu_k = 1$ it was shown that the above probability exhibits a power law decay like $k^{-0.5}$. Namely, at steady state, the probability that site k is occupied decreases like $k^{0.5}$. In a later study [31] this result was analytically calculated, and it was shown (Equation (6)) there that

$$\lim_{k \rightarrow \infty} P(X_k > 0) \approx \frac{1}{\sqrt{\pi k}} \quad (1)$$

It was also argued that Equation (1) is universal in a stronger sense, as it is independent even of the arrival rate λ , where ‘ \approx ’ denotes asymptotic equivalence to leading order in k . This result led to the immediate conclusion that the conditional mean occupancy of site k is given by

$$E[X_k | X_k > 0] \approx \frac{\lambda}{\mu} \sqrt{\pi k} \quad (2)$$

However, we assert that $\lambda \neq 1$, $P(X_k > 0)$ as well as $E[X_k | X_k > 0]$ are functions of the arrival rate λ . First, in the case that $\lambda \rightarrow 0$, the term $P(X_k > 0)$ should tend to 0, whereas the approximation in Equation (1) does not satisfy this condition. Furthermore, the conditional mean occupancy must satisfy that $E[X_k | X_k > 0] \geq 1$ for any arrival rate, in particular when $\lambda \rightarrow 0$. Yet, Equation (2) does not satisfy this quality either.

We claim that, while $P(X_k > 0) \approx \frac{1}{\sqrt{\pi k}}$ is a good approximation for the site occupancies when the inward rate of arrival is $\lambda = 1$, a corrected formula of $P(X_k > 0)$ depends on the income flux λ . We derive an explicit formula for $P(X_k > 0)$ for all cases of λ . As a result, we also adjust Formula (2) and present an explicit result (depending on λ) for the conditional mean occupancy of site k .

Denote by I_k the indicator occupancy of site k . That is, $I_k = \begin{cases} 1 & X_k > 0 \\ 0 & X_k = 0 \end{cases}$, and denote by $P_\lambda(I_k = 1) \equiv P_\lambda(k)$ the two-variable occupancy function (discrete variable k and continuous variable λ). Viewing $P_\lambda(k)$ as a function of two variables gives a deeper insight into this complex model. We show the interesting relation: $P_{\lambda=1}(k) = \lim_{\lambda \rightarrow \infty} P_\lambda(k+1)$, which has two significant implications. First, the use of the latter relation simplifies the calculation of $P_1(k)$ since it releases $P_\lambda(k)$ from its stochastic dependents on the inward flux. The implication of $\lambda \rightarrow \infty$ is that the first site is always occupied. Thus, the probability of site occupancies is dependent on gate openings only. Second, we show that $P_\lambda(k)$ is a monotone increasing function of λ with an upper bound when $\lambda \rightarrow \infty$, which is equal to $P_1(k-1)$. Furthermore, we show that $\frac{1}{\sqrt{\pi(k-1+\frac{1}{\lambda})+\frac{1}{\lambda^2}}}$ is a more accurate approximation for $P_\lambda(k)$ for λ ranging from 0 to infinity and leads to a refined result for the conditional mean occupancy.

2.2. Laws of Motion in an n -Site G-ASIP

Consider an n -site G-ASIP. During the time between two successive gate openings, arrivals occur to the first site only according to a general arrival process. In a steady state, let A denote the number of arrivals to site Q_1 during an inter-gate opening interval, and let

$\hat{A}(z) = E[z^A]$, $|z| \leq 1$, denote the corresponding probability-generating function (PGF). Let X_j , $j = 1, \dots, n$ denote the number of particles (occupancy) in site j right after a gate opening, and let $X_j^{(r)}$ denote the number of particles in Q_j right after the r^{th} gate opening. Let $A^{(r)}$ be the number of arrivals to site j during the interval between the $(r-1)^{th}$ and r^{th} gate openings. Let $G(z_1, \dots, z_n) = E[z_1^{X_1} \dots z_n^{X_n}]$, $|z_j| \leq 1$ be the PGF of the joint probability mass function of the site occupancies at steady state.

The occupancies Law of Motion yields

If gate 1 opens at the $(r+1)^{st}$ gate opening, then:

$$X_1^{(r+1)} = 0, \quad X_2^{(r+1)} = X_1^{(r)} + A^{(r+1)} + X_2^{(r)}, \quad \dots, \quad X_n^{(r+1)} = X_n^{(r)}$$

If gate 2 opens at step $r+1$:

$$X_1^{(r+1)} = X_1^{(r)} + A^{(r+1)}, \quad X_2^{(r+1)} = 0, \quad X_3^{(r+1)} = X_2^{(r)} + X_3^{(r)}, \quad \dots, \quad X_n^{(r+1)} = X_n^{(r)}$$

If gate j opens at step $r+1$:

$$X_1^{(r+1)} = X_1^{(r)} + A^{(r+1)}, \quad X_2^{(r+1)} = X_2^{(r)}, \quad \dots, \quad X_j^{(r+1)} = 0, \quad X_{j+1}^{(r+1)} = X_j^{(r)} + X_{j+1}^{(r)}, \quad \dots, \quad X_n^{(r+1)} = X_n^{(r)}$$

If gate n opens at step $r+1$:

$$X_1^{(r+1)} = X_1^{(r)} + A^{(r+1)}, \quad X_2^{(r+1)} = X_2^{(r)}, \quad X_3^{(r+1)} = X_3^{(r)}, \quad \dots, \quad X_n^{(r+1)} = 0$$

In steady state we obtain

$$\begin{aligned} G(z_1, z_2, \dots, z_n) &= E[z_1^{X_1} \dots z_n^{X_n}] \\ &= p_1 \hat{A}(z_2, z_2, z_3, \dots, z_n) G(z_2, z_2, z_3, \dots, z_n) + p_2 \hat{A}(z_1, z_3, \dots, z_n) G(z_1, z_3, \dots, z_n) \\ &\quad + \dots + p_n \hat{A}(z_1, z_2, \dots, z_{n-1}) G(z_1, z_2, \dots, z_{n-1}) \end{aligned} \quad (3)$$

Let $\vec{z} = (z_1, z_2, \dots, z_n)$, and let $\vec{z}_k = \left(z_1, z_2, \dots, z_{k-1}, \underbrace{z_{k+1}}_{k\text{-th position}}, z_{k+1}, \dots, z_n \right)$, $k = 1, 2, \dots, n-1$ and $\vec{z}_n = (z_1, z_2, \dots, z_{n-1})$. Then, Equation (3) becomes

$$G(\vec{z}) = p_1 \hat{A}(\vec{z}_1) G(\vec{z}_1) + p_2 \hat{A}(\vec{z}_2) G(\vec{z}_2) + \dots + p_n \hat{A}(\vec{z}_n) G(\vec{z}_n) \quad (4)$$

Obtaining an expression for $G(\vec{z})$ allows us to explicitly calculate the occupation probability $P(X_k > 0)$ for any k in an n -size G-ASIP. The optimal allocation of p_j , $j = 1, \dots, n$ with the constraint $\sum_{j=1}^n p_j = 1$ of the G-ASIP was studied in [32,34] and it was shown that a homogeneous G-ASIP where $p_j = \frac{1}{n}$, $j = 1, \dots, n$, is optimal. In what follows, we study the homogeneous G-ASIP.

2.3. Calculation of $P_\lambda(k)$

Utilizing Equation (4) enables a simple calculation of $P_\lambda(k)$ for any inter-gate opening process and any arrival rate. It is simpler to use in the sequel the ‘emptiness function’ $Q_\lambda(k) \equiv P_\lambda(I_k = 0)$.

The site occupancies PGF, $G(z_1, z_2, \dots, z_n)$, can be applied in order to obtain the probability that site k is empty. That is, substituting $z_k = 0$ and $z_j = 1$ for all $j \neq k$ gives

$$G\left(1, \dots, 1, \underbrace{0}_k, 1, \dots, 1\right) = Q_\lambda(k).$$

Theorem 1. In a homogeneous n -site G -ASIP where arrivals occur at the first site only, the probability that site k is empty is

$$Q_\lambda(1) = \frac{1}{n - (n-1)\hat{A}(0)}$$

$$Q_\lambda(k) = \left(2^{k-2}, 2^{k-3}, \dots, 1, \frac{\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \right) \prod_{j=2}^k \frac{1}{2^{j-1}} C_{k,j} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ Q_\lambda(1) \end{pmatrix}, \quad k \geq 2$$

where,

$$C_{k,j} = \begin{pmatrix} 2^{j-2} & 0 & \dots & & & & 0 \\ 0 & \ddots & 0 & \dots & & & \vdots \\ \vdots & 0 & 2^{j-2} & 0 & \dots & & 0 \\ & \vdots & 0 & 2^{j-3} & \dots & 2 & 1 & \frac{\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \\ & & \vdots & \ddots & \ddots & 4 & 2 & \frac{2\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \\ & & & & \ddots & \vdots & \vdots & \vdots \\ & & & & & 2^{j-3} & \frac{2^{j-3}\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \\ 0 & \dots & & & & & 0 & \frac{2^{j-2}\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \end{pmatrix}$$

To simplify, setting

$$C_j = \begin{pmatrix} 2^{j-3} & \dots & 2 & 1 \\ 0 & \ddots & & 2 \\ \vdots & \ddots & & \vdots \\ 0 & \dots & 0 & 2^{j-3} \end{pmatrix} \rightarrow \vec{A}_j = \begin{pmatrix} \frac{\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \\ \frac{2\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \\ \vdots \\ \frac{2^{j-2}\hat{A}^2(0)}{n - (n-1)\hat{A}(0)} \end{pmatrix}$$

gives a compact presentation

$$C_{k,j} = \begin{pmatrix} 2^{k-2}I_{k-j} & 0 & 0 \\ 0 & C_j & \vec{A}_j \\ 0 & 0 & \vdots \end{pmatrix}$$

Proof. Let $\vec{0}_{(n,k)} = \left(\underbrace{1, 1, \dots, 0, \dots, 0}_n \right)$, then substituting $\vec{z} = \vec{0}_{(n,n-1)}$ in Equation (4) yields

$$G(0) = Q_\lambda(1) = \frac{1}{n - (n-1)\hat{A}(0)}$$

By substituting $\vec{z} = \vec{0}_{(n,k)}, k = 1, \dots, n$ in Equation (4), a linear set is obtained

$$\begin{pmatrix} 2^{n-2} & -1 & 0 & \cdots & 0 \\ 0 & 2^{n-2} & -1 & & \vdots \\ \vdots & \ddots & \ddots & -1 & 0 \\ & & 2^{n-2} & & -\hat{A}(0) \\ 0 & \cdots & 0 & n - (n-1)\hat{A}(0) & \end{pmatrix} \begin{pmatrix} G(\vec{0}_{(n,1)}) \\ G(\vec{0}_{(n,2)}) \\ \vdots \\ G(\vec{0}_{(n,n)}) \end{pmatrix} = \begin{pmatrix} 1 \\ G(\vec{0}_{(n-1,1)}) \\ \vdots \\ \hat{A}(0)G(\vec{0}_{(n-1,n-1)}) \end{pmatrix}$$

The above yields a recursive formula,

$$\begin{pmatrix} G(\vec{0}_{(n,1)}) \\ G(\vec{0}_{(n,2)}) \\ \vdots \\ G(\vec{0}_{(n,n)}) \end{pmatrix} = \frac{1}{2^{n-1}} C_n \begin{pmatrix} 1 \\ G(\vec{0}_{(n-1,1)}) \\ \vdots \\ G(\vec{0}_{(n-1,n-1)}) \end{pmatrix}$$

By using the initial condition $Q_\lambda(1) = \frac{1}{n - (n-1)\hat{A}(0)}$, the proof is complete. \square

Theorem 2. In a homogeneous n -site G -ASIP where arrivals occur at the first site only with $\lambda \rightarrow \infty$, the emptiness probabilities are given by

$$Q_\lambda(1) = \frac{1}{n}$$

$$Q_\lambda(k) = (2^{k-2}, \dots, 4, 2, 1) \prod_{j=2}^k \frac{1}{2^{j-1}} C_{k,j} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} + \frac{1}{2}, \quad k \geq 2$$

where,

$$C_{k,j} = \begin{pmatrix} 2^{k-2} I_{k-j} & 0 \\ 0 & C_j \end{pmatrix}, \quad C_j = \begin{pmatrix} 2^{j-3} & \cdots & 2 & 1 \\ 0 & \ddots & & 2 \\ \vdots & \ddots & & \vdots \\ 0 & \cdots & 0 & 2^{j-3} \end{pmatrix}$$

Proof. Applying Theorem 1 when $\lambda \rightarrow \infty$ (namely, the probability of no arrivals between two gate openings tends to zero) leads to $\lim_{\lambda \rightarrow \infty} \hat{A}(0)(\lambda) = 0$, which leads to

$$Q_\lambda(1) = \frac{1}{n}$$

$$Q_\lambda(k) = (2^{k-2}, \dots, 4, 2, 1) \prod_{j=2}^k \frac{1}{2^{j-1}} C_j \begin{pmatrix} 1 \\ 1 \\ \vdots \\ \frac{1}{n} \end{pmatrix}$$

where,

$$C_{k,j} = \begin{pmatrix} 2^{k-2} I_{k-j} & 0 \\ 0 & C_j \end{pmatrix}, C_j = \begin{pmatrix} 2^{j-3} & \dots & 2 & 1 \\ 0 & \ddots & & 2 \\ \vdots & \ddots & & \vdots \\ 0 & \dots & 0 & 2^{j-3} \end{pmatrix}$$

The proof is complete. \square

The result $Q_\lambda(1) = \frac{1}{n}$ deserves further explanation: When $\lambda \rightarrow \infty$, an unlimited number of particles accumulate in front of Q_1 . Right after the opening of gate 1, Q_1 becomes empty, and will stay empty until another gate opening occurs. At that instant, all accumulated particles will move to Q_1 . At the next gate opening Q_1 opens (and becomes empty) with probability $\frac{1}{n}$, which leads to the above result.

2.4. Calculation of $P_\lambda(k)$ for the Classical ASIP

We now derive an explicit formula for the site occupancy probabilities for the case when gate k opens independently every Exponential ($\mu_k = 1$) time, $k = 1, 2, \dots, n$, ($\mu = \sum_{k=1}^n \mu_k = n$) and the arrival is Poisson (λ). We further show that $\lim_{\lambda \rightarrow \infty} P_\lambda(k+1) = P_1(k)$.

Substituting $\hat{A}(0) = \frac{\mu}{\mu+\lambda} = \frac{n}{n+\lambda}$ in Theorem 1 yields

$$Q_\lambda(1) = \frac{n+\lambda}{n(\lambda+1)}$$

$$Q_\lambda(k) = \left(2^{k-2}, \dots, 2, 1, \frac{n}{(\lambda+1)(\lambda+n)} \right) \prod_{j=2}^k \frac{1}{2^{j-1}} C_j \begin{pmatrix} 1 \\ 1 \\ \vdots \\ \frac{n+\lambda}{n(\lambda+1)} \end{pmatrix}, \quad k \geq 2$$

where,

$$C_j = \begin{pmatrix} 2^{j-2} & 0 & \dots & & & & 0 \\ 0 & \ddots & 0 & \dots & & & \vdots \\ \vdots & 0 & 2^{j-2} & 0 & \dots & & 0 \\ & \vdots & 0 & 2^{j-3} & \dots & 2 & 1 & \frac{n}{(\lambda+1)(\lambda+n)} \\ & & \vdots & \ddots & \ddots & 4 & 2 & \frac{2n}{(\lambda+1)(\lambda+n)} \\ & & & & \ddots & \vdots & \vdots & \vdots \\ & & & & & 2^{j-3} & \frac{2^{j-3}n}{(\lambda+1)(\lambda+n)} \\ 0 & \dots & & & & 0 & \frac{2^{j-2}}{\lambda+1} \end{pmatrix}$$

To illustrate, we calculate the occupancy probabilities of the first five sites in Appendix A and get:

$$Q_\lambda(1) = \frac{n+\lambda}{n(\lambda+1)}, \quad Q_\lambda(2) = \frac{1}{2} \left(1 + \frac{\lambda+1}{(\lambda+1)^3} \right), \quad Q_\lambda(3) = \frac{1}{8} \left(5 + \frac{\lambda+3}{(\lambda+1)^3} \right),$$

$$Q_\lambda(4) = \frac{1}{16} \left(11 + \frac{\lambda+3}{(\lambda+1)^3} + \frac{2}{(\lambda+1)^4} \right)$$

$$Q_\lambda(5) = \frac{1}{128} \left(93 + \frac{2(\lambda+5)}{(\lambda+1)^5} + \frac{10(\lambda+2)}{(\lambda+1)^4} + \frac{5}{(\lambda+1)^2} \right)$$

It is clearly seen that the site occupancy probabilities are functions of the inward flux and have an upper limit for the probability $P_\lambda(k)$ as $\lambda \rightarrow \infty$. This follows, since $Q_\lambda(k)$ is a monotone decreasing function of λ for every k . It is readily seen that $\lim_{\lambda \rightarrow \infty} P_\lambda(2) = \frac{1}{2}$, $\lim_{\lambda \rightarrow \infty} P_\lambda(3) = \frac{3}{8}$, $\lim_{\lambda \rightarrow \infty} P_\lambda(4) = \frac{5}{16}$ and $\lim_{\lambda \rightarrow \infty} P_\lambda(5) = \frac{35}{128}$. On the other hand, as λ becomes small and tends to zero, $\lim_{\lambda \rightarrow 0} P_\lambda(k) = 0$ for all k .

Studying the probabilities $Q_\lambda(k)$ as functions λ , we observe that direct substitution in $Q_\lambda(k)$ of $\lambda = 1$ and of $\lambda \rightarrow \infty$ in the first six sites, leads to an interesting relationship, as depicted in Table 2, namely $\lim_{\lambda \rightarrow \infty} P_\lambda(k+1) = P_1(k)$, $k \geq 2$.

Table 2. $Q_\lambda(k)$ when $\lambda = 1$ and when $\lambda \rightarrow \infty$ for the first six sites.

$Q_\lambda(k)$	$\lambda = 1$	$\lambda \rightarrow \infty$
$k = 1$	$\frac{n+1}{2n}$	$\frac{1}{n}$
$k = 2$	$\frac{5}{8}$	$\frac{1}{2}$
$k = 3$	$\frac{11}{16}$	$\frac{5}{8}$
$k = 4$	$\frac{93}{128}$	$\frac{11}{16}$
$k = 5$	$\frac{722}{1024}$	$\frac{93}{128}$
$k = 6$	$\frac{25,376}{32,768}$	$\frac{722}{1024}$

Figure 2 below depicts the behavior of $P_\lambda(k)$ (abbreviated $P(k)$) as function of λ for $k = 2, 3, 4, 5$, and exhibits the corresponding upper bound.

The relation $\lim_{\lambda \rightarrow \infty} P_\lambda(k+1) = P_1(k)$ simplifies the calculation of $P_1(k)$ since it releases $P_\lambda(k)$ from its dependents on the inward flux. The implication of $\lambda \rightarrow \infty$ is that the first site is always occupied and allows the use of Theorem 2 to calculate $P_1(k)$.

While it was demonstrated in [31] that $\frac{1}{\sqrt{\pi k}}$ is a good approximation for the site occupancy probabilities when the number of sites become large, independent of λ , Monte Carlo simulations show that for $0 < \lambda < 1$ the occupancy probabilities graph fluctuates between 0 and $\frac{1}{\sqrt{\pi k}}$. For $\lambda > 1$ and as λ tends to infinity, the occupancy graph sways away from $\frac{1}{\sqrt{\pi k}}$.

Following extensive simulations of the G-ASIP model with varying arrival rates λ , we propose a refined and better approximation for $P_\lambda(k)$, namely, $P_\lambda(k) \approx \frac{1}{\sqrt{\pi(k-(1-\frac{1}{\lambda})) + \frac{1}{\lambda^2}}}$.




Figure 2. $P_\lambda(k)$ for $k = 2, 3, 4, 5$. Horizontal lines depict the limiting values $\lim_{\lambda \rightarrow \infty} P_\lambda(k+1)$ and their interactions with $P_1(k)$.

In the following set of results obtained using Monte Carlo simulations, we compare the two approximations for five different arrival rates λ . Figure 3 below shows that for arrival rates $\lambda = 8$, $\lambda = 4$, $\lambda = 1$, $\lambda = \frac{1}{2}$ and $\lambda = \frac{1}{4}$ the simulated occupancy probabilities is better approximated by $\frac{1}{\sqrt{\pi(k - (1 - \frac{1}{\lambda})) + \frac{1}{\lambda^2}}}$ rather than $\frac{1}{\sqrt{\pi k}}$.

Figure 4 shows that the suggested approximation of $P_\lambda(k)$ is an increasing function of the variable λ as should be expected, and it is a decreasing function of the index k . It also shows, as its contours suggest, the very interesting property that $P_\infty(k+1) = P_1(k)$. For example, when $\lambda = 1$ the probability that site $k = 4$ is occupied is $P_1(4) = \frac{35}{128} \approx 0.273$, while the approximated value is $\frac{1}{\sqrt{4\pi+1}} \approx 0.2715$.

With a better approximation for $P_\lambda(k)$, we now present a refined formula for the conditional mean occupancy of site k , namely,

$$E[X_k | I_k = 1] \approx \lambda \sqrt{\pi \left(k - \left(1 - \frac{1}{\lambda} \right) \right) + \frac{1}{\lambda^2}} \quad (5)$$




Figure 3. Cont.




Figure 3. Monte Carlo simulations comparing the approximations $\frac{1}{\sqrt{\pi(k-(1-\frac{1}{\lambda})) + \frac{1}{\lambda^2}}}$ and $\frac{1}{\sqrt{\pi k}}$. As shown by the simulations, $\frac{1}{\sqrt{\pi k}}$ is a good approximation when $\lambda = 1$ but it loses its accuracy as λ shifts away from 1. The two-dimensional function $P_\lambda(k) \approx \frac{1}{\sqrt{\pi(k-(1-\frac{1}{\lambda})) + \frac{1}{\lambda^2}}}$ is better suited to approximate the site's occupancy probabilities.




Figure 4. The two-dimensional function $P_\lambda(k) = \frac{1}{\sqrt{\pi(k-(1-\frac{1}{\lambda})) + \frac{1}{\lambda^2}}}$.

Equation (10) in [31] claims that $E[X_k|I_k = 1] = E[X_k|X_k > 0] \approx \frac{\lambda}{\mu_k} \sqrt{\pi k}$, implying that, when $\mu_k = 1$, $E[X_k|I_k = 1] = \lambda \sqrt{\pi k}$. However, Equation (10) infers that $\lim_{\lambda \rightarrow 0} E[X_k|X_k > 0] = 0$, which cannot be true since the conditional mean occupancy of site k must satisfy $E[X_k|I_k = 1] \geq 1$ for all $\lambda > 0$. If the Poisson arrival rate is low, it should be expected that most sites will be vacant, and those who are occupied will be occupied by

a single particle only. The limit $\lim_{\lambda \rightarrow 0} E[X_k | X_k > 0] = 1$ is to be expected. Namely, the limit $\lim_{\lambda \rightarrow 0} [X_k | I_k = 1] = \lim_{\lambda \rightarrow 0} \frac{\lambda}{P_\lambda(k)} = 1$ should be obtained. This condition demands that for very small λ ($\lambda \ll 1$) the occupancy probability function should satisfy $P_\lambda(k) \sim \lambda$.

In conclusion, equipped with analytical calculations, as well as with Monte Carlo observations, we can claim that the function $P_\lambda(k)$ must satisfy:

1. $P_\infty(k+1) = P_1(k)$.
2. $\lim_{\lambda \rightarrow 0} P_\lambda(k) = \lambda$.
3. $P_\lambda(k)$ decreases like $k^{0.5}$.

Our suggested corrected two-variable site occupancy function approximation is

$$P_\lambda(k) \approx \frac{1}{\sqrt{\pi \left(k - \left(1 - \frac{1}{\lambda} \right) \right) + \frac{1}{\lambda^2}}}$$

This approximation is also valid for smaller values of k and satisfies conditions 1, 2 and 3 above.

This leads to the corrected conditional mean occupancy

$$E[X_k | I_k = 1] \approx \lambda \sqrt{\pi \left(k - \left(1 - \frac{1}{\lambda} \right) \right) + \frac{1}{\lambda^2}}$$

2.5. Site Occupancy Probabilities When Arrivals Occur to All Sites

Consider a 3-site G-ASIP where arrivals occur to all sites. The site occupancy probabilities are (the derivation is given in Appendix B).

$$P(I_1 = 0) = \frac{p_1}{1 - (p_2 + p_3)\hat{A}(0)}$$

$$P(I_2 = 0) = \frac{p_1^2 p_2 \hat{A}(0) \hat{A}(0,0)}{\left(1 - (p_1 + p_3)\hat{A}(0,0)\right) \left(1 - (p_2 + p_3)\hat{A}(0)\right) \left(1 - p_3 \hat{A}(1,0)\right)} + \frac{p_2}{1 - p_3 \hat{A}(1,0)}$$

$$P(I_3 = 0) = \frac{p_3}{1 - p_1 \hat{A}(1,1,0)} + \frac{p_1^2 p_2^2 p_3 \hat{A}(0) \hat{A}(0,0) \hat{A}(0,0,0) \hat{A}(1,0,0)}{\left(1 - p_1 \hat{A}(1,1,0)\right) \left(1 - p_2 \hat{A}(1,0,0)\right) \left(1 - (p_1 + p_2) \hat{A}(0,0,0)\right) \left(1 - (p_2 + p_3) \hat{A}(0)\right) \left(1 - (p_1 + p_3) \hat{A}(0,0)\right)} \\ + \frac{p_2 \hat{A}(1,0,0) p_3 \hat{A}(1,0) \left(\frac{p_1^2 p_2 \hat{A}(0) \hat{A}(0,0)}{\left(1 - (p_1 + p_3) \hat{A}(0,0)\right) \left(1 - (p_2 + p_3) \hat{A}(0)\right) \left(1 - p_3 \hat{A}(1,0)\right)} + \frac{p_2}{1 - p_3 \hat{A}(1,0)} \right)}{\left(1 - p_1 \hat{A}(1,1,0)\right) \left(1 - p_2 \hat{A}(1,0,0)\right)}$$

The construction of the set of equations for calculating site occupancy probabilities for a 4-site G-ASIP where arrivals occur to all sites are presented in Appendix C.

3. Site Occupancy PGF in a G-ASIP with Arrivals to All Sites: $n = 3$

As indicated in the introduction, the PGF of the joint probability mass function of the site occupancies of a G-ASIP network was calculated in [32] only for the case of $n = 2$. In this section we extend the analysis and calculate the PGF for a 3-site G-ASIP. Arrivals occur at all sites Q_1, Q_2 and Q_3 during the time between two successive gate openings with a general arrival process. In a steady state, let $A_j, j = 1, 2, 3$ denote the number of arrivals to site Q_j during an inter-gate opening interval and let $\hat{A}(z_1, z_2, z_3) = E[z_1^{A_1} z_2^{A_2} z_3^{A_3}]$ denote the corresponding joint PGF. Let $X_j, j = 1, 2, 3$ denote the number of particles (occupancy)

in site j right after a gate opening, and let $X_j^{(r)}$ denote the number of particles in Q_j right after the r^{th} gate opening. Let $A_j^{(r)}$ be the number of arrivals to site j during the interval between the $(r-1)^{th}$ and r^{th} gate openings. Let $G(z_1, z_2, z_3) = E[z_1^{X_1} z_2^{X_2} z_3^{X_3}]$, $|z_j| \leq 1$ be the PGF of the joint probability mass function of the sites' occupancies at steady state.

3.1. Successive Substitutions

The laws of motion of an n -site G-ASIP were presented in Section 2.2. Substituting $n = 3$ yields

$$G(\vec{z}) = p_1 \hat{A}(\vec{z}_1) G(\vec{z}_1) + p_2 \hat{A}(\vec{z}_2) G(\vec{z}_2) + p_3 \hat{A}(\vec{z}_3) G(\vec{z}_3) \quad (6)$$

The PGF of the joint probability mass function of the site occupancies of a 3-site classical ASIP was calculated in [28] using a successive substitution method. The tedious successive substitution procedure to calculate $G(z_1, z_2, z_3)$ is detailed in Appendix C. The final result is:

$$\begin{aligned}
G(z_1, z_2, z_3) = & \left(\frac{p_1^2 p_2^2 p_3 \hat{A}(z_3, z_3, z_3) \hat{A}(z_3, z_3) \hat{A}(z_3) \hat{A}(z_2, z_3, z_3)}{\left(\left(1 - (p_2 + p_3) \hat{A}(z_3) \right) \right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3) \right) \left(1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3) \right)} \right. \\
& \left. \frac{\left(1 - p_1 \hat{A}(z_2, z_2, z_3) \right) \left(1 - p_2 \hat{A}(z_2, z_3, z_3) \right)}{p_1^2 p_2^2 p_3 \hat{A}(z_2, z_3) \hat{A}(z_3) \hat{A}(z_3, z_3) \hat{A}(z_2, z_3, z_3)} + \frac{p_1 p_2^2 p_3 \hat{A}(z_2, z_3) \hat{A}(z_2) \hat{A}(z_2, z_3, z_3)}{\left(1 - (p_1 + p_3) \hat{A}(z_3, z_3) \right) \left(1 - (p_2 + p_3) \hat{A}(z_3) \right) 1 - (p_2 + p_3) \hat{A}(z_2)} \right. \\
& \left. + \frac{\left(1 - p_1 \hat{A}(z_2, z_2, z_3) \right) \left(1 - p_2 \hat{A}(z_2, z_3, z_3) \right) \left(1 - p_3 \hat{A}(z_2, z_3) \right)}{\left(1 - p_1 \hat{A}(z_2, z_2, z_3) \right) \left(1 - p_2 \hat{A}(z_2, z_3, z_3) \right) \left(1 - p_3 \hat{A}(z_2, z_3) \right)} \right. \\
& \left. + \frac{\frac{p_1 p_2 p_3 \hat{A}(z_2, z_2) \hat{A}(z_2)}{\left(1 - (p_1 + p_3) \hat{A}(z_2, z_2) \right) \left(1 - (p_2 + p_3) \hat{A}(z_2) \right)}}{1 - p_1 \hat{A}(z_2, z_2, z_3)} \right) \\
& + \\
& p_2 \hat{A}(z_1, z_3, z_3) \left(\frac{p_1^2 p_2 p_3 \hat{A}(z_3, z_3, z_3) \hat{A}(z_3, z_3) \hat{A}(z_3)}{\left(1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3) \right) \left(\left(1 - (p_2 + p_3) \hat{A}(z_3) \right) \right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3) \right) \left(1 - p_2 \hat{A}(z_1, z_3, z_3) \right)} \right. \\
& \left. + \frac{\frac{p_1^2 p_2 p_3 \hat{A}(z_1, z_3) \hat{A}(z_3) \hat{A}(z_3, z_3)}{\left(1 - (p_1 + p_3) \hat{A}(z_3, z_3) \right) \left(1 - (p_2 + p_3) \hat{A}(z_3) \right)} + \frac{p_1 p_2 p_3 \hat{A}(z_1, z_3) \hat{A}(z_1)}{1 - (p_2 + p_3) \hat{A}(z_1)}}{\left(1 - p_3 \hat{A}(z_1, z_3) \right) \left(1 - p_2 \hat{A}(z_1, z_3, z_3) \right)} \right) \\
& + \\
& p_3 \hat{A}(z_1, z_2) \left(\frac{\frac{p_1^2 p_2 \hat{A}(z_2) \hat{A}(z_2, z_2)}{\left(1 - (p_1 + p_3) \hat{A}(z_2, z_2) \right) \left(1 - (p_2 + p_3) \hat{A}(z_2) \right)} + \frac{p_1 p_2 \hat{A}(z_1)}{1 - (p_2 + p_3) \hat{A}(z_1)}}{1 - p_3 \hat{A}(z_1, z_2)} \right)
\end{aligned} \tag{7}$$

3.2. Matrix Representation

3.2.1. Linear Equations

We argue and show that the tedious successive substitution process used to derive the explicit solution of Equation (7), as depicted by the tree in Figure 2, can be modified and simplified with a formulated matrix representation approach leading to a set of linear equations,

Proposition 1. *There are 20 distinct variables $G(z_i, z_j, z_k)$, $1 \leq i \leq j \leq k \leq 4$, $z_4 := 1$ in Equation (6).*

Proof. Consider the 3 variables z_i, z_j, z_k , as well as the digit 1, such that $1 \leq i \leq j \leq k \leq 4$. The number of vertices and leaves that exist in the set

$$\{G(z_i, z_j, z_k) \mid 1 \leq i \leq j \leq k \leq 4, \quad z_4 := 1\}$$

is equal to the number of ways one can create distinct 3-digit numbers from the digits $\{1, 2, 3, 4\}$ with repetition and in increasing order. There are $D(4, 3) = \binom{4-1+3}{4-1} = \binom{6}{3} = 20$ possible ways. \square

With a straightforward substitution in Equation (6), one can generate 20 linear equations of the variables $G(z_i, z_j, z_k)$, $1 \leq i \leq j \leq k \leq 4$, $z_4 := 1$.

The solution derived from this set can then be substituted in Equation (6) to finalize the process.

Still, solving a 20×20 linear equations set can be tedious in itself. Thus, we take a step forward and present the following improving approach.

Proposition 2. *There are only 10 variables needed to derive $\{G(z_i, z_j, z_k) \mid 1 \leq i \leq j \leq k \leq 3\}$.*

Proof. The number of vertices and leaves that exist in the set

$$\{G(z_i, z_j, z_k) \mid 1 \leq i \leq j \leq k \leq 3\}$$

is equal to the number of ways one can create a 3-digit number from the digits $\{1, 2, 3\}$ with repetition and in increasing order. There are $D(3, 3) = \binom{5}{2} = 10$ possible ways. \square

3.2.2. Solution

We bring to use Equation (6):

$$G(z_1, z_2, z_3) = p_1 \hat{A}(z_2, z_2, z_3) G(z_2, z_2, z_3) + p_2 \hat{A}(z_1, z_3, z_3) G(z_1, z_3, z_3) + p_3 \hat{A}(z_1, z_2) G(z_1, z_2)$$

and apply the following eight substitutions, which yield eight linear equations as follows:

$$\begin{aligned} z_2 = 1, z_3 = 1 &\rightarrow G(z_1) \\ z_1 = z_2, z_2 = 1, z_3 = 1 &\rightarrow G(z_2) \\ z_2 = z_1, z_3 = 1 &\rightarrow G(z_1, z_1) \\ z_1 = z_2, z_3 = 1 &\rightarrow G(z_2, z_2) \\ z_3 = 1 &\rightarrow G(z_1, z_2) \\ z_2 = z_1, z_3 = 1 &\rightarrow G(z_1, z_1, z_2) \\ z_3 = z_2 &\rightarrow G(z_1, z_2, z_2) \\ z_1 = z_2, z_3 = z_2 &\rightarrow G(z_2, z_2, z_2) \end{aligned}$$

Consequently, we obtain the following set of eight linear equations:

$$\begin{aligned}
G(z_1) &= p_1 + p_2 \hat{A}(z_1)G(z_1) + p_3 \hat{A}(z_1)G(z_1) \\
G(z_2) &= p_1 + p_2 \hat{A}(z_2)G(z_2) + p_3 \hat{A}(z_2)G(z_2) \\
G(z_1, z_1) &= p_1 \hat{A}(z_1, z_1)G(z_1, z_1) + p_2 \hat{A}(z_1)G(z_1) + p_3 \hat{A}(z_1, z_1)G(z_1, z_1) \\
G(z_2, z_2) &= p_1 \hat{A}(z_2, z_2)G(z_2, z_2) + p_2 \hat{A}(z_2)G(z_2) + p_3 \hat{A}(z_2, z_2)G(z_2, z_2) \\
G(z_1, z_2) &= p_1 \hat{A}(z_2, z_2)G(z_2, z_2) + p_2 \hat{A}(z_1)G(z_1) + p_3 \hat{A}(z_1, z_2)G(z_1, z_2) \\
G(z_1, z_1, z_2) &= p_1 \hat{A}(z_1, z_1, z_2)G(z_1, z_1, z_2) + p_2 \hat{A}(z_1, z_2, z_2)G(z_1, z_2, z_2) + p_3 \hat{A}(z_1, z_1)G(z_1, z_1) \\
G(z_1, z_2, z_2) &= p_1 \hat{A}(z_2, z_2, z_2)G(z_2, z_2, z_2) + p_2 \hat{A}(z_1, z_2, z_2)G(z_1, z_2, z_2) + p_3 \hat{A}(z_1, z_2)G(z_1, z_2) \\
G(z_2, z_2, z_2) &= p_1 \hat{A}(z_2, z_2, z_2)G(z_2, z_2, z_2) + p_2 \hat{A}(z_2, z_2, z_2)G(z_2, z_2, z_2) + p_3 \hat{A}(z_2, z_2)G(z_2, z_2)
\end{aligned}$$

The latter set can be solved directly in a matrix form $H \cdot g = p$, where

$$H = \begin{pmatrix}
1 - (p_2 + p_3)\hat{A}(z_1) & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 - (p_2 + p_3)\hat{A}(z_2) & 0 & 0 & 0 & 0 & 0 & 0 \\
-p_2\hat{A}(z_1) & 0 & 1 - (p_1 + p_3)\hat{A}(z_1, z_1) & 0 & 0 & 0 & 0 & 0 \\
0 & -p_2\hat{A}(z_2) & 0 & 1 - (p_2 + p_3)\hat{A}(z_2, z_2) & 0 & 0 & 0 & 0 \\
-p_2\hat{A}(z_1) & 0 & 0 & -p_1\hat{A}(z_2, z_2) & 1 - p_3\hat{A}(z_1, z_2) & 0 & 0 & 0 \\
0 & 0 & -p_3\hat{A}(z_1, z_1) & 0 & 0 & 1 - p_1\hat{A}(z_1, z_1, z_2) & -p_2\hat{A}(z_1, z_2, z_2) & 0 \\
0 & 0 & 0 & 0 & -p_3\hat{A}(z_1, z_2) & 0 & 1 - p_2\hat{A}(z_1, z_2, z_2) & -p_1\hat{A}(z_2, z_2, z_2) \\
0 & 0 & 0 & -p_3\hat{A}(z_2, z_2) & 0 & 0 & 0 & 1 - (p_1 + p_2)\hat{A}(z_2, z_2, z_2)
\end{pmatrix}$$

$$g = \begin{pmatrix} G(z_1) \\ G(z_2) \\ G(z_1, z_1) \\ G(z_2, z_2) \\ G(z_1, z_2) \\ G(z_1, z_1, z_2) \\ G(z_1, z_2, z_2) \\ G(z_2, z_2, z_2) \end{pmatrix} \quad \text{and} \quad p = \begin{pmatrix} p_1 \\ p_1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

In the Section 3.2.3, we further reduce the computational effort by splitting the 8×8 linear set into two subsets of order 2×2 and 3×3 , respectively.

3.2.3. Matrix Representation Simplification

The 8×8 linear set constructed in Section 3.2.2 can further be contracted into two smaller systems of triangular matrices, one of order 2×2 the other of order 3×3 .

$G(z)$ is derived immediately from Equation (6)

$$G(z) = \frac{p_1}{1 - (p_2 + p_3)\hat{A}(z)}, \quad z = z_1, z_2, z_3$$

$G(z_2, z_2)$ and $G(z_1, z_2)$ are obtained from

$$\begin{pmatrix} 1 - (p_1 + p_3)\hat{A}(z_2, z_2) & 0 \\ -p_1\hat{A}(z_2, z_2) & 1 - p_3\hat{A}(z_1, z_2) \end{pmatrix} \begin{pmatrix} G(z_2, z_2) \\ G(z_1, z_2) \end{pmatrix} = \begin{pmatrix} p_2\hat{A}(z_2)G(z_2) \\ p_2\hat{A}(z_1)G(z_1) \end{pmatrix}$$

Finally, $G(z_1, z_1, z_2)$, $G(z_1, z_2, z_2)$ and $G(z_2, z_2, z_2)$ are calculated by

$$\begin{pmatrix} 1 - p_1 \hat{A}(z_1, z_1, z_2) & -p_2 \hat{A}(z_1, z_2, z_2) & 0 \\ 0 & 1 - p_2 \hat{A}(z_1, z_2, z_2) & -p_1 \hat{A}(z_2, z_2, z_2) \\ 0 & 0 & 1 - (p_1 + p_2) \hat{A}(z_2, z_2, z_2) \end{pmatrix} \begin{pmatrix} G(z_1, z_1, z_2) \\ G(z_1, z_2, z_2) \\ G(z_2, z_2, z_2) \end{pmatrix} = \begin{pmatrix} p_3 \hat{A}(z_1, z_1) G(z_1, z_1) \\ p_3 \hat{A}(z_1, z_2) G(z_1, z_2) \\ p_3 \hat{A}(z_2, z_2) G(z_2, z_2) \end{pmatrix}$$

where $G(z_1, z_1)$ is calculated in step 2 by substituting $z_2 = z_1$ in the variable $G(z_2, z_2)$.

Now, with an appropriate substitution in $G(z_1, z_1, z_2)$, the PGF $G(z_2, z_2, z_3)$ is obtained. Similarly, $G(z_1, z_3, z_3)$ is obtained from $G(z_1, z_2, z_2)$. Finally, by substituting the above as well as $G(z_1, z_2)$, in (6), the explicit solution given in Equation (7) is rederived.

Proposition 3. Only eight variables are required to solve $\{G(z_i, z_j, z_k) | 1 \leq i \leq j \leq k \leq 3\}$.

Proof. Suppose one obtains an explicit formula for $G(z_1, z_2, z_2)$. Then, by substituting $z_2 = z_3$, $G(z_1, z_3, z_3)$ is obtained. Furthermore, by substituting $z_1 = z_2$, the PGF $G(z_2, z_2, z_3)$ is given. Thus, for a complete solution, one needs only eight PGF variables, namely, $G(z_1)$, $G(z_1, z_1)$, $G(z_1, z_2)$, $G(z_1, z_1, z_2)$, $G(z_1, z_2, z_2)$, $G(z_2, z_2, z_2)$ together with $G(z_2)$ and $G(z_2, z_2)$.

These eight variables in eight linear equations construct two distinct triangular linear sets of order 3×3 and 2×2 , leading to a unique solution. \square

To summarize, instead of originally having a 20×20 size linear system, one can solve the problem with only eight linear equations that can be further decomposed to smaller subsets.

3.2.4. Arrivals to the First Site Only and Gamma Distributions Inter-Gate Openings Times

In the special case that particles arrive to site Q_1 only, such that the PGF of the number of arrivals per gate opening is given by $\hat{A}(z_1) = \hat{A}(z_1, 1, 1)$, Equation (7) becomes simpler:

$$\begin{aligned} G(z_1, z_2, z_3) = & \frac{\frac{p_1^3 p_2^2 p_3 \hat{A}^3(z_3) \hat{A}^2(z_2)}{(1 - (p_2 + p_3) \hat{A}(z_3)) (1 - (p_1 + p_3) \hat{A}(z_3)) (1 - (p_1 + p_2) \hat{A}(z_3))} + \frac{p_1^2 p_2 p_3 \hat{A}^3(z_2) (1 - p_3 \hat{A}(z_2) - p_1 p_2 \hat{A}^2(z_2))}{(1 - (p_1 + p_3) \hat{A}(z_2)) (1 - (p_2 + p_3) \hat{A}(z_2))}}{(1 - p_1 \hat{A}(z_2)) (1 - p_2 \hat{A}(z_2)) (1 - p_3 \hat{A}(z_2))} \\ & + \frac{\frac{p_1^2 p_2^2 p_3 \hat{A}^3(z_3) \hat{A}(z_1)}{(1 - (p_2 + p_3) \hat{A}(z_3)) (1 - (p_1 + p_3) \hat{A}(z_3)) (1 - (p_1 + p_2) \hat{A}(z_3))} + \frac{p_1 p_2^2 p_3 \hat{A}^3(z_1)}{1 - \hat{A}(z_1) (p_2 + p_3)}}{(1 - p_2 \hat{A}(z_1)) (1 - p_3 \hat{A}(z_1))} \\ & + \frac{\frac{p_1^2 p_2 p_3 \hat{A}^2(z_2) \hat{A}(z_1)}{(1 - \hat{A}(z_2) (p_2 + p_3)) (1 - \hat{A}(z_2) (p_1 + p_3))} + \frac{p_1 p_2 p_3 \hat{A}^2(z_1)}{1 - \hat{A}(z_1) (p_2 + p_3)}}{1 - p_3 \hat{A}(z_1)} \end{aligned} \quad (8)$$

We now calculate explicit results for the wide family of Gamma distributed inter-opening times.

The Family of Gamma Distributions as the Inter-Gate Openings Times

Suppose that the time O between two successive gate openings is Gamma distributed, $\Gamma(\alpha, \alpha\mu)$, with density $f(t) = \frac{e^{-\alpha\mu t} (\alpha\mu)^\alpha t^{\alpha-1}}{\Gamma(\alpha)}$ and mean $E[O] = \frac{\alpha}{\alpha\mu} = \frac{1}{\mu}$. We assume that the arrival process during an inter-opening time interval is composed of three in-

dependent Poisson processes, each with intensity λ_i to site i . Then, the PGF is given by

$$\hat{A}(z_1, z_2, z_3) = \left(\frac{\alpha\mu}{\alpha\mu + \sum_{i=1}^3 \lambda_i(1-z_i)} \right)^\alpha.$$

Substituting the above in Equation (7) for the 20 possible values of $\hat{A}(z_1, z_2, z_3)$ leads to an explicit expression for the PGF $G(z_1, z_2, z_3)$.

If particles arrive to site Q_1 only, then (denoting $\lambda_1 = \lambda$), $\hat{A}(z) = \left(\frac{\alpha\mu}{\alpha\mu + \lambda(1-z)} \right)^\alpha$, so that

$$G(z_1, z_2, z_3) = \left(\begin{aligned} & \left(\frac{p_1^3 p_2^2 p_3 \frac{(\alpha\mu)^{3\alpha}}{(\alpha\mu + \lambda(1-z_3))^{3\alpha}} \frac{(\alpha\mu + \lambda(1-z_2))^\alpha}{(\alpha\mu)^\alpha}}{\left(1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_3))^\alpha}\right) \left(1 - \frac{(p_1+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_3))^\alpha}\right) \left(1 - \frac{(p_1+p_2)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_3))^\alpha}\right)} + \frac{p_1^2 p_2 p_3 \left(1 - \frac{p_3(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_2))^\alpha} - \frac{p_1 p_2 (\alpha\mu)^{2\alpha}}{(\alpha\mu + \lambda(1-z_2))^{2\alpha}}\right)}{\left(1 - \frac{(p_1+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_2))^\alpha}\right) \left(1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_2))^\alpha}\right)} \right. \\ & \left. \frac{\left(1 + \frac{\lambda(1-z_2)}{\alpha\mu}\right)^\alpha - p_1}{\left(1 + \frac{\lambda(1-z_2)}{\alpha\mu}\right)^\alpha - p_2} \frac{\left(1 + \frac{\lambda(1-z_2)}{\alpha\mu}\right)^\alpha - p_3}{\left(1 + \frac{\lambda(1-z_2)}{\alpha\mu}\right)^\alpha - p_3} \right) \\ & + \left(\frac{p_1^2 p_2^2 p_3 \frac{(\alpha\mu)^{3\alpha}}{(\alpha\mu + \lambda(1-z_3))^{3\alpha}}}{\left(1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_3))^\alpha}\right) \left(1 - \frac{(p_1+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_3))^\alpha}\right)} + \frac{p_1 p_2^2 p_3 \frac{(\alpha\mu)^{2\alpha}}{(\alpha\mu + \lambda(1-z_1))^{2\alpha}}}{1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_1))^\alpha}} \right. \\ & \left. \frac{\left(\frac{(\alpha\mu + \lambda(1-z_1))^\alpha}{(\alpha\mu)^\alpha} - p_2\right) \left(1 - \frac{p_3(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_1))^\alpha}\right)}{\left(\frac{(\alpha\mu + \lambda(1-z_1))^\alpha}{(\alpha\mu)^\alpha} - p_3\right)} \right) \\ & + \left(\frac{p_1^2 p_2 p_3 \frac{(\alpha\mu)^{2\alpha}}{(\alpha\mu + \lambda(1-z_2))^{2\alpha}}}{\left(1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_2))^\alpha}\right) \left(1 - \frac{(p_1+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_2))^\alpha}\right)} + \frac{p_1 p_2 p_3 \frac{(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_1))^\alpha}}{1 - \frac{(p_2+p_3)(\alpha\mu)^\alpha}{(\alpha\mu + \lambda(1-z_1))^\alpha}} \right. \\ & \left. \frac{\left(\frac{(\alpha\mu + \lambda(1-z_1))^\alpha}{(\alpha\mu)^\alpha} - p_3\right)}{\left(\frac{(\alpha\mu + \lambda(1-z_1))^\alpha}{(\alpha\mu)^\alpha} - p_3\right)} \right) \end{aligned} \right) \quad (9)$$

Note that Equation (9) generalizes the result obtained in [28] where $\hat{A}(z) = \frac{\mu}{\mu + \lambda(1-z)}$ is the PGF of the Poisson (λ) number of arrivals to Q_1 during an Exponential inter-opening time with rate $\mu = \mu_1 + \mu_2 + \mu_3$. In this case $p_i = \frac{\mu_i}{\mu}$, so that $p_i \hat{A}(z_j) = \frac{\mu_i}{\mu + \lambda(1-z_j)}$. Similarly, setting $\alpha = 1$ in (9) yields Equation (40) in [28].

Deterministic Inter-Opening Times

When $\alpha \rightarrow \infty$, the gates inter-opening time follows the Deterministic distribution. If arrivals occur to the first site only the PGF $\hat{A}(z) = \left(1 + \frac{\lambda(1-z)}{\alpha\mu}\right)^{-\alpha}$ becomes $\hat{A}(z) = e^{-\frac{\lambda(1-z)}{\mu}}$ and Equation (9) becomes

$$\begin{aligned}
G(z_1, z_2, z_3) = & \frac{p_1^2 p_2 p_3 e^{\frac{\lambda(1-z_2)}{\mu}}}{\left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_2 + p_3)\right) \left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_1 + p_3)\right) \left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_1 + p_2)\right)} + \frac{p_1^2 p_2 p_3 \left(e^{\frac{2\lambda(1-z_2)}{\mu}} - p_3 - p_1 p_2\right)}{\left(e^{\frac{\lambda(1-z_2)}{\mu}} - (p_1 + p_3)\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - (p_2 + p_3)\right)} \\
= & \frac{\left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_1\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_2\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_3\right)}{\left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_1\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_2\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - p_3\right)} \\
& + \frac{p_1^2 p_2 p_3 e^{\frac{\lambda(1-z_1)}{\mu}}}{\left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_2 + p_3)\right) \left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_1 + p_3)\right) \left(e^{\frac{\lambda(1-z_3)}{\mu}} - (p_1 + p_2)\right)} + \frac{p_1 p_2^2 p_3}{e^{\frac{\lambda(1-z_1)}{\mu}} - (p_2 + p_3)} \\
& \frac{\left(e^{\frac{\lambda(1-z_1)}{\mu}} - p_2\right) \left(e^{\frac{\lambda(1-z_1)}{\mu}} - p_3\right)}{\left(e^{\frac{\lambda(1-z_1)}{\mu}} - p_2\right) \left(e^{\frac{\lambda(1-z_1)}{\mu}} - p_3\right)} \\
& + \frac{p_1^2 p_2 p_3}{\left(e^{\frac{\lambda(1-z_2)}{\mu}} - (p_2 + p_3)\right) \left(e^{\frac{\lambda(1-z_2)}{\mu}} - (p_1 + p_3)\right)} + \frac{p_1 p_2 p_3}{e^{\frac{\lambda(1-z_1)}{\mu}} - (p_2 + p_3)} \\
& \frac{\left(e^{\frac{\lambda(1-z_1)}{\mu}} - p_3\right)}{e^{\frac{\lambda(1-z_1)}{\mu}} - p_3}
\end{aligned} \tag{10}$$

4. Site Occupancy PGF in a G-ASIP: $n = 4$

In this section, we apply the matrix approach introduced and utilized in Section 3 to derive the PGF of a 4-site G-ASIP. A general arrival process occurs at all sites Q_1, Q_2, Q_3 and Q_4 during the time between two successive gate openings where $\hat{A}(z_1, z_2, z_3, z_4)$ is the PGF of the number of arrivals to the sites. All other model assumptions are similar to those in Section 3. $G(z_1, z_2, z_3, z_4) = E[z_1^{X_1} z_2^{X_2} z_3^{X_3} z_4^{X_4}]$, $|z_i| \leq 1$ is the PGF of the site occupancies.

In what follows, we derive an explicit solution for the PGF $G(z_1, z_2, z_3, z_4)$.

4.1. Derivation

The multidimensional law of motion yields

If gate 1 opens at step $r + 1$:

$$X_1^{(r+1)} = 0, \quad X_2^{(r+1)} = X_1^{(r)} + A_1^{(r+1)} + X_2^{(r)} + A_2^{(r+1)}, \quad X_3^{(r+1)} = X_3^{(r)} + A_3^{(r+1)}, \quad X_4^{(r+1)} = X_4^{(r)} + A_4^{(r+1)}$$

If gate 2 opens at step $r + 1$:

$$X_1^{(r+1)} = X_1^{(r)} + A_1^{(r+1)}, \quad X_2^{(r+1)} = 0, \quad X_3^{(r+1)} = X_3^{(r)} + A_3^{(r+1)} + X_2^{(r)} + A_2^{(r+1)}, \quad X_4^{(r+1)} = X_4^{(r)} + A_4^{(r+1)}$$

If gate 3 opens at step $r + 1$:

$$X_1^{(r+1)} = X_1^{(r)} + A_1^{(r+1)}, \quad X_2^{(r+1)} = X_2^{(r)} + A_2^{(r+1)}, \quad X_3^{(r+1)} = 0, \quad X_4^{(r+1)} = X_3^{(r)} + A_3^{(r+1)} + X_4^{(r)} + A_4^{(r+1)}$$

If gate 4 opens at step $r + 1$:

$$X_1^{(r+1)} = X_1^{(r)} + A_1^{(r+1)}, \quad X_2^{(r+1)} = X_2^{(r)} + A_2^{(r+1)}, \quad X_3^{(r+1)} = X_3^{(r)} + A_3^{(r+1)}, \quad X_4^{(r+1)} = 0$$

In steady state we have:

$$\begin{aligned}
G(z_1, z_2, z_3, z_4) &= E[z_1^{X_1} z_2^{X_2} z_3^{X_3} z_4^{X_4}] \\
&= p_1 E[z_1^0 z_2^{X_1+X_2+A_1+A_2} z_3^{X_3+A_3} z_4^{X_4+A_4}] + p_2 E[z_1^{X_1+A_1} z_2^0 z_3^{X_2+A_2+X_3+A_3} z_4^{X_4+A_4}] = \\
&\quad + p_3 E[z_1^{X_1+A_1} z_2^{X_2+A_2} z_3^0 z_4^{X_3+A_3+X_4+A_4}] + p_4 E[z_1^{X_1+A_1} z_2^{X_2+A_2} z_3^{X_3+A_3} z_4^0] \\
&= p_1 \hat{A}(z_2, z_2, z_3, z_4) G(z_2, z_2, z_3, z_4) + p_2 \hat{A}(z_1, z_3, z_3, z_4) G(z_1, z_3, z_3, z_4) \\
&\quad + p_3 \hat{A}(z_1, z_2, z_4, z_4) G(z_1, z_2, z_4, z_4) + p_4 \hat{A}(z_1, z_2, z_3) G(z_1, z_2, z_3)
\end{aligned}$$

That is,

$$\begin{aligned}
G(z_1, z_2, z_3, z_4) &= p_1 \hat{A}(z_2, z_2, z_3, z_4) G(z_2, z_2, z_3, z_4) + p_2 \hat{A}(z_1, z_3, z_3, z_4) G(z_1, z_3, z_3, z_4) \\
&\quad + p_3 \hat{A}(z_1, z_2, z_4, z_4) G(z_1, z_2, z_4, z_4) + p_4 \hat{A}(z_1, z_2, z_3) G(z_1, z_2, z_3)
\end{aligned} \quad (11)$$

As indicated, one can tediously apply a successive substitutions approach. Instead, we use the concise matrix representation approach developed in Section 3.2.3.

4.2. Direct Matrix Solution

Following the approach of Proposition 1 (for $n = 3$ system), when $n = 4$ the site occupancies PGF can be constructed from Equation (11) by solving a 70×70 linear system $\left(D(5, 4) = \binom{8}{4} = 70\right)$, or more efficiently, following Proposition 2, with a 35×35 linear system $\left(D(4, 4) = \binom{7}{3} = 35\right)$. The 5-steps substitutions are described in Appendix D and yield the following sets

$G(z)$ is derived immediately from Equation (11)

$$G(z) = \frac{p_1}{1 - (p_2 + p_3 + p_4) \hat{A}(z)} \quad (12)$$

$G(z_2, z_2)$ and $G(z_1, z_2)$ are obtained from

$$\begin{pmatrix} 1 - (p_3 + p_4) \hat{A}(z_1, z_2) & -p_1 \hat{A}(z_2, z_2) \\ 0 & 1 - (p_1 + p_3 + p_4) \hat{A}(z_2, z_2) \end{pmatrix} \begin{pmatrix} G(z_1, z_2) \\ G(z_2, z_2) \end{pmatrix} = \begin{pmatrix} p_2 \hat{A}(z_1) G(z_1) \\ p_2 \hat{A}(z_2) G(z_2) \end{pmatrix} \quad (13)$$

$G(z_1, z_1, z_2)$, $G(z_1, z_2, z_2)$ and $G(z_2, z_2, z_2)$ are calculated from

$$\begin{pmatrix} 1 - (p_1 + p_4) \hat{A}(z_1, z_1, z_2) & -p_2 \hat{A}(z_1, z_2, z_2) & 0 \\ 0 & 1 - (p_2 + p_4) \hat{A}(z_1, z_2, z_2) & -p_1 \hat{A}(z_2, z_2, z_2) \\ 0 & 0 & 1 - (p_1 + p_2 + p_4) \hat{A}(z_2, z_2, z_2) \end{pmatrix} \begin{pmatrix} G(z_1, z_1, z_2) \\ G(z_1, z_2, z_2) \\ G(z_2, z_2, z_2) \end{pmatrix} = \begin{pmatrix} p_3 \hat{A}(z_1, z_1) G(z_1, z_1) \\ p_3 \hat{A}(z_1, z_2) G(z_1, z_2) \\ p_3 \hat{A}(z_2, z_2) G(z_2, z_2) \end{pmatrix} \quad (14)$$

From the fourth step in Appendix D we obtain:

$$\begin{pmatrix} 1 - (p_1 + p_2) \hat{A}(z_1, z_1, z_1, z_2) & -p_3 \hat{A}(z_1, z_1, z_2, z_2) & 0 & 0 \\ 0 & 1 - (p_1 + p_3) \hat{A}(z_1, z_1, z_2, z_2) & -p_2 \hat{A}(z_1, z_2, z_2, z_2) & 0 \\ 0 & 0 & 1 - (p_2 + p_3) \hat{A}(z_1, z_2, z_2, z_2) & -p_1 \hat{A}(z_2, z_2, z_2, z_2) \\ 0 & 0 & 0 & 1 - (p_1 + p_2 + p_3) \hat{A}(z_2, z_2, z_2, z_2) \end{pmatrix} \begin{pmatrix} G(z_1, z_1, z_1, z_2) \\ G(z_1, z_1, z_2, z_2) \\ G(z_1, z_2, z_2, z_2) \\ G(z_2, z_2, z_2, z_2) \end{pmatrix} = \begin{pmatrix} p_4 \hat{A}(z_1, z_1, z_1) G(z_1, z_1, z_1) \\ p_4 \hat{A}(z_1, z_1, z_2) G(z_1, z_1, z_2) \\ p_4 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2) \\ p_4 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2) \end{pmatrix} \quad (15)$$

The last set gives

$$\begin{pmatrix} 1 - \hat{p}_1 A(z_1, z_1, z_2, z_3) & 0 & 0 \\ 0 & 1 - \hat{p}_2 A(z_1, z_2, z_2, z_3) & -\hat{p}_3 A(z_1, z_2, z_3, z_3) \\ 0 & 0 & 1 - \hat{p}_3 A(z_1, z_2, z_3, z_3) \end{pmatrix} \begin{pmatrix} G(z_1, z_1, z_2, z_3) \\ G(z_1, z_2, z_2, z_3) \\ G(z_1, z_2, z_3, z_3) \end{pmatrix} \\
= \begin{pmatrix} \hat{p}_2 A(z_1, z_2, z_2, z_2) G(z_1, z_2, z_2, z_2) + \hat{p}_3 A(z_1, z_1, z_3, z_3) G(z_1, z_1, z_3, z_3) + \hat{p}_4 A(z_1, z_1, z_2) G(z_1, z_1, z_2) \\ \hat{p}_1 A(z_2, z_2, z_2, z_3) G(z_2, z_2, z_2, z_3) + \hat{p}_4 A(z_1, z_2, z_2) G(z_1, z_2, z_2) \\ \hat{p}_1 A(z_2, z_2, z_3, z_3) G(z_2, z_2, z_3, z_3) + \hat{p}_2 A(z_1, z_3, z_3, z_3) G(z_1, z_3, z_3, z_3) + \hat{p}_4 A(z_1, z_2, z_3) G(z_1, z_2, z_3) \end{pmatrix}$$

By substituting the PGFs $G(z_2, z_2, z_3, z_4)$, $G(z_1, z_3, z_3, z_4)$, $G(z_1, z_2, z_4, z_4)$ and $G(z_1, z_2, z_3)$ in Equation (11), the required site occupancies PGF $G(z_1, z_2, z_3, z_4)$ is obtained.

5. Matrix Representation for an n -Site System

5.1. Computational Effort

Proposition 4. The G-ASIP model of n queues with arrivals at all sites can be solved by a linear system of size $D(n+1, n) = \binom{2n}{n}$.

Proof. The set $\{G(z_{i_1}, z_{i_2}, z_{i_3}, \dots, z_{i_n}) | 1 \leq i_1 \leq i_2 \leq \dots \leq i_n \leq n+1, z_{i_{n+1}} := 1\}$ contains all vertices and leaves representing a G-ASIP model with n sites. With appropriate substitutions of the equation derived from the laws of motion, one can create a set of $D(n+1, n) = \binom{2n}{n}$ linear equations with the same number of variables leading to a unique solution. This follows, since the number of variables in the set $\{G(z_{i_1}, z_{i_2}, z_{i_3}, \dots, z_{i_n}) | 1 \leq i_1 \leq i_2 \leq \dots \leq i_n \leq n+1, z_{i_{n+1}} := 1\}$ is equal to the amount of numbers one can create using the digits i_1, i_2, \dots, i_n in a non-decreasing order with repetition. \square

Proposition 5. The above G-ASIP model can be solved by a linear system of half the size, namely, with size $D(n, n) = \binom{2n-1}{n}$.

Proof. The set $\{G(z_{i_1}, z_{i_2}, z_{i_3}, \dots, z_{i_n}) | 1 \leq i_1 \leq i_2 \leq \dots \leq i_n \leq n\}$ contains all vertices and leaves representing a G-ASIP model with n sites except for variables that contain $z = 1$. Once the latter system is solved with its $D(n, n)$ variables, the digit 1 can be substituted as $z_{i_k} = 1$ to achieve all remaining needed vertices. Since $D(n, n) = \frac{1}{2}D(n+1, n)$, only half of the variables are required. \square

Let us denote the PGF $G_{i_1, i_2, \dots, i_k}^n \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right)$ where i_j denotes the number of z_j element repetitions in $G_{i_1, i_2, \dots, i_k}^n(\cdot)$, such that $\sum_{j=1}^k i_j = n$.

Lemma 1. For a given k there are $\binom{n-1}{k-1}$ PGFs variables

$$G_{i_1, i_2, \dots, i_k}^n \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right)$$

Proof. The number of PGFs $G_{i_1, i_2, \dots, i_k}^n(\cdot)$ where i_j denotes the number of z_j element repetitions such that $\sum_{j=1}^k i_j = n$, is equal to the number of n -digit numbers produced by the k digits $1, 2, \dots, k$ ($k \leq n$) in an increasing order, where all digits are used at least once. It is equal to the number of combinations to distribute $n - k$ identical elements in k cells, being equal to $D(k, n - k) = \binom{k-1+n-k}{k-1} = \binom{n-1}{k-1}$. \square

Lemma 2. There is a total of $2^{n-1} - 1$ PGFs distinct variables

$$G_{i_1, i_2, \dots, i_k}^n \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right).$$

Proof. According to Lemma 1, there are $\binom{n-1}{k-1}$ PGFs $G_{i_1, i_2, \dots, i_k}^n(\cdot)$ variables. Summing over all $k = 1, \dots, n-1$, yields, $\sum_{k=1}^{n-1} \binom{n-1}{k-1} = 2^{n-1} - 1$. \square

Proposition 6. The n -site G-ASIP can be solved by a set of $2^n - (n+1)$ linear equations, which can be decomposed into $n-1$ subsystems, each of $2^{k-1} - 1$ linear equations, $k = 1, 2, \dots, n$.

Proof. The set of variables needed to retrieve $G(z_1, z_2, \dots, z_n)$ is $\left\{ G_{i_1, i_2, \dots, i_k}^j \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq j, 1 \leq j \leq n \right\}$. This set can be decomposed into the disjoint union $\bigcup_{j=1}^n \left\{ G_{i_1, i_2, \dots, i_k}^j(\cdot) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq j, 1 \leq j \leq n \right\}$.

Using the disjoint property,

$$\left| \bigcup_{j=1}^n \left\{ G_{i_1, i_2, \dots, i_k}^j(\cdot) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq j, 1 \leq j \leq n \right\} \right| = \sum_{j=1}^n \left| \left\{ G_{i_1, i_2, \dots, i_k}^j(\cdot) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq j, 1 \leq j \leq n \right\} \right|$$

According to Lemma 2, the number of distinct variables in $\left\{ G_{i_1, i_2, \dots, i_k}^j \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right) \mid 1 \leq i_1 \leq i_2 \leq \dots \leq i_k \leq j, 1 \leq j \leq n \right\}$ is $2^{j-1} - 1$.

Hence, $\sum_{j=1}^n (2^{j-1} - 1) = (2^n - 1) - n = 2^n - (n+1)$. \square

5.2. The Effort Required to Move from a Solution of an n -Site Network to a Solution of an $(n + 1)$ -Site Network

In previous sections, we developed a procedure describing how to derive a solution for the site occupancies PGF in the G-ASIP networks with n sites. Specifically, we first showed that its solution requires a set of $\binom{2n}{n}$ equations, and then the cut of the number of equations in half to $\binom{2n-1}{n}$. It was further shown that the latter set can be reduced to a set of $2^n - (n + 1)$ equations.

For example, the site occupancies PGF of the $n = 3$ site G-ASIP could be obtained by a 20×20 system $\left(D(4, 3) = \binom{6}{3} = 20\right)$, according to Proposition 4. Then, according to Proposition 5, the system can be reduced and solved by a 10×10 system $\left(D(3, 3) = \binom{5}{3} = 10\right)$. Furthermore, the system was solved in Section 3.2.3 by as little as two subsystems of sizes 2×2 and 3×3 only.

Similarly, to obtain the site occupancies PGF for the $n = 4$ site G-ASIP, instead of solving a 70×70 system $\left(D(5, 4) = \binom{8}{4} = 70\right)$, according to Proposition 4, or a 35×35 system $\left(D(7, 4) = \binom{7}{4} = 35\right)$ according to Proposition 5, the PGF can be obtained with as few as 3 systems of order 2×2 , 3×3 , and 7×7 only, as shown in Section 4.2. Furthermore, the latter 7×7 system is decomposed into two subsystems of order 4×4 and 3×3 .

We now present a scheme showing how to expand the solution of the n -site G-ASIP into an $(n + 1)$ -site network.

Proposition 7. Given the sites occupancies PGF $G(z_1, z_2, \dots, z_n)$ of an n -site G-ASIP, one can expand the solution to an $(n + 1)$ -site network with a system of $2^n - 1$ linear equations.

Proof. Denoting by $G(z_1, z_2, \dots, z_{n+1})$ the sites occupancies PGF of the $(n + 1)$ -site network. As seen in Proposition 3, the set of variables needed to retrieve $G(z_1, z_2, \dots, z_{n+1})$ is $\{G(z_{i_1}, z_{i_2}, z_{i_3}, \dots, z_{i_{n+1}}) | 1 \leq i_1 \leq i_2 \leq \dots \leq i_{n+1} \leq n + 1\}$. With $G(z_1, z_2, \dots, z_n)$ given, one needs only the reduced set $\left\{G_{i_1, i_2, \dots, i_k}^{n+1} \left(\underbrace{z_1, \dots, z_1}_{i_1}, \underbrace{z_2, \dots, z_2}_{i_2}, \dots, \underbrace{z_k, \dots, z_k}_{i_k} \right) | 1 \leq i_1 \leq i_2 \leq \dots \leq i_{n+1} \leq n + 1 \right\}$. According to Lemma 2, there are only $2^n - 1$ such distinct variables. \square

The site occupancies PGF of the $n = 5$ site G-ASIP can be obtained by a 252×252 system $\left(D(6, 5) = \binom{10}{5} = 252\right)$ according to Proposition 4. However, given $G(z_1, z_2, z_3, z_4)$ the site occupancies PGF of the 4-site G-ASIP, one can expand it to the $n = 5$ sites with as little as $2^4 - 1 = 15$ additional equations, according to Proposition 7.

6. Mean Site Occupancies

6.1. Mean Site Occupancies in a G-ASIP with $n = 3$ Sites

Focusing on mean site occupancies, one can take a direct approach while using the occupancies laws of motion from Section 3.1:

If gate 1 opens at step $r + 1$:

$$X_1^{(r+1)} = 0, \quad X_2^{(r+1)} = X_1^{(r)} + A_1^{(r+1)} + X_2^{(r)} + A_2^{(r+1)}, \quad X_3^{(r+1)} = X_3^{(r)} + A_3^{(r+1)}$$

If gate 2 opens at step $r + 1$:

$$X_1^{(r+1)} = X_1^{(r)} + A_1^{(r+1)}, \quad X_2^{(r+1)} = 0, \quad X_3^{(r+1)} = X_2^{(r)} + A_2^{(r+1)} + X_3^{(r)} + A_3^{(r+1)}$$

If gate 3 opens at step $r + 1$:

$$X_1^{(r+1)} = X_1^{(r)} + A_1^{(r+1)}, \quad X_2^{(r+1)} = X_2^{(r)} + A_2^{(r+1)}, \quad X_3^{(r+1)} = 0$$

Thus, the mean number of particles in site Q_1 is given by:

$$E[X_1] = p_2(E[X_1] + E[A_1]) + p_3(E[X_1] + E[A_1]) = (p_2 + p_3)E[X_1] + (p_2 + p_3)E[A_1]$$

Simplification yields,

$$p_1 E[X_1] = (1 - p_1)E[A_1]$$

Thus,

$$E[X_1] = \frac{1}{p_1} E[A_1] - E[A_1]$$

Next,

$$\begin{aligned} E[X_2] &= p_1(E[X_1] + E[A_1] + E[X_2] + E[A_2]) + p_3(E[X_2] + E[A_2]) \\ &= p_1(E[X_1] + E[A_1]) + (p_1 + p_3)E[X_2] + (p_1 + p_3)E[A_2], \end{aligned} \quad (16)$$

which leads to,

$$p_2 E[X_2] = p_1(E[X_1] + E[A_1]) + (p_1 + p_3)E[A_2]$$

Substituting (17) in (18) yields,

$$p_2 E[X_2] = p_1 \left(\frac{(1 - p_1)}{p_1} E[A_1] + E[A_1] \right) + (p_1 + p_3)E[A_2] = E[A_1] + (p_1 + p_3)E[A_2]$$

resulting in

$$E[X_2] = \frac{1}{p_2} \sum_{j=1}^2 E[A_j] - E[A_2] \quad (17)$$

Similarly, considering site Q_3 :

$$\begin{aligned} E[X_3] &= p_1(E[X_3] + E[A_3]) + p_2(E[X_2] + E[A_2] + E[X_3] + E[A_3]) \\ &= (p_1 + p_2)E[X_3] + (p_1 + p_2)E[A_3] + p_2(E[X_2] + E[A_2]) \end{aligned}$$

$$p_3 E[X_3] = (p_1 + p_2)E[A_3] + p_2(E[X_2] + E[A_2]) \quad (18)$$

Substituting (17) in (18) yields

$$p_3 E[X_3] = (p_1 + p_2)E[A_3] + p_2 \left(\frac{1}{p_2} (E[A_1] + (1 - p_2)E[A_2]) + E[A_2] \right) = (p_1 + p_2)E[A_3] + (E[A_1] + E[A_2])$$

Therefore,

$$E[X_3] = \frac{1}{p_3} (E[A_1] + E[A_2] + (1 - p_3)E[A_3]) = \frac{1}{p_3} \sum_{j=1}^3 E[A_j] - E[A_3]$$

6.2. Mean Site Occupancies in an n -Site G-ASIP

Expanding the solution of Section 6.1 to the non-homogeneous n -site G-ASIP we get

$$E[X_1] = \frac{1}{p_1} E[A_1] - E[A_1] \quad (19)$$

$$E[X_k] = \frac{1}{p_k} \sum_{j=1}^k E[A_j] - E[A_k], \quad k = 1, \dots, n \quad (20)$$

The mean site occupancies of the n -site classical ASIP was analyzed in [28], where arrivals occur only to Q_1 with Poisson rate λ , and each gate k opens independently of other gates following an Exponential distribution with mean $\frac{1}{\mu_k}$. It was shown that under stationary conditions $E[X_k] = \frac{\lambda}{\mu_k}$, $k = 1, \dots, n$. Our general model can be compared to the above model by substituting $p_k = \frac{\mu_k}{\mu}$, $E[A_1] = \frac{\lambda}{\mu}$ while for $k = 2, \dots, n$, $E[A_k] = 0$, where $\mu = \sum_{k=1}^n \mu_k$. Equations (19) and (20) become, respectively,

$$E[X_1] = \frac{\lambda}{\mu_1} \left(1 - \frac{\mu_1}{\mu}\right) \quad (21)$$

$$E[X_k] = \frac{\lambda}{\mu_k}, \quad k = 2, \dots, n \quad (22)$$

Note that X_k is defined to be the number of particles in site Q_k right after a gate opening instant. As such, the mean $E[X_k]$ is also calculated right after gate openings. An important observation is that, when arrivals occur only to Q_1 , the number of particles present in each site Q_k , $k = 2, \dots, n$ stays constant between instants of gate openings. Thus, taking expectation right after a gate opening or any other time between gate openings yields the same result. Hence, (22) coincides with Equation (16) in [28]. However, in site Q_1 , since Poisson arrivals occur continuously in between gate openings, the mean $E[X_1]$ is affected by the sampling instant. While the mean site occupancies in [28] was taken uniformly over time, in the general G-ASIP the mean is taken right after gate openings, which leads us to expect a smaller term for this mean, as obtained in (21).

Consider now the case where the arriving particles are spread uniformly between the n sites such that $E[A_k] = \frac{\lambda}{n\mu}$, for all k . Gate opening probabilities are as before, $p_k = \frac{\mu_k}{\mu}$. Then, (20) becomes,

$$E[X_k] = \frac{1}{p_k} \sum_{j=1}^k E[A_j] - E[A_k] = \frac{\mu}{\mu_k} \sum_{j=1}^k \frac{\lambda}{\mu n} - \frac{\lambda}{\mu n} = \frac{\mu}{\mu_k} \frac{\lambda k}{\mu n} - \frac{\lambda}{\mu n} = \frac{\lambda k}{\mu_k n} - \frac{\lambda}{\mu n} = \frac{\lambda}{n\mu_k} \left(k - \frac{\mu_k}{\mu}\right)$$

for all $k = 1, \dots, n$.

If $\mu_k = \mu_1$, ($\mu = n\mu_1$) for all k , then

$$E[X_k] = \frac{\lambda}{n\mu_1} \left(k - \frac{\mu_1}{\mu}\right) = \frac{\lambda}{\mu} \left(k - \frac{1}{n}\right), \quad k = 1, \dots, n$$

Notice that the mean difference in occupancy between two consecutive sites, right after gate opening, yields,

$$E[X_{k+1}] - E[X_k] = \frac{\lambda}{\mu} \left(k + 1 - \frac{1}{n}\right) - \frac{\lambda}{\mu} \left(k - \frac{1}{n}\right) = \frac{\lambda}{\mu} \quad (23)$$

That is, the mean difference is constant and equals $\frac{\lambda}{\mu}$, which is the mean total number of arrivals to the entire system between two gate openings.

The mean load of the first k sites can now be calculated as follows:

Denote by $X_{(k)} = \sum_{j=1}^k X_j$ the total load of the first k sites, i.e., the total number of particles in the first k sites right after gate openings. Equation (24) becomes,

$$E[X_{(k)}] = E\left[\sum_{m=1}^k X_m\right] = \sum_{m=1}^k E[X_m] = \sum_{m=1}^k \left(\frac{1}{p_m} \sum_{j=1}^m E[A_j] - E[A_m]\right) \quad (24)$$

$$E[X_{(k)}] = E\left[\sum_{m=1}^k X_m\right] = \sum_{m=1}^k E[X_m] = \sum_{m=1}^k \left(\frac{1}{p_m} \sum_{j=1}^m E[A_j] - E[A_m]\right) \quad (25)$$

Substituting in Equation (24), $p_k = \frac{\mu_k}{\mu}$, $E[A_1] = \frac{\lambda}{\mu}$ and $E[A_k] = 0, k = 2, \dots, n$, where $\mu = \sum_{k=1}^n \mu_k$, leads to

$$E[X_{(k)}] = \sum_{m=1}^k \left(\frac{1}{p_m} E[A_1] - E[A_m]\right) = \sum_{m=1}^k \left(\frac{1}{p_m} E[A_1]\right) - E[A_1] = \sum_{m=1}^k \left(\frac{\mu}{\mu_k} \frac{\lambda}{\mu}\right) - \frac{\lambda}{\mu} = \sum_{m=1}^k \frac{\lambda}{\mu_k} - \frac{\lambda}{\mu}$$

whereas the result in [28] is $E[X_{(k)}] = \sum_{m=1}^k \frac{\lambda}{\mu_k}$. The difference of $\frac{\lambda}{\mu}$ between the two results is due to the same reason explained above, which is the instant in which the system is observed.

Since $E[X_{k+1}] - E[X_k] = \frac{\lambda}{\mu}$, the load of the first k sites is

$$E[X_{(k)}] = \sum_{j=1}^k E[X_j] = \sum_{j=1}^k \left(E[X_1] + \frac{j\lambda}{\mu}\right) = kE[X_1] + \frac{\lambda k(k-1)}{2\mu} = \frac{\lambda k}{\mu} \left(1 - \frac{1}{n}\right) + \frac{\lambda k(k-1)}{2\mu} = \frac{\lambda k}{\mu} \left(\frac{k+1}{2} - \frac{1}{n}\right)$$

while the load of the entire system is

$$E[X_{(n)}] = \frac{\lambda(n+2)(n-1)}{2\mu}$$

7. Conclusions

An innovative matrix approach to derive the multidimensional probability-generating function (PGF) of the site occupancies in a generalized n -site ASIP network (G-ASIP) with arrivals to all sites is developed. The family $\Gamma(\alpha, \alpha\mu)$ of Gamma distributed inter-gate opening instants is analyzed and its extreme cases, Exponential ($\alpha = 1$), and Deterministic ($\alpha \rightarrow \infty$), are further investigated. It is then shown how the matrix approach considerably reduces the required computational effort to obtain the occupancy PGF. Explicit results for the cases $n = 3$ and $n = 4$ sites are derived. Furthermore, a procedure to move from a PGF of an n -site network to the PGF of an $(n+1)$ -site network is constructed. It is shown that the probability that site k is occupied is a function of both the site's index and the arrival flux and not solely of the site's index, which leads to refined formulae for the probability that site k is occupied and for its conditional mean site's occupancy. Consequently, it is shown that, in the case where the arrival process to the first site is Poisson with rate λ , the following interesting property holds: $P(\text{site } k \text{ is occupied} \mid \lambda = 1) = P(\text{site } k+1 \text{ is occupied} \mid \lambda \rightarrow \infty)$. The results are enhanced by numerical calculations exhibited in graphs.

Author Contributions: Conceptualization, U.Y.; formal analysis, Y.Y. and U.Y.; investigation, Y.Y. and U.Y.; writing—original draft, Y.Y.; writing—review and editing, Y.Y. and U.Y. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

TSS	Tandem Stochastic System
TJN	Tandem Jackson Network
ASEP	Asymmetric Simple Exclusion Process
ASIP	Asymmetric Simple Inclusion Process
G-ASIP	Generalized Asymmetric Simple Inclusion Process
PGF	Probability-generating Function

Appendix A. Occupancy Probabilities of the First Five Sites

Direct calculation from Theorem 1 yields the following occupancy probabilities of the first five sites:

$$\begin{aligned}
 Q_\lambda(1) &= \frac{n + \lambda}{n(\lambda + 1)} \\
 Q_\lambda(2) &= \frac{1}{2} \left(1 + \frac{n^2}{(n\lambda + n)^2} \right) = \frac{1}{2} \left(1 + \frac{\lambda + 1}{(\lambda + 1)^3} \right) \\
 Q_\lambda(3) &= \frac{1}{8} \left(2 \quad 1 \quad \frac{n}{(\lambda+n)(\lambda+1)} \right) \begin{pmatrix} 2 & 0 & 0 \\ 0 & 1 & \frac{n}{(\lambda+n)(\lambda+1)} \\ 0 & 0 & \frac{2}{\lambda+1} \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ \frac{n+\lambda}{n(\lambda+1)} \end{pmatrix} = \frac{1}{8} \left(5 + \frac{\lambda + 3}{(\lambda + 1)^3} \right) \\
 Q_\lambda(4) &= \frac{1}{64} \left(4 \quad 2 \quad 1 \quad \frac{n}{(\lambda+n)(\lambda+1)} \right) \begin{pmatrix} 4 & 0 & 0 & 0 \\ 0 & 2 & 1 & \frac{n}{(\lambda+n)(\lambda+1)} \\ 0 & 0 & 2 & \frac{2n}{(\lambda+n)(\lambda+1)} \\ 0 & 0 & 0 & \frac{4}{\lambda+1} \end{pmatrix} \begin{pmatrix} 2 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & \frac{n}{(\lambda+n)(\lambda+1)} \\ 0 & 0 & 0 & \frac{2}{\lambda+1} \end{pmatrix} \begin{pmatrix} 1 \\ 1 \\ 1 \\ \frac{n+\lambda}{n(\lambda+1)} \end{pmatrix} \\
 &= \frac{1}{16} \left(11 + \frac{\lambda^2 + 4\lambda + 5}{(\lambda + 1)^4} \right) = \frac{1}{16} \left(11 + \frac{\lambda + 3}{(\lambda + 1)^3} + \frac{2}{(\lambda + 1)^4} \right) \\
 Q_\lambda(5) &= \frac{1}{128} \left(93 + \frac{5\lambda^3 + 25\lambda^2 + 47\lambda + 35}{(\lambda + 1)^5} \right) = \frac{1}{128} \left(93 + \frac{2(\lambda + 5)}{(\lambda + 1)^5} + \frac{10(\lambda + 2)}{(\lambda + 1)^4} + \frac{5}{(\lambda + 1)^2} \right)
 \end{aligned}$$

Appendix B. Site Occupancy Probabilities of a 3-Site G-ASIP

With proper substitutions in the matrix system in Section 3.2.3 we get

$$G(0) = P(I_1 = 0) = \frac{p_1}{1 - (p_2 + p_3)\hat{A}(0)} \quad (\text{A1})$$

$$\begin{pmatrix} 1 - p_3\hat{A}(1,0) & -p_1\hat{A}(0,0) \\ 0 & 1 - (p_1 + p_3)\hat{A}(0,0) \end{pmatrix} \begin{pmatrix} G(1,0) \\ G(0,0) \end{pmatrix} = \begin{pmatrix} p_2 \\ p_2\hat{A}(0)G(0) \end{pmatrix} \quad (\text{A2})$$

and

$$\begin{pmatrix} 1 - p_1\hat{A}(1,1,0) & -p_2\hat{A}(1,0,0) & 0 \\ 0 & 1 - p_2\hat{A}(1,0,0) & -p_1\hat{A}(0,0,0) \\ 0 & 0 & 1 - (p_1 + p_2)\hat{A}(0,0,0) \end{pmatrix} \begin{pmatrix} G(1,1,0) \\ G(1,0,0) \\ G(0,0,0) \end{pmatrix} = \begin{pmatrix} p_3 \\ p_3\hat{A}(1,0)G(1,0) \\ p_3\hat{A}(0,0)G(0,0) \end{pmatrix} \quad (\text{A3})$$

Solving the set (A2) gives

$$\begin{pmatrix} G(0,0) \\ G(1,0) \end{pmatrix} = \frac{1}{\left(1 - (p_1 + p_3)\hat{A}(0,0)\right)\left(1 - p_3\hat{A}(1,0)\right)} \begin{pmatrix} 1 - p_3\hat{A}(1,0) & 0 \\ p_1\hat{A}(0,0) & 1 - (p_1 + p_3)\hat{A}(0,0) \end{pmatrix} \begin{pmatrix} p_2\hat{A}(0)G(0) \\ p_2 \end{pmatrix} \quad (\text{A4})$$

Substituting (A1) in (A4) gives

$$G(0,0) = \frac{p_1 p_2 \hat{A}(0)}{\left(1 - (p_2 + p_3)\hat{A}(0)\right)\left(1 - (p_1 + p_3)\hat{A}(0,0)\right)}$$

$$G(1,0) = P(I_2 = 0) = \frac{p_1^2 p_2 \hat{A}(0) \hat{A}(0,0)}{\left(1 - (p_1 + p_3)\hat{A}(0,0)\right)\left(1 - (p_2 + p_3)\hat{A}(0)\right)\left(1 - p_3\hat{A}(1,0)\right)} + \frac{p_2}{1 - p_3\hat{A}(1,0)}$$

Solving the set (A3) leads to

$$\begin{pmatrix} G(1,1,0) \\ G(1,0,0) \\ G(0,0,0) \end{pmatrix} = \begin{pmatrix} \frac{1}{1 - p_1\hat{A}(1,1,0)} & \frac{p_2\hat{A}(1,0,0)}{\left(1 - p_1\hat{A}(1,1,0)\right)\left(1 - p_2\hat{A}(1,0,0)\right)} & \frac{p_1 p_2 \hat{A}(0,0,0) \hat{A}(1,0,0)}{\left(1 - p_1\hat{A}(1,1,0)\right)\left(1 - p_2\hat{A}(1,0,0)\right)\left(1 - (p_1 + p_2)\hat{A}(0,0,0)\right)} \\ 0 & \frac{1}{1 - p_2\hat{A}(1,0,0)} & \frac{p_1\hat{A}(0,0,0)}{\left(1 - p_2\hat{A}(1,0,0)\right)\left(1 - (p_1 + p_2)\hat{A}(0,0,0)\right)} \\ 0 & 0 & \frac{1}{1 - (p_1 + p_2)\hat{A}(0,0,0)} \end{pmatrix} \begin{pmatrix} p_3 \\ p_3\hat{A}(1,0)G(1,0) \\ p_3\hat{A}(0,0)G(0,0) \end{pmatrix}$$

which gives

$$G(1,1,0) = P(I_3 = 0) = \frac{p_3}{1 - p_1\hat{A}(1,1,0)} + \frac{p_2\hat{A}(1,0,0)p_3\hat{A}(1,0)G(1,0)}{\left(1 - p_1\hat{A}(1,1,0)\right)\left(1 - p_2\hat{A}(1,0,0)\right)} + \frac{p_1 p_2 p_3 \hat{A}(0,0) \hat{A}(0,0,0) \hat{A}(1,0,0) G(0,0)}{\left(1 - p_1\hat{A}(1,1,0)\right)\left(1 - p_2\hat{A}(1,0,0)\right)\left(1 - (p_1 + p_2)\hat{A}(0,0,0)\right)}.$$

Appendix C. Site Occupancy Probabilities of a 4-Site G-ASIP

The following 4 substitutions in the sets presented in Section 3.2 yield the site occupancy probabilities of a G-ASIP where arrivals occur to all sites.

Substituting $z = 0$ in Equation (7) gives

$$G(0) = P(I_1 = 0) = \frac{p_1}{1 - (p_2 + p_3 + p_4)\hat{A}(0)}$$

Substituting $z_1 = 1$ and $z_2 = 0$ in Equation (8) yields

$$\begin{pmatrix} 1 - (p_3 + p_4)\hat{A}(1,0) & -p_1\hat{A}(0,0) \\ 0 & 1 - (p_1 + p_3 + p_4)\hat{A}(0,0) \end{pmatrix} \begin{pmatrix} G(1,0) \\ G(0,0) \end{pmatrix} = \begin{pmatrix} p_2 \\ p_2\hat{A}(0)G(0) \end{pmatrix}$$

Substituting $z_1 = 1$ and $z_2 = 0$ in Equation (9) leads to

$$\begin{pmatrix} 1 - (p_1 + p_4)\hat{A}(1,1,0) & -p_2\hat{A}(1,0,0) & 0 \\ 0 & 1 - (p_2 + p_4)\hat{A}(1,0,0) & -p_1\hat{A}(0,0,0) \\ 0 & 0 & 1 - (p_1 + p_2 + p_4)\hat{A}(0,0,0) \end{pmatrix} \begin{pmatrix} G(1,1,0) \\ G(1,0,0) \\ G(0,0,0) \end{pmatrix} = \begin{pmatrix} p_3 \\ p_3\hat{A}(1,0)G(1,0) \\ p_3\hat{A}(0,0)G(0,0) \end{pmatrix}$$

Substituting $z_1 = 1$ and $z_2 = 0$ in Equation (10) gives

$$\begin{pmatrix} 1 - (p_1 + p_2)\hat{A}(1,1,1,0) & -p_3\hat{A}(1,1,0,0) & 0 & 0 \\ 0 & 1 - (p_1 + p_3)\hat{A}(1,1,0,0) & -p_2\hat{A}(1,0,0,0) & 0 \\ 0 & 0 & 1 - (p_2 + p_3)\hat{A}(1,0,0,0) & -p_1\hat{A}(0,0,0,0) \\ 0 & 0 & 0 & 1 - (p_1 + p_2 + p_3)\hat{A}(0,0,0,0) \end{pmatrix} \begin{pmatrix} G(1,1,1,0) \\ G(1,1,0,0) \\ G(1,0,0,0) \\ G(0,0,0,0) \end{pmatrix} = \begin{pmatrix} p_4 \\ p_4\hat{A}(1,1,0)G(1,1,0) \\ p_4\hat{A}(1,0,0)G(1,0,0) \\ p_4\hat{A}(0,0,0)G(0,0,0) \end{pmatrix}$$

Solving the sets above leads to the explicit expressions of the site occupancy probabilities in a 4-site G-ASIP.

Appendix D. Derivation of $G(z_1, z_2, z_3)$

The successive substitution procedure to calculate $G(z_1, z_2, z_3)$ is detailed below.

The procedure is to iterate Equation (6) repeatedly, in a branching tree structure, until reaching the leaves $G(1, 1, 1) = 1$. Then, the tree is folded back, yielding the value of the root $G(z_1, z_2, z_3)$. See also [28].

Figure A1 illustrates the iterative solution when $n = 3$, following Equation (6).




Figure A1. Tree representation of the solution procedure for $G(z_1, z_2, z_3)$.

Since there are various leaves that repeat themselves in Figure A1, one can shrink Figure A1 into Figure A2 below:




Figure A2. A reduced representation of Figure A1.

Starting with the term $G(z_1, z_2, 1)$ and substituting $z_3 = 1$ in Equation (6) results in

$$G(z_1, z_2, 1) = p_1 \hat{A}(z_2, z_2, 1)G(z_2, z_2, 1) + p_2 \hat{A}(z_1, 1, 1)G(z_1, 1, 1) + p_3 \hat{A}(z_1, z_2, 1)G(z_1, z_2, 1) \quad (\text{A5})$$

Hence,

$$G(z_1, z_2, 1) = \frac{p_1 \hat{A}(z_2, z_2, 1)G(z_2, z_2, 1) + p_2 \hat{A}(z_1, 1, 1)G(z_1, 1, 1)}{1 - p_3 \hat{A}(z_1, z_2, 1)} \quad (\text{A6})$$

Substituting $z_2 = 1$ in Equation (A6) leads to

$$G(z_1, 1, 1) = \frac{p_1}{1 - (p_2 + p_3) \hat{A}(z_1, 1, 1)} \quad (\text{A7})$$

Substituting (A7) in (A6) yields

$$G(z_1, z_2, 1) = \frac{p_1 \hat{A}(z_2, z_2, 1)G(z_2, z_2, 1) + p_2 \hat{A}(z_1, 1, 1) \frac{p_1}{1 - (p_2 + p_3) \hat{A}(z_1, 1, 1)}}{1 - p_3 \hat{A}(z_1, z_2, 1)} \quad (\text{A8})$$

To calculate $G(z_1, z_2, 1)$, we substitute $z_1 = z_2$ in (A8) and obtain

$$G(z_2, z_2, 1) = \frac{p_1 p_2 \hat{A}(z_2, 1, 1)}{\left(1 - (p_1 + p_3) \hat{A}(z_2, z_2, 1)\right) \left(1 - (p_2 + p_3) \hat{A}(z_2, 1, 1)\right)} \quad (\text{A9})$$

Substituting (A9) in (A8) results in

$$G(z_1, z_2, 1) = \frac{\frac{p_1^2 p_2 \hat{A}(z_2, 1, 1) \hat{A}(z_2, z_2, 1)}{\left(1 - (p_1 + p_3) \hat{A}(z_2, z_2, 1)\right) \left(1 - (p_2 + p_3) \hat{A}(z_2, 1, 1)\right)} + \frac{p_1 p_2 \hat{A}(z_1, 1, 1)}{1 - (p_2 + p_3) \hat{A}(z_1, 1, 1)}}{1 - p_3 \hat{A}(z_1, z_2, 1)} \quad (\text{A10})$$

We move on to calculate $G(z_1, z_3, z_3)$

To simplify notation, set $\hat{A}(z_i, z_j, 1) \equiv \hat{A}(z_i, z_j)$ and $\hat{A}(z_i, 1, 1) \equiv \hat{A}(z_i)$. Similarly, set $G(z_i, z_j, 1) \equiv G(z_i, z_j)$ and $G(z_i, 1, 1) \equiv G(z_i)$.

Substituting $z_2 = z_3$ in (6) gives

$$G(z_1, z_3, z_3) = p_1 \hat{A}(z_3, z_3, z_3)G(z_3, z_3, z_3) + p_2 \hat{A}(z_1, z_3, z_3)G(z_1, z_3, z_3) + p_3 \hat{A}(z_1, z_3)G(z_1, z_3)$$

Hence,

$$G(z_1, z_3, z_3) = \frac{p_1 \hat{A}(z_3, z_3, z_3)G(z_3, z_3, z_3) + p_3 \hat{A}(z_1, z_3)G(z_1, z_3)}{1 - p_2 \hat{A}(z_1, z_3, z_3)} \quad (\text{A11})$$

Substituting $z_2 = z_3$ in (A10) leads to

$$G(z_1, z_3) = \frac{\frac{p_1^2 p_2 \hat{A}(z_3) \hat{A}(z_3, z_3)}{\left(1 - (p_1 + p_3) \hat{A}(z_3, z_3)\right) \left(1 - (p_2 + p_3) \hat{A}(z_3)\right)} + \frac{p_1 p_2 \hat{A}(z_1)}{1 - (p_2 + p_3) \hat{A}(z_1)}}{1 - p_3 \hat{A}(z_1, z_3)} \quad (\text{A12})$$

Now, to calculate $G(z_3, z_3, z_3)$, we substitute $z_1 = z_3$ in (A11) to get

$$G(z_3, z_3, z_3) = \frac{p_3 \hat{A}(z_3, z_3) G(z_3, z_3)}{1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3)} \quad (\text{A13})$$

Substituting $z_1 = z_3$ in (A12) yields

$$G(z_3, z_3) = \frac{\frac{p_1^2 p_2 \hat{A}(z_3) \hat{A}(z_3, z_3)}{1 - (p_1 + p_3) \hat{A}(z_3, z_3)} + p_1 p_2 \hat{A}(z_3)}{\left(1 - p_3 \hat{A}(z_3, z_3)\right) \left(1 - (p_2 + p_3) \hat{A}(z_3)\right)}$$

Hence,

$$G(z_3, z_3) = \frac{p_1 p_2 \hat{A}(z_3)}{\left(1 - (p_2 + p_3) \hat{A}(z_3)\right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3)\right)} \quad (\text{A14})$$

Substituting (A14) in (A13) yields

$$G(z_3, z_3, z_3) = \frac{p_1 p_2 p_3 \hat{A}(z_3, z_3) \hat{A}(z_3)}{\left(1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3)\right) \left(\left(1 - (p_2 + p_3) \hat{A}(z_3)\right)\right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3)\right)} \quad (\text{A15})$$

Finally, substituting (A15) and (A12) in (A11) leads to

$$G(z_1, z_3, z_3) = \frac{\frac{p_1^2 p_2 p_3 \hat{A}(z_3, z_3, z_3) \hat{A}(z_3, z_3) \hat{A}(z_3)}{\left(1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3)\right) \left(\left(1 - (p_2 + p_3) \hat{A}(z_3)\right)\right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3)\right)} + \frac{\frac{p_1^2 p_2 p_3 \hat{A}(z_1, z_3) \hat{A}(z_3) \hat{A}(z_3, z_3)}{\left(1 - (p_1 + p_3) \hat{A}(z_3, z_3)\right) \left(1 - (p_2 + p_3) \hat{A}(z_3)\right)} + \frac{p_1 p_2 p_3 \hat{A}(z_1, z_3) \hat{A}(z_1)}{1 - (p_2 + p_3) \hat{A}(z_1)}}{1 - p_2 \hat{A}(z_1, z_3, z_3)} \quad (\text{A16})$$

(c) The last step is to get $G(z_2, z_2, z_3)$

Substituting $z_1 = z_2$ in (A5) results in

$$G(z_2, z_2, z_3) = p_1 \hat{A}(z_2, z_2, z_3) G(z_2, z_2, z_3) + p_2 \hat{A}(z_2, z_3, z_3) G(z_2, z_3, z_3) + p_3 \hat{A}(z_2, z_2) G(z_2, z_2)$$

Hence,

$$G(z_2, z_2, z_3) = \frac{p_2 \hat{A}(z_2, z_3, z_3) G(z_2, z_3, z_3) + p_3 \hat{A}(z_2, z_2) G(z_2, z_2)}{1 - p_1 \hat{A}(z_2, z_2, z_3)} \quad (\text{A17})$$

$G(z_2, z_2)$ was already calculated in (A9), so to find $G(z_2, z_2, z_3)$ we substitute $z_1 = z_2$ in Equation (A16) and get

$$\begin{aligned}
 G(z_2, z_2, z_3) = & \frac{\hat{A}(z_2, z_3, z_3) \left(\frac{p_1^2 p_2^2 p_3 \hat{A}(z_3, z_3, z_3) \hat{A}(z_3, z_3) \hat{A}(z_3)}{\left((1 - (p_1 + p_2) \hat{A}(z_3, z_3, z_3)) \left((1 - (p_2 + p_3) \hat{A}(z_3)) \right) \left(1 - (p_1 + p_3) \hat{A}(z_3, z_3) \right) \right)} + \frac{p_1^2 p_2^2 p_3 \hat{A}(z_2, z_3) \hat{A}(z_3) \hat{A}(z_3)}{\left((1 - (p_1 + p_3) \hat{A}(z_3, z_3)) \left(1 - (p_2 + p_3) \hat{A}(z_3) \right) \right)} + \frac{p_1 p_2^2 p_3 \hat{A}(z_2, z_3) \hat{A}(z_2)}{1 - (p_2 + p_3) \hat{A}(z_2)} \right)}{1 - p_2 \hat{A}(z_2, z_3, z_3)} \\
 & + \frac{1 - p_1 \hat{A}(z_2, z_2, z_3)}{1 - p_1 \hat{A}(z_2, z_2, z_3)} + \frac{\frac{p_1 p_2 p_3 \hat{A}(z_2, z_2) \hat{A}(z_2)}{\left((1 - (p_1 + p_3) \hat{A}(z_2, z_2)) \left(1 - (p_2 + p_3) \hat{A}(z_2) \right) \right)}}{1 - p_1 \hat{A}(z_2, z_2, z_3)}
 \end{aligned} \quad (A18)$$

Thus, by substituting (A18), (A16) and (A10) in (6), the PGF $G(z_1, z_2, z_3)$ is obtained.

Appendix E. 4-Site G-ASIP Substitutions

The substitutions that are required in order to calculate the sites occupancy PGF are as follows

Step 1. Substituting $z_2 = 1, z_3 = 1, z_4 = 1$ in Equation (11) leads to

$$G(z_1) = p_1 + p_2 \hat{A}(z_1) G(z_1) + p_3 \hat{A}(z_1) G(z_1) + p_4 \hat{A}(z_1) G(z_1)$$

Step 2. Substituting $z_2 = 1, z_3 = 1, z_4 = 1$ in Equation (11) results in

$$G(z_1, z_2) = p_1 \hat{A}(z_2, z_2) G(z_2, z_2) + p_2 \hat{A}(z_1) G(z_1) + p_3 \hat{A}(z_1, z_2) G(z_1, z_2) + p_4 \hat{A}(z_1, z_2) G(z_1, z_2)$$

By substituting $z_1 = z_2, z_3 = 1, z_4 = 1$ in Equation (11) we get

$$G(z_2, z_2) = p_1 \hat{A}(z_2, z_2) G(z_2, z_2) + p_2 \hat{A}(z_2) G(z_2) + p_3 \hat{A}(z_2, z_2) G(z_2, z_2) + p_4 \hat{A}(z_2, z_2) G(z_2, z_2)$$

Step 3. Substituting $z_2 = z_1, z_3 = z_2, z_4 = 1$ in Equation (11) leads to

$$\begin{aligned}
 G(z_1, z_1, z_2) = & p_1 \hat{A}(z_1, z_1, z_2) G(z_1, z_1, z_2) + p_2 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2) \\
 & + p_3 \hat{A}(z_1, z_1) G(z_1, z_1) + p_4 \hat{A}(z_1, z_1, z_2) G(z_1, z_1, z_2)
 \end{aligned}$$

Substituting $z_3 = z_2, z_4 = 1$ in Equation (11) results in

$$\begin{aligned}
 G(z_1, z_2, z_2) = & p_1 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2) + p_2 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2) \\
 & + p_3 \hat{A}(z_1, z_2) G(z_1, z_2) + p_4 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2)
 \end{aligned}$$

By substituting $z_1 = z_2, z_3 = z_2, z_4 = 1$ in Equation (11) we get

$$\begin{aligned}
 G(z_2, z_2, z_2) = & p_1 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2) + p_2 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2) \\
 & + p_3 \hat{A}(z_2, z_2) G(z_2, z_2) + p_4 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2)
 \end{aligned}$$

Step 4. Substituting $z_2 = z_1, z_3 = z_1, z_4 = z_2$ in Equation (11) leads to

$$\begin{aligned}
 G(z_1, z_1, z_1, z_2) = & p_1 \hat{A}(z_1, z_1, z_1, z_2) G(z_1, z_1, z_1, z_2) + p_2 \hat{A}(z_1, z_1, z_1, z_2) G(z_1, z_1, z_1, z_2) \\
 & + p_3 \hat{A}(z_1, z_1, z_2, z_2) G(z_1, z_1, z_2, z_2) + p_4 \hat{A}(z_1, z_1, z_1) G(z_1, z_1, z_1)
 \end{aligned}$$

Substituting $z_2 = z_1, z_3 = z_2, z_4 = z_2$ in Equation (11) results in

$$G(z_1, z_1, z_2, z_2) = p_1 \hat{A}(z_1, z_1, z_2, z_2) G(z_1, z_1, z_2, z_2) + p_2 \hat{A}(z_1, z_2, z_2, z_2) G(z_1, z_2, z_2, z_2) \\ + p_3 \hat{A}(z_1, z_1, z_2, z_2) G(z_1, z_1, z_2, z_2) + p_4 \hat{A}(z_1, z_1, z_2) G(z_1, z_1, z_2)$$

By substituting $z_3 = z_1, z_4 = z_2$ in Equation (11) we get

$$G(z_1, z_2, z_2, z_2) = p_1 \hat{A}(z_2, z_2, z_2, z_2) G(z_2, z_2, z_2, z_2) + p_2 \hat{A}(z_1, z_2, z_2, z_2) G(z_1, z_2, z_2, z_2) \\ + p_3 \hat{A}(z_1, z_2, z_2, z_2) G(z_1, z_2, z_2, z_2) + p_4 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2)$$

Finally, substituting $z_1 = z_2, z_3 = z_2, z_4 = z_2$ in Equation (11) gives

$$G(z_2, z_2, z_2, z_2) = p_1 \hat{A}(z_2, z_2, z_2, z_2) G(z_2, z_2, z_2, z_2) + p_2 \hat{A}(z_2, z_2, z_2, z_2) G(z_2, z_2, z_2, z_2) \\ + p_3 \hat{A}(z_2, z_2, z_2, z_2) G(z_2, z_2, z_2, z_2) + p_4 \hat{A}(z_2, z_2, z_2) G(z_2, z_2, z_2)$$

Step 5. Substituting $z_2 = z_1, z_3 = z_2, z_4 = z_3$ in Equation (11) leads to

$$G(z_1, z_1, z_2, z_3) = p_1 \hat{A}(z_1, z_1, z_2, z_3) G(z_1, z_1, z_2, z_3) + p_2 \hat{A}(z_1, z_2, z_2, z_2) G(z_1, z_2, z_2, z_2) \\ + p_3 \hat{A}(z_1, z_1, z_3, z_3) G(z_1, z_1, z_3, z_3) + p_4 \hat{A}(z_1, z_1, z_2) G(z_1, z_1, z_2)$$

Substituting $z_3 = z_2, z_4 = z_3$ in Equation (11) results in

$$G(z_1, z_2, z_2, z_3) = p_1 \hat{A}(z_2, z_2, z_2, z_3) G(z_2, z_2, z_2, z_3) + p_2 \hat{A}(z_1, z_2, z_2, z_3) G(z_1, z_2, z_2, z_3) \\ + p_3 \hat{A}(z_1, z_2, z_3, z_3) G(z_1, z_2, z_3, z_3) + p_4 \hat{A}(z_1, z_2, z_2) G(z_1, z_2, z_2)$$

Substituting $z_4 = z_3$ in Equation (11) we get

$$G(z_1, z_2, z_3, z_3) = p_1 \hat{A}(z_2, z_2, z_3, z_3) G(z_2, z_2, z_3, z_3) + p_2 \hat{A}(z_1, z_3, z_3, z_3) G(z_1, z_3, z_3, z_3) \\ + p_3 \hat{A}(z_1, z_2, z_3, z_3) G(z_1, z_2, z_3, z_3) + p_4 \hat{A}(z_1, z_2, z_3) G(z_1, z_2, z_3)$$

With the above substitutions, the sets are constructed.

References

1. Chernyak, V.Y.; Chertkov, M.; Goldberg, D.A.; Turitsyn, K.J. Non-Equilibrium Statistical Physics of Currents in Queuing Networks. *J. Stat. Phys.* **2010**, *140*, 819. [\[CrossRef\]](#)
2. Merhav, N.; Kafri, Y.J. Bose–Einstein condensation in large deviations with applications to information systems. *J. Stat. Mech.* **2010**, *2010*, P02011. [\[CrossRef\]](#)
3. Derrida, B. An exactly soluble non-equilibrium system: The asymmetric simple exclusion process. *Phys. Rep.* **1998**, *301*, 65. [\[CrossRef\]](#)
4. Golinelli, O.; Mallick, K. The asymmetric simple exclusion process: An integrable model for non-equilibrium statistical mechanics. *J. Phys. A: Math. Gen.* **2006**, *39*, 12679. [\[CrossRef\]](#)
5. Kaspi, H.; Kella, O.; Perry, D. Dam processes with state dependent batch sizes and intermittent production processes with state dependent rates. *Queueing Syst.* **1996**, *24*, 37–57. [\[CrossRef\]](#)
6. Jackson, J.R. Networks of Waiting Lines. *Oper. Res.* **1957**, *5*, 518–521. [\[CrossRef\]](#)
7. Jackson, J.R. Jobshop-Like Queueing Systems. *Manag. Sci.* **1963**, *10*, 131–142. [\[CrossRef\]](#)
8. Jackson, R.R.P. Random Queueing Processes with Phase-Type Service. *J. R. Stat. Soc. Ser. B (Methodol.)* **1956**, *18*, 129–132. [\[CrossRef\]](#)
9. Chen, H.; Yao, D.D. *Fundamentals of Queueing Networks*; Springer: Berlin/Heidelberg, Germany, 2001.
10. Bertsekas, D.P.; Gallager, R.G. *Data Networks*, 2nd ed.; Prentice Hall: Hoboken, NJ, USA, 1992.
11. Yechiali, U. Sequencing an N-Stage Process with Feedback. *Probab. Eng. Inf. Sci.* **1988**, *2*, 263–265. [\[CrossRef\]](#)

12. Blythe, R.A.; Evans, M.R. Nonequilibrium steady states of matrix-product form: A solver's guide. *J. Phys. A: Math. Theor.* **2007**, *40*, R333–R441. [\[CrossRef\]](#)
13. MacDonald, C.T.; Gibbs, J.H.; Pipkin, A.C. Kinetics of biopolymerization on nucleic acid templates. *Biopolymers* **1968**, *6*, 1–25. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Heckmann, K. Biomembranes: Passive Permeability of Cell Membranes. In *Proceedings of a Satellite Symposium*; Kreuzer, F., Slegers, J.F.G., Eds.; Plenum: New York, NY, USA, 1971.
15. Levitt, D.G. Dynamics of a Single-File Pore: Non-Fickian Behavior. *Phys. Rev. A* **1973**, *8*, 3050. [\[CrossRef\]](#)
16. Richards, P.M. Theory of one-dimensional hopping conductivity and diffusion. *Phys. Rev. B* **1977**, *16*, 1393. [\[CrossRef\]](#)
17. Wolf, D.E.; Schreckenberg, M.; Bachem, A. *Traffic and Granular Flow*; World Scientific: New York, NY, USA, 1996.
18. Shaw, L.B.; Zia, R.K.; Lee, K.H. Totally asymmetric exclusion process with extended objects: A model for protein synthesis. *Phys. Rev. E* **2003**, *68*, 021910. [\[CrossRef\]](#)
19. Reuveni, S.; Meilijson, I.; Kupiec, M.; Rupp, E.; Tuller, T. Genome-Scale Analysis of Translation Elongation with a Ribosome Flow Model. *PLoS Comput. Biol.* **2011**, *7*, e1002127. [\[CrossRef\]](#) [\[PubMed\]](#)
20. Halpin-Healy, T.; Zhang, Y.C. Kinetic roughening phenomena, stochastic growth, directed polymers and all that. *Asp. Multidiscip. Stat. Mech. Phys. Rep.* **1995**, *254*, 215–414.
21. Krug, J. Origins of scale invariance in growth processes. *Adv. Phys.* **1997**, *46*, 139–282. [\[CrossRef\]](#)
22. Bundschuh, R. Asymmetric exclusion process and extremal statistics of random sequences. *Phys. Rev. E* **2002**, *65*, 031911. [\[CrossRef\]](#)
23. Klumpp, S.; Lipowsky, R. Traffic of Molecular Motors Through Tube-Like Compartments. *J. Stat. Phys.* **2003**, *113*, 233–268. [\[CrossRef\]](#)
24. Burlatsky, S.F.; Oshanin, G.S.; Mogutov, A.V.; Moreau, M. Directed walk in a one-dimensional lattice gas. *Phys. Lett. A* **1992**, *166*, 230–234. [\[CrossRef\]](#)
25. Burlatsky, S.F.; Oshanin, G.; Moreau, M.; Reinhardt, W.P. Motion of a driven tracer particle in a one-dimensional symmetric lattice gas. *Phys. Rev. E* **1996**, *54*, 3165. [\[CrossRef\]](#) [\[PubMed\]](#)
26. Benichou, O.; Cazabat, A.M.; de Coninck, J.; Moreau, M.; Oshanin, G. Stokes Formula and Density Perturbances for Driven Tracer Diffusion in an Adsorbed Monolayer. *Phys. Rev. Lett.* **2000**, *84*, 511. [\[CrossRef\]](#) [\[PubMed\]](#)
27. Monasterio, C.M.; Oshanin, G. Bias- and bath-mediated pairing of particles driven through a quiescent medium. *Soft Matter* **2011**, *7*, 993–1000. [\[CrossRef\]](#)
28. Reuveni, S.; Eliazar, I.; Yechiali, U. Asymmetric Inclusion Process. *Phys. Rev. E* **2011**, *84*, 041101. [\[CrossRef\]](#)
29. Reuveni, S.; Eliazar, I.; Yechiali, U. Asymmetric Inclusion Process as a Showcase of Complexity. *Phys. Rev. Lett.* **2012**, *109*, 020603. [\[CrossRef\]](#)
30. Reuveni, S.; Eliazar, I.; Yechiali, U. Limit laws for the asymmetric inclusion process. *Phys. Rev. E* **2012**, *86*, 061133. [\[CrossRef\]](#)
31. Reuveni, S.; Eliazar, I.; Hirschberg, O.; Yechiali, U. Occupation Probabilities and Fluctuations in the Asymmetric Inclusion Process. *Phys. Rev. E* **2014**, *89*, 042109. [\[CrossRef\]](#) [\[PubMed\]](#)
32. Boxma, O.; Kella, O.; Yechiali, U. An ASIP Model with General Gate Opening Intervals. *Queueing Syst.* **2016**, *84*, 1–20. [\[CrossRef\]](#)
33. Boxma, O.; Kella, O.; Yechiali, U. Workload Distributions in ASIP Queueing Networks. *Queueing Syst.* **2021**, *97*, 81–100. [\[CrossRef\]](#)
34. Yeger, Y.; Yechiali, U. Performance Measures in a Generalized Asymmetric Simple Inclusion Process. *Mathematics* **2022**, *10*, 594. [\[CrossRef\]](#)