

Supplementary Materials

Table S1. Number of participants with missing data for each variable, expressed in absolute and relative frequencies, for the final sample composed by 307 children.

	<i>n</i> = 307
Parental characteristics	
Mother age at delivery (years)	0 (0.0%)
Mother origin country	0 (0.0%)
Mother level of education	0 (0.0%)
Mother social class	1 (0.3%)
Mother smoking during pregnancy	15 (4.9%)
Mother BMI (kg/m ²)	0 (0.0%)
Father BMI (kg/m ²)	11 (3.6%)
Parental cardiovascular antecedents	0 (0.0%)
Child characteristics	
Sex	0 (0.0%)
Age (years)	5 (1.6%)
Week of gestation at delivery	6 (2.0%)
Predominant breastfeeding duration (weeks)	37 (12.1%)
Birth weight (gr)	6 (2.0%)
At 4 years	
Mean daily energy intake (cals)	8 (2.6%)
Weekly out-of-school physical activity time (hours)	12 (3.9%)
Weight (kg)	6 (2.0%)
Height (cm)	6 (2.0%)
BMI (kg/m ²)	4 (1.3%)
Waist circumference (cm)	7 (2.3%)
Waist circumference/Height ratio	7 (2.3%)
Triponderal index (kg/m ³)	6 (2.0%)
Systolic blood pressure	7 (2.3%)
Diastolic blood pressure	7 (2.3%)
Mean arterial pressure	7 (2.3%)
Total cholesterol (mg/dL)	60 (19.5%)
cHDL (mg/dL)	60 (19.5%)
cLDL (mg/dL)	60 (19.5%)
Triglycerides (mg/dL)	60 (19.5%)
Atherogenic index	60 (19.5%)
Glucose (mg/dL)	59 (19.2%)
Insulin (μU/mL)	77 (25.1%)
HOMA index	77 (25.1%)
At 8 years	
Mean daily energy intake (cals)	21 (6.8%)
Weekly out-of-school physical activity time (hours)	19 (6.2%)
Weight (kg)	15 (4.9%)
Height (cm)	15 (4.9%)
BMI (kg/m ²)	15 (4.9%)
Waist circumference (cm)	15 (4.9%)
Waist circumference/Height ratio	15 (4.9%)
Triponderal index (kg/m ³)	15 (4.9%)
Systolic blood pressure	19 (6.2%)
Diastolic blood pressure	19 (6.2%)
Mean arterial pressure	19 (6.2%)

Total cholesterol (mg/dL)	61 (19.9%)
cHDL (mg/dL)	63 (20.5%)
cLDL (mg/dL)	63 (20.5%)
Triglycerides (mg/dL)	61 (19.9%)
Atherogenic index	63 (20.5%)
Glucose (mg/dL)	61 (19.9%)
Insulin (μU/mL)	71 (23.1%)
HOMA index	72 (23.5%)
